

Entrywise Inference for Causal Panel Data: A Simple and Instance-Optimal Approach

Yuling Yan^{*,†} Martin J. Wainwright^{*,†,‡}

Institute for Data, Systems, and Society^{*}
Department of Electrical Engineering and Computer Sciences[†]
Department of Mathematics[‡]
Massachusetts Institute of Technology
{yulingy,mjwain}@mit.edu

Abstract

In causal inference with panel data under staggered adoption, the goal is to estimate and derive confidence intervals for potential outcomes and treatment effects. We propose a computationally efficient procedure, involving only simple matrix algebra and singular value decomposition. We derive non-asymptotic bounds on the entrywise error, establishing its proximity to a suitably scaled Gaussian variable. Despite its simplicity, our procedure turns out to be instance-optimal, in that our theoretical scaling matches a local instance-wise lower bound derived via a Bayesian Cramér–Rao argument. Using our insights, we develop a data-driven procedure for constructing entrywise confidence intervals with pre-specified coverage guarantees. Our analysis is based on a general inferential toolbox for the SVD algorithm applied to the matrix denoising model, which might be of independent interest.

1 Introduction

Consider a study in which observations of units (e.g., individuals, companies, countries) are collected over time, and a subset of units are exposed to a “treatment” (e.g., drugs, regulations, or governmental interventions) beginning at some time. Given such data, how to estimate counterfactuals such as the treatment effect for a given unit-time pair? This problem and its variants, known as *causal inference with panel data*, find wide applications in economics, social sciences, and biomedical research (e.g., [HI08; LBNE08; HHY08; IR15]). In this paper, we assume that the treatment assignments follow the staggered adoption design [AI22; ST21], meaning that units may begin treatment at possibly different times, but that once initiated, the treatment is irreversible. We can collect the observed data for treated and untreated unit/period pairs into two different matrices, whose rows index units and columns index periods. Both matrices have complementary missing entries, since we cannot observe the counterfactual outcomes. See Figure 1 for an illustration of the induced missing pattern.

Broadly speaking, in the literature on causal inference, there are at least two main approaches for estimating causal effects with panel data: those based on unconfoundedness (e.g., [RR83; IR15]), and those based on synthetic controls (e.g., [AG03; ADH15; Aba21]). In order to understand how these methods work, suppose that we are interested in estimating the counterfactual outcome for unit i at period t . Approaches based on unconfoundedness seek to identify a subset of untreated units whose outcomes (before unit i was treated) are similar to those of unit i , and use their observed outcomes at period t to estimate unit i ’s missing outcome. On the other hand, approaches based on synthetic controls estimate unit i ’s counterfactual outcome at time t by using a weighted average

	time periods																	
	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	units	group 1
	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C		
	C	C	C	C	C	C	C	C	C	C	T	T	T	T	T	T		group 2
	C	C	C	C	C	C	C	C	C	C	T	T	T	T	T	T		
	C	C	C	C	C	C	C	T	T	T	T	T	T	T	T	T		
	C	C	C	C	C	C	C	T	T	T	T	T	T	T	T	T		group 3
	C	C	C	C	C	C	C	T	T	T	T	T	T	T	T	T		
	C	C	C	C	T	T	T	T	T	T	T	T	T	T	T	T		group 4
	C	C	C	C	T	T	T	T	T	T	T	T	T	T	T	T		
	stage 1				stage 2				stage 3				stage 4					

Figure 1. An illustration of a causal panel data with staggered adoption design. The labels C and T refer to “control” (i.e., untreated) and “treated” respectively.

of all untreated units’ observed outcomes at that time, where the weights are typically determined by solving a regression problem using a set of predictors that are unaffected by the treatment.

Athey et al. [Ath+21] observed that both the unconfoundedness and synthetic control methods can be related to the literature on low-rank matrix completion (e.g., [CR09; NW12]). Based on this connection, they proposed to impute the missing potential outcomes via the standard convex relaxation for low-rank matrix completion—namely, minimizing a least-squares objective with a nuclear norm regularization. When the potential outcome matrix is approximately low-rank, this approach can be equipped with attractive guarantees. Following this general avenue, the past few years have witnessed substantial progress in the development and analysis of matrix completion algorithms tailored for casual panel data (e.g., [BN21; CBN23; Aga+23; CY23]).

1.1 Our contributions

With this context, the main contribution of this paper is to propose a simple procedure, substantially less computationally intensive than nuclear norm relaxation, for inferring individual treatment effects in causal panel data. Moreover, we are able to show that our procedure—despite its simplicity—is unimprovable in a sharp instance-optimal sense. More precisely:

- **Entrywise guarantees:** We give non-asymptotic bounds and distributional characterizations of the error in estimating each matrix entry. Using the distributional characterization, we provide a data-driven procedure for computing confidence intervals for the unobserved entries with prescribed coverage.
- **Sharp instance-wise optimality:** We demonstrate that the length of our confidence intervals match the Bayesian Cramér-Rao lower bound associated with a genie-aided version of the problem (in which an oracle provides partial information about the unknown truth).
- **Computational simplicity:** Our algorithm only involves basic matrix operations and singular

value decompositions, making it computationally faster than standard convex relaxations (which typically involve solving semi-definite programs).

Our procedure and theory is built upon an inferential toolbox developed for the SVD algorithm in the context of low-rank matrix denoising, which might be of independent interest. See [Appendix F](#) for details.

A preview: [Figure 2](#) provides a high-level preview of some of the consequences of our results. In particular, let $\mathbf{M}^* \in \mathbb{R}^{N \times T}$ be the underlying matrix of outcomes—say for the control group.




Figure 2. Illustration of the asymptotic normality of the rescaled error $R_{i,t}$ from equation (1.1) for $(i, t) = (800, 800)$ (top panel) and $(i, t) = (800, 500)$ (bottom panels). Left panels show histograms based on 1000 independent trials with the density function of a standard $\mathcal{N}(0, 1)$ variate overlaid with a thin red line. Right panels show Q-Q plots of the empirical quantiles versus the standard normal quantiles.

Our methodological innovation is to propose an estimate $\widehat{\mathbf{M}}$, along with an analogous one for the unknown matrix of treated outcomes, whose computation involves only SVDs and other basic matrix operations. For each unobserved entry (i, t) , we define the rescaled error term

$$R_{i,t} = (\widehat{M}_{i,t} - M_{i,t}^*) / \sqrt{\widehat{\gamma}_{i,t}}, \quad (1.1)$$

where $\widehat{\gamma}_{i,t}$ is a data-dependent scaling factor. On the achievable side, we provide a non-asymptotic decomposition of the rescaled error $R_{i,t}$, a particular consequence of which is that it converges in

distribution to a standard $N(0, 1)$ variate; see [Theorems 1](#) and [3](#) for statements of these results. [Figure 2](#) provides an empirical demonstration of this predicted entrywise asymptotic normality, for two particular choices of the index pair (i, t) . Moreover, we prove that these estimation-theoretic and inferential guarantees are *optimal in a strong sense*: our variance estimates $\hat{\gamma}_{i,t}$ converge to a population quantity $\gamma_{i,t}^*$ that is defined via a non-asymptotic Bayesian Cramér–Rao lower bound; see [Theorem 2](#) and the discussion following [Theorem 3](#) for these claims.

1.2 Related work

So as to put our contributions in context, let us discuss some related lines of work. Low-rank matrix completion has been extensively studied under the random missing assumption, with convex relaxations based on the nuclear norm minimization being one of the most widely-studied methods. Initial investigations [[CR09](#); [RFP10](#); [CT10](#)] primarily focused on the noiseless scenario and examined the minimal sample size required for exact recovery. In the context of noisy matrix completion (where the observed entries are corrupted by random noise), the last decade has seen the establishment of optimal estimation guarantees for convex relaxation [[NW12](#); [Klo14](#); [Che+20](#)]. A more recent line of work [[Che+19](#); [XY21](#)] has focused on debiasing techniques for convex relaxation, showing how it is possible to construct confidence intervals for each entry of the unknown matrix. We refer to the reader to the survey [[CLC19](#)] for an overview of other non-convex approaches to matrix completion (e.g., [[BM03](#); [KMO10](#); [Sre04](#)]).

Athey et al. [[Ath+21](#)] pioneered the use of convex matrix completion for imputing potential outcomes in causal panel data. They provided Frobenius norm bounds on the error in estimating the full matrix; such a bound can be viewed as providing estimation guarantees for averaged treatment effects. More recently, Choi and Yuan [[CY23](#)] proposed a more refined approach that first divides the missing entries into smaller groups, and then applies the convex relaxation to each group. They derived entrywise error bounds that are sharper than the Frobenius norm error bound from the paper [[Ath+21](#)], and also established asymptotic normality for some statistics of interest. Agarwal et al. [[Aga+23](#)] proposed an algorithm based on synthetic nearest-neighbors, for which they established error bounds and asymptotic normality. There is also a line of work [[BN21](#); [CBN23](#)] on models satisfying certain factor constraints, to which we compare in more detail in [Section 4.3](#). Other work on related but distinct estimation problems for causal panel data include the papers [[FLP21](#); [LR23](#); [CKL23](#); [ASS20](#)].

Our algorithm involves spectral techniques, and consequently, our analysis of it has connections to past studies of spectral methods [[Abb+20](#); [Cai+21](#); [YCF21](#); [ZC23](#)]. Of particular relevance is our use of leave-one-out approaches [[EK15](#); [SCC17](#); [Ma+20](#); [Che+21b](#)] in giving a sharp characterization of the entrywise behavior; see the monograph [[Che+21a](#)] for more detail and background.

Notation: For a positive integer n , we adopt the shorthand $[n] := \{1, \dots, n\}$. For a matrix $\mathbf{A} \in \mathbb{R}^{n_1 \times n_2}$ and subsets $\mathcal{I} = \{i_1, i_2, \dots, i_I\} \subseteq [n_1]$ and $\mathcal{J} = \{j_1, j_2, \dots, j_J\} \subseteq [n_2]$, define $\mathbf{A}_{\mathcal{I}, \mathcal{J}} \in \mathbb{R}^{I \times J}$ to be the submatrix of \mathbf{A} comprising the intersection of its I rows indexed by \mathcal{I} and J columns indexed by \mathcal{J} , specifically $(\mathbf{A}_{\mathcal{I}, \mathcal{J}})_{k,l} = A_{i_k, j_l}$ for any $k \in [I]$ and $l \in [J]$. In addition, we abbreviate $\mathbf{A}_{\mathcal{I}, \cdot} := \mathbf{A}_{\mathcal{I}, [n_2]}$ and $\mathbf{A}_{\cdot, \mathcal{J}} := \mathbf{A}_{[n_1], \mathcal{J}}$. If a subset $\mathcal{I} = \{i\}$ is a singleton, we may directly use the index i to replace \mathcal{I} for simplicity. We use $\|\mathbf{A}\|$, $\|\mathbf{A}\|_{\text{F}}$ and $\|\mathbf{A}\|_{\infty}$ (respectively) to denote the spectral norm, Frobenius norm and entrywise ℓ_{∞} norms of a matrix \mathbf{A} . For any symmetric matrices \mathbf{A} and $\mathbf{B} \in \mathbb{R}^{n \times n}$, the relation $\mathbf{A} \succeq \mathbf{B}$ (resp. $\mathbf{A} \preceq \mathbf{B}$) means that $\mathbf{A} - \mathbf{B}$ (resp. $\mathbf{B} - \mathbf{A}$) is positive semidefinite. For any invertible matrix $\mathbf{H} \in \mathbb{R}^{r \times r}$ with SVD $\mathbf{U}\Sigma\mathbf{V}^{\top}$, define its sign

matrix $\text{sgn}(\mathbf{H}) := \mathbf{U}\mathbf{V}^\top$. We also define the function $\text{svds}(\mathbf{A}, r)$ to output the truncated rank- r SVD $(\mathbf{U}, \mathbf{\Sigma}, \mathbf{V})$ of \mathbf{A} , where the columns of $\mathbf{U} \in \mathbb{R}^{n_1 \times r}$ and $\mathbf{V} \in \mathbb{R}^{n_2 \times r}$ are the top- r left/right singular vectors, and $\mathbf{\Sigma} \in \mathbb{R}^{r \times r}$ is a diagonal matrix containing the top- r singular values.

2 Problem set-up

We now turn to a more precise description of estimation and inference based on causal panel data.

2.1 Basic observation model

We consider a panel data setting in which there are N units over T periods. Associated with each unit $i \in [N]$ at each time period $t \in [T]$ are the two potential outcomes $Y_{i,t}(0)$ and $Y_{i,t}(1)$, as well as a binary variable $\omega_{i,t} \in \{0, 1\}$. This binary variable indicates whether or not treatment has occurred; in particular, we have $\omega_{i,t} = 1$ whenever unit i is exposed to a treatment at time t , and $\omega_{i,t} = 0$ otherwise. With this set-up, we observe the collection of pairs $(\omega_{i,t}, Y_{i,t})$ where $Y_{i,t} = Y_{i,t}(\omega_{i,t})$ for all units $i \in [N]$ and times $t \in [T]$. Given this collection of observations, our goal is to estimate the treatment effect $\tau_{i,t} := Y_{i,t}(1) - Y_{i,t}(0)$ for each (i, t) pairs.

Of course, this problem is ill-posed in general, so that additional assumptions are required. The analysis of this paper is based on the following conditions:

- **Staggered adoption design:** The units may differ in the time they are first exposed to the treatment, but the treatment is irreversible (e.g., [AI22; Ath+21; ST21]).
- **Low-rank structure:** The potential outcome matrices have ranks much smaller than $\min\{N, T\}$.

2.2 Formulation as low-rank matrix completion

We now reformulate this problem more succinctly as an instance of low-rank matrix completion. For each of the two potential outcomes (treated versus untreated), we can introduce an $N \times T$ matrix that is populated by the outcomes—that is, matrices $\mathbf{M}^* \in \mathbb{R}^{N \times T}$ and $\mathbf{N}^* \in \mathbb{R}^{N \times T}$ with entries

$$M_{i,t}^* = Y_{i,t}(0) \quad \text{and} \quad N_{i,t}^* = Y_{i,t}(1).$$

We assume that \mathbf{M}^* and \mathbf{N}^* have ranks r and r' respectively, with both r and r' much smaller than $\min\{N, T\}$. Letting $\Omega \subseteq [N] \times [T]$ be an index subset where $(i, t) \in \Omega$ indicates that unit i is treated at time t , we observe the matrices

$$\mathbf{M} = \mathcal{P}_{\Omega^c}(\mathbf{M}^* + \mathbf{E}) \quad \text{and} \quad \mathbf{N} = \mathcal{P}_{\Omega}(\mathbf{N}^* + \mathbf{E}), \quad (2.1)$$

where $\mathcal{P}_{\Omega}(\mathbf{X})$ denotes the Euclidean projection of a matrix \mathbf{X} onto the subspace of matrices supported on Ω , and \mathbf{E} is a random noise matrix with i.i.d. $\mathcal{N}(0, \omega^2/(NT))$ entries. Our scaling of the noise variance ensures that \mathbf{E} has Frobenius norm close to one with high probability, and facilitates interpretation of results in the sequel.

Under the staggered adoption design, we can sort the N units according to the time t_i at which they were first exposed to the treatment; if unit i is never exposed to the treatment, then $t_i = +\infty$. With this sorting, we have $t_1 \geq t_2 \geq \dots \geq t_N$, and we are guaranteed the existence of an integer




Figure 3. An illustration of the the indicator matrix Ω , the noise matrix E , and the two potential outcome matrices M^* and N^* under staggered adoption design. Here $\mathbf{1}$ (resp. $\mathbf{0}$) denotes a matrix whose entries are all one (resp. zero). The light (resp. dark) gray blocks are the potential outcomes and noises associated with the untreated (resp. treated) unit/period pairs.

$k \geq 1$ and partitions N_1, \dots, N_k and T_1, \dots, T_k with $\sum_{i=1}^k N_i = N$ and $\sum_{j=1}^k T_j = T$ such that the Boolean matrix

$$\Omega \in \{0, 1\}^{N \times T} \quad \text{with entries } \Omega_{i,t} = \mathbb{1}_{(i,t) \in \Omega}, \quad (2.2)$$

associated with the subset Ω has the block structure as in the top left panel of Figure 3. We also partition M^* , N^* and E according to the pattern in Ω to get the submatrices $M_{i,j}^*, N_{i,j}^*, E_{i,j} \in \mathbb{R}^{N_i \times T_j}$, as shown in Figure 3. For each $i \in [k]$ and $1 \leq j \leq k - i$ define $M_{i,j} = M_{i,j}^* + E_{i,j}$, and for each $i \in [k]$ and $k - i < j \leq k$ define $N_{i,j} = N_{i,j}^* + E_{i,j}$.

2.3 Estimation and confidence intervals

Based on the observations M and N , our goal is to estimate and infer (e.g. constructing confidence intervals) the entries of $M^* - N^*$. Since we do not assume any joint structure on the pair M^* and N^* , it is equivalent to estimate each of M^* and N^* . Moreover, since the first N_1 units are never exposed to the treatment, it is in general impossible to estimate and infer the counterfactual outcomes $N_{1,j}^*$ for $j = 1, \dots, k$. Indeed, the first N_1 rows of N^* are unidentifiable, since they could be any linear combination of the other rows. For the same reason, we cannot estimate the counterfactual outcomes $N_{i,1}^*$ for $i = 1, \dots, k$. Therefore, our goal is to design an algorithm that

provides point estimates, along with confidence intervals, for the pairs $M_{i,t}^*$, $N_{i,t}^*$, as well as the treatment effect

$$\tau_{i,t}^* := N_{i,t}^* - M_{i,t}^*, \quad (2.3)$$

for any pair of indices $i \in \{N_1 + 1, N_1 + 2, \dots, N\}$ and $t \in \{T_1 + 1, T_1 + 2, \dots, T\}$.

In fact, it is sufficient to study estimation and inference for the missing blocks of \mathbf{M} . This is because the missing blocks of \mathbf{N} can be estimated in the same way if we rotate \mathbf{N} by 180 degrees, and the observed entries in \mathbf{M} (respectively \mathbf{N}) already provide reasonable estimates for the untreated (respectively treated) outcomes in \mathbf{M}^* (respectively \mathbf{N}^*). Although it is possible to denoise the observed outcomes by exploiting the low-rank structure¹, so as to bring focus to missing data aspect of the problem—which is the more challenging in nature—we focus on estimating and constructing confidence intervals for the unobserved entries in \mathbf{M} , and by analogy, the unobserved entries of \mathbf{N} as well.

2.4 A simple four-block structure

Our estimator for the general case can be understood most easily by understanding how it applies to a special four-block structure. More precisely, suppose that a subset of N_1 units *never* receive the treatment, while the other $N_2 = N - N_1$ units are exposed to the irreversible treatment at time $T_1 + 1$. Then our general observation model can be written in the four-block form

$$\mathbf{M}^* = \begin{bmatrix} \mathbf{M}_a^* & \mathbf{M}_b^* \\ \mathbf{M}_c^* & \mathbf{M}_d^* \end{bmatrix}, \quad \mathbf{N}^* = \begin{bmatrix} \mathbf{N}_a^* & \mathbf{N}_b^* \\ \mathbf{N}_c^* & \mathbf{N}_d^* \end{bmatrix}, \quad \text{and} \quad \mathbf{E} = \begin{bmatrix} \mathbf{E}_a & \mathbf{E}_b \\ \mathbf{E}_c & \mathbf{E}_d \end{bmatrix}. \quad (2.4)$$

Here the subscript a denotes matrices of size N_1 by T_1 ; the subscript b denotes matrices of size N_1 by $T - T_1$; and so on for the subscripts c and d . As shown in the sequel, it suffices to understand how to perform estimation and inference in this scenario because our algorithm for the general staggered design first reduces the problem to a simpler problem with four-block structure. In view of the discussion at the end of the previous section, our goal is to estimate and construct confidence intervals for the entries of \mathbf{M}_d^* , since they correspond to the counterfactual outcomes associated with the control group.

3 Estimation algorithms

In this section, we first design an algorithm that is applicable to the simpler four-block model (2.4). This algorithm is relatively easy to describe, and is the basic building block for our algorithm that applies to the general staggered design.

3.1 Algorithm for the four-block design

We first define some necessary notation for the four-block model. We begin by writing our observation matrix \mathbf{M} in the form

$$\mathbf{M} = \begin{bmatrix} \mathbf{M}_a & \mathbf{M}_b \\ \mathbf{M}_c & ? \end{bmatrix} = \begin{bmatrix} \mathbf{M}_a^* + \mathbf{E}_a & \mathbf{M}_b^* + \mathbf{E}_b \\ \mathbf{M}_c^* + \mathbf{E}_c & ? \end{bmatrix}.$$

¹This can be done by applying the truncated SVD algorithm to a submatrix of \mathbf{M} or \mathbf{N} that contains this observed outcome. See our inferential toolbox for the SVD algorithm in [Appendix F](#) for details.

Algorithm 1: Estimating counterfactual outcomes: four-block design

- 1 **Input:** Data matrix M , rank r
 /* Step 1: Subspace Estimation */
 - 2 Compute the truncated rank- r SVD $(U_{\text{left}}, \Sigma_{\text{left}}, V_{\text{left}})$ of M_{left} .
 - 3 Partition U_{left} into two submatrices U_1 and U_2 , where $U_1 \in \mathbb{R}^{N_1 \times r}$ consists of its top N_1 rows and $U_2 \in \mathbb{R}^{N_2 \times r}$ consists of its bottom N_2 rows.
 /* Step 2: Matrix Denoising */
 - 4 Compute the truncated rank- r SVD $(U_{\text{upper}}, \Sigma_{\text{upper}}, V_{\text{upper}})$ of M_{upper}
 - 5 Partition V_{upper} into two submatrices V_1 and V_2 , where $V_1 \in \mathbb{R}^{T_1 \times r}$ consists of its top T_1 rows and $V_2 \in \mathbb{R}^{T_2 \times r}$ consists of its bottom T_2 rows.
 - 6 Compute the estimate $\widehat{M}_b = U_{\text{upper}} \Sigma_{\text{upper}} V_2^\top$ of M_b^* .
 /* Step 3: Linear Regression */
 - 7 Compute the matrix $\widehat{M}_d = U_2 (U_1^\top U_1)^{-1} U_1^\top \widehat{M}_b$.
 - 8 **Output:** \widehat{M}_d as estimate of M_d^*
-

The unknown matrix M^* has a (compact) singular value decomposition, which we write in the form $M^* = U^* \Sigma^* V^{*\top}$, where $U^* \in \mathbb{R}^{N \times r}$ and $V^* \in \mathbb{R}^{T \times r}$ have orthonormal columns, and $\Sigma^* = \text{diag}(\sigma_1^*, \dots, \sigma_r^*)$ is a diagonal matrix consisting of the ordered singular values $\sigma_1^* \geq \sigma_2^* \geq \dots \geq \sigma_r^*$. We partition the matrices of singular vectors $U^* \in \mathbb{R}^{N \times r}$ and $V^* \in \mathbb{R}^{T \times r}$ into two blocks as

$$U^* = \begin{bmatrix} U_1^* \in \mathbb{R}^{N_1 \times r} \\ U_2^* \in \mathbb{R}^{N_2 \times r} \end{bmatrix} \quad \text{and} \quad V^* = \begin{bmatrix} V_1^* \in \mathbb{R}^{T_1 \times r} \\ V_2^* \in \mathbb{R}^{T_2 \times r} \end{bmatrix}. \quad (3.1)$$

It is also convenient to define

$$M_{\text{left}} = \begin{bmatrix} M_a \\ M_c \end{bmatrix}, \quad M_{\text{left}}^* = \begin{bmatrix} M_a^* \\ M_c^* \end{bmatrix}, \quad M_{\text{upper}} = [M_a \ M_b], \quad M_{\text{upper}}^* = [M_a^* \ M_b^*].$$

To motivate our algorithm, consider the ideal setting where there is no noise (i.e. $E = \mathbf{0}$). Given that M_a^* is rank- r , we can first compute the matrix U^* (up to a rotation) by using the SVD of the left submatrix M_{left}^* , and then recover M_d^* by

$$M_d^* = U_2^* (U_1^{*\top} U_1^*)^{-1} U_1^{*\top} M_b^*.$$

The idea behind the above formula is that, each column of M^* can be written uniquely as a linear combination of the columns of U^* . When U_1^* has full rank, we can find the coefficients by regressing each column of M_b^* over the columns of U_1^* . These coefficients in turn allow reconstruction of M_d^* using the columns of U_2^* .

In the presence of random noise, the naïve procedure needs to be adjusted. Doing so leads to the following three-step procedure:

- **(Subspace estimation)** The goal of this step is to estimate U^* . We compute the truncated rank- r SVD of M_{left} as

$$(U_{\text{left}}, \Sigma_{\text{left}}, V_{\text{left}}) \longleftarrow \text{svds}(M_{\text{left}}, r),$$

and use the top- r column subspace U_{left} as an estimate of U^* up to rotations. We partition U_{left} into two submatrices U_1 and U_2 , where $U_1 \in \mathbb{R}^{N_1 \times r}$ consists of its top N_1 rows and $U_2 \in \mathbb{R}^{N_2 \times r}$ consists of its bottom N_2 rows.

	T_1	T_2	T_3	T_4	T_5	T_6
N_1	$M_{1,1}$	$M_{1,2}$	$M_{1,3}$	$M_{1,4}$	$M_{1,5}$	$M_{1,6}$
N_2	$M_{2,1}$	$M_{2,2}$	$M_{2,3}$	$M_{2,4}$	$M_{2,5}$	
N_3	$M_{3,1}$	$M_{3,2}$	$M_{3,3}$	$M_{3,4}$		
N_4	$M_{4,1}$	$M_{4,2}$	$M_{4,3}$			
N_5	$M_{5,1}$	$M_{5,2}$?		
N_6	$M_{6,1}$					

	T_1	T_2	T_3	T_4	T_5	T_6
N_1	$M_{1,1}$	$M_{1,2}$	$M_{1,3}$	$M_{1,4}$	$M_{1,5}$	$M_{1,6}$
N_2	$M_{2,1}$	$M_{2,2}$	$M_{2,3}$	$M_{2,4}$	$M_{2,5}$	
N_3	$M_{3,1}$	$M_{3,2}$	$M_{3,3}$	$M_{3,4}$		
N_4	$M_{4,1}$	$M_{4,2}$	$M_{4,3}$			
N_5	$M_{5,1}$	$M_{5,2}$				
N_6	$M_{6,1}$?

(a)
(b)

Figure 4. Two examples on how to construct a four-block data matrix $\mathbf{M}^{(i_0, j_0)}$ to estimate \mathbf{M}_{i_0, j_0}^* . We consider $k = 6$, and the left and right panel correspond to $(i_0, j_0) = (5, 4)$ and $(6, 6)$ respectively. We use the question mark to denote the unobserved block that we want to estimate, and we use the bold line to single out the corresponding four-block design. For each observed block, we use different colors to distinguish their roles: red, blue and yellow blocks constitute \mathbf{M}_a , \mathbf{M}_b and \mathbf{M}_c (cf. (3.2)), while gray blocks are those unimportant data that we discard while estimating \mathbf{M}_{i_0, j_0}^* .

- **(Matrix denoising)** Then we need an estimate of \mathbf{M}_b^* . The most natural idea is to use \mathbf{M}_b , which is an unbiased estimator for \mathbf{M}_b^* . However, due to the noise in the observation, this naive estimate will incur much larger estimation error, leading to suboptimal statistical efficiency. Instead we compute the truncated rank- r SVD of $\mathbf{M}_{\text{upper}}$, given by

$$(\mathbf{U}_{\text{upper}}, \mathbf{\Sigma}_{\text{upper}}, \mathbf{V}_{\text{upper}}) \leftarrow \text{svds}(\mathbf{M}_{\text{upper}}, r),$$

We partition $\mathbf{V}_{\text{upper}}$ into two submatrices \mathbf{V}_1 and \mathbf{V}_2 , where $\mathbf{V}_1 \in \mathbb{R}^{T_1 \times r}$ consists of its top T_1 rows and $\mathbf{V}_2 \in \mathbb{R}^{T_2 \times r}$ consists of its bottom T_2 rows. We use $\widehat{\mathbf{M}}_b = \mathbf{U}_{\text{upper}} \mathbf{\Sigma}_{\text{upper}} \mathbf{V}_2^\top$ as an estimate of \mathbf{M}_b^* .

- **(Linear regression)** For each $1 \leq t \leq T_2$, we use the OLS solution to estimate the coefficient of the linear combination of the t -th column of \mathbf{M}_b^* with respect to the basis \mathbf{U}_1^*

$$\widehat{\beta}_t = \arg \min_{\beta \in \mathbb{R}^r} \|(\widehat{\mathbf{M}}_b)_{\cdot, t} - \mathbf{U}_1 \beta\|_2^2 = (\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top (\widehat{\mathbf{M}}_b)_{\cdot, t} \quad \text{for all } 1 \leq t \leq T_2.$$

This allows us to estimate the t -th column of \mathbf{M}_d^* by $\mathbf{U}_2 \widehat{\beta}_t$. In matrix form, our final estimate for \mathbf{M}_d^* is $\widehat{\mathbf{M}}_d = \mathbf{U}_2 (\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top \widehat{\mathbf{M}}_b$.

This procedure is summarized in [Algorithm 1](#). As we will see in the following sections, under mild assumptions the estimate $\widehat{\mathbf{M}}_d$ achieves full statistical efficiency for estimating \mathbf{M}_d^* (even including the preconstant).

3.2 Algorithm for the staggered adoption design

We move on to discuss the general staggered adoption design. At a high level, our procedure is based on reducing to the four-block case. Concretely, for any block (i_0, j_0) in [Figure 3](#), we define an associated four-block estimation problem. The solution to this problem—which can be

Algorithm 2: Estimating counterfactual outcomes: staggered adoption design

- 1 **Input:** Data matrix \mathbf{M} , Boolean matrix $\mathbf{\Omega}$, rank r .
 - 2 Extract the dimension information $\{N_i\}_{1 \leq i \leq k}$ and $\{T_j\}_{1 \leq j \leq k}$ from $\mathbf{\Omega}$.
 - 3 **for** $i_0 = 1$ **to** k **do**
 - 4 **for** $j_0 = k + 1 - i_0$ **to** k **do**
 - 5 Construct the data matrix $\mathbf{M}^{(i_0, j_0)}$ via equation (3.2).
 - 6 Call [Algorithm 1](#) with input $\mathbf{M}^{(i_0, j_0)}$ and rank r to compute an estimate $\widehat{\mathbf{M}}_d$ of its bottom left block.
 - 7 Extract the submatrix $\widehat{\mathbf{M}}_{i_0, j_0}$ from $\widehat{\mathbf{M}}_d$ corresponding to \mathbf{M}_{i_0, j_0}^* .
 - 8 **Output:** $\widehat{\mathbf{M}}_{i_0, j_0}$ as an estimate of \mathbf{M}_{i_0, j_0}^* for each (i_0, j_0)
-

obtained using the algorithm from the preceding section—allows us to the block \mathbf{M}_{i_0, j_0}^* . We obtain the four-block problem by removing a subset of data so as to simplify the observation structure; importantly, our theory to be described in the next section shows that the left-out data is ultimately not of significant statistical utility.

Let us now describe how to estimate a block \mathbf{M}_{i_0, j_0}^* , for some index (i_0, j_0) such that $j_0 > k + 1 - i_0$, so that it is not observed. Defining the indices $k_1 = k + 1 - j_0$ and $k_2 = k + 1 - i_0$, we construct a four-block data matrix $\mathbf{M}^{(i_0, j_0)}$ as follows

$$\mathbf{M}^{(i_0, j_0)} = \begin{bmatrix} \mathbf{M}_a & \mathbf{M}_b \\ \mathbf{M}_c & ? \end{bmatrix}, \quad \text{where} \quad \begin{cases} \mathbf{M}_a = [\mathbf{M}_{i,j}]_{1 \leq i \leq k_1, 1 \leq j \leq k_2}, \\ \mathbf{M}_b = [\mathbf{M}_{i,j}]_{1 \leq i \leq k_1, k_2 < j \leq j_0}, \\ \mathbf{M}_c = [\mathbf{M}_{i,j}]_{k_1 < i \leq i_0, 1 \leq j \leq k_2}. \end{cases} \quad (3.2)$$

Next we call [Algorithm 1](#) with input $\mathbf{M}^{(i_0, j_0)}$ to estimate its bottom-right block. Although the output provides estimation jointly for all blocks $\mathbf{M}_{i,j}^*$ where $k_1 < i \leq i_0$ and $k_2 < j \leq j_0$, we only use it to estimate the block \mathbf{M}_{i_0, j_0}^* . (For other blocks, this estimate may not be statistically efficient.) [Figure 4](#) and its caption provide two examples to help illustrate how we construct the four-block data matrix for estimating each unobserved block. The complete estimation procedure is summarized in [Algorithm 2](#).

As noted above, we show in the sequel that this algorithm achieves full statistical efficiency (including the constant pre-factors). Consequently, the data discarded while estimating each block are indeed unimportant for estimating that block, due to the lack of increase in statistical error.

4 Main results

We are now positioned to present our main theoretical guarantees for the algorithms described in [Section 3](#). We begin with theory for the four-block design in [Section 4.1](#), before extending it to the general case in [Section 4.2](#).

4.1 Theory for four-block design

As noted previously, low-rank matrix recovery is an under-determined problem, and certain regularity conditions are required to provide theoretical guarantees. We begin by stating the conditions used in our analysis; see the discussion in [Section 4.1.3](#) for some interpretation and intuition.

Sub-block conditioning: Our first set of conditions involve the top sub-blocks $\mathbf{U}_1^* \in \mathbb{R}^{N_1 \times r}$ and $\mathbf{V}_1^* \in \mathbb{R}^{T_1 \times r}$ of \mathbf{U}^* and \mathbf{V}^* , respectively. In particular, we assume that there are constants $0 < c_\ell \leq c_u < \infty$ such that

$$c_\ell \frac{N_1}{N} \mathbf{I}_r \preceq \mathbf{U}_1^{*\top} \mathbf{U}_1^* \preceq c_u \frac{N_1}{N} \mathbf{I}_r, \quad \text{and} \quad c_\ell \frac{T_1}{T} \mathbf{I}_r \preceq \mathbf{V}_1^{*\top} \mathbf{V}_1^* \preceq c_u \frac{T_1}{T} \mathbf{I}_r. \quad (4.1)$$

Noise level: Throughout, so as to avoid degenerate corner cases, we assume that $\min\{N_1, T_1\} \geq c \log(N+T)$. In addition, some of our results require an upper bound on the noise level, as measured by the ratio ω/σ_r^* . In particular, we define the noise functional

$$\rho_{N,T} := \frac{\omega}{\sigma_r^*} \frac{1}{\sqrt{\min\{N_1, T_1\}}} \quad (4.2)$$

For a given target error level $\delta > 0$, we require that $\min\{N_1, T_1\}$ is sufficiently large so as to ensure that

$$\rho_{N,T} \sqrt{r + \log(N+T)} \leq c_{\text{noise}} \delta \quad (4.3)$$

for sufficiently small constant $c_{\text{noise}} > 0$.

Local incoherence: For each $i \in [N]$ and $t \in [T]$, we define the local incoherence parameters

$$\mu_i := \sqrt{\frac{N}{r}} \|\mathbf{U}_{i,\cdot}^*\|_2 \quad \text{and} \quad \nu_t = \sqrt{\frac{T}{r}} \|\mathbf{V}_{t,\cdot}^*\|_2, \quad (4.4a)$$

along with the noise levels $\rho_N := \frac{\sigma}{\sigma_r^*} \sqrt{\frac{1}{N_1}}$ and $\rho_T := \frac{\sigma}{\sigma_r^*} \sqrt{\frac{1}{T_1}}$. For any given target error level $\delta > 0$, in order to provide recovery guarantees for entry (i, t) , we require that

$$\max\{\mu_i \rho_T, \nu_t \rho_N\} \sqrt{r} \leq c_{\text{inc}} \delta \quad \text{and} \quad \min\left\{\frac{\mu_i}{\sqrt{N_1}}, \frac{\nu_t}{\sqrt{T_1}}\right\} \sqrt{r \log(N+T)} \leq c_{\text{inc}} \delta \quad (4.4b)$$

for some sufficiently small constant $c_{\text{inc}} > 0$. For proving proximity to a Gaussian variable (as stated in the main text), we also require that

$$\min\left\{\frac{\rho_T}{\mu_i}, \frac{\rho_N}{\nu_t}\right\} \left\{\sqrt{\log(N+T)} + \frac{\log(N+T)}{\sqrt{r}}\right\} \leq c_{\text{inc}} \delta. \quad (4.4c)$$

We refer the reader to [Lemma 8](#) for some analysis not requiring this last condition.

4.1.1 An achievable result and its consequences

With these conditions in hand, we are now ready to state an achievable result for our estimator. It provides a non-asymptotic bound, for any unobserved index (i, t) , on the rescaled error of our estimator in terms of a standard Gaussian variable $G_{i,t} \sim \mathcal{N}(0, 1)$.

Theorem 1 (Non-asymptotics and distributional theory). *Under the sub-matrix condition (4.1), consider some $\delta > 0$ for which the noise condition (4.3) and incoherence conditions (4.4) hold. Then with probability at least $1 - O((N+T)^{-10})$, [Algorithm 1](#) produces an estimate $\widehat{\mathbf{M}}_d$ that for any unobserved index (i, t) , we have*

$$\left| \frac{1}{\sqrt{\gamma_{i,t}^*}} (\widehat{\mathbf{M}}_d - \mathbf{M}_d^*)_{i,t} - G_{i,t} \right| \leq \delta, \quad (4.5a)$$

and the variance takes the form

$$\gamma_{i,t}^* = \frac{\omega^2}{NT} \left\{ \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_{i,\cdot}^{*\top} + \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{*\top} \right\}. \quad (4.5b)$$

Let us explore some consequences of [Theorem 1](#), beginning with its uses in constructing confidence intervals. Observe that the bound [\(4.5a\)](#) guarantees that the rescaled error is close to a standard Gaussian variable. The scaling factor $\gamma_{i,t}^*$, corresponding to the error variance, depends on unknown population quantities. However, it can be estimated accurately by standard plug-in approach, leading to data-driven construction of entrywise confidence intervals. In particular, we define

$$\widehat{\gamma}_{i,t} := \frac{\widehat{\omega}^2}{NT} \left\{ \mathbf{U}_{i,\cdot} (\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_{i,\cdot}^\top + \mathbf{V}_{t,\cdot} (\mathbf{V}_1^\top \mathbf{V}_1)^{-1} \mathbf{V}_{t,\cdot}^\top \right\} \quad \text{where} \quad \widehat{\omega}^2 = \frac{N}{N_1} \|\mathbf{M}_{\text{upper}} - \widehat{\mathbf{M}}_{\text{upper}}\|_{\text{F}}^2.$$

Here all of the matrices $\mathbf{U} = \mathbf{U}_{\text{left}}$, $\mathbf{V} = \mathbf{V}_{\text{left}}$ and $\widehat{\mathbf{M}}_{\text{upper}} = \mathbf{U}_{\text{upper}} \boldsymbol{\Sigma}_{\text{upper}} \mathbf{V}_{\text{upper}}^\top$ can be computed from [Algorithm 1](#). Letting Φ denote the CDF of standard Gaussian distribution, we then define the interval

$$\text{CI}_{i,t}^{1-\alpha} := [\widehat{M}_{i,t} \pm \Phi^{-1}(1 - \alpha/2) \sqrt{\widehat{\gamma}_{i,t}}], \quad (4.6a)$$

With this definition, we have the following guarantee:

Proposition 1. *In addition to the conditions of [Theorem 1](#), suppose that $\min\{N_1, T\} \geq \delta^{-2} r \log(N+T)$. Then the interval $\text{CI}_{i,t}^{1-\alpha}$ has the coverage guarantee*

$$\mathbb{P}\left(\text{CI}_{i,t}^{1-\alpha} \ni M_{i,t}^*\right) = 1 - \alpha + O(\delta + (N+T)^{-10}). \quad (4.6b)$$

See [Section 5.2](#) for the proof.

Second, while [Theorem 1](#) is stated in a way that facilitates its inferential use, it also allows us to derive entrywise error bounds for $\widehat{\mathbf{M}}_d$. If the conditions of [Theorem 1](#) hold with some δ decreasing to zero as $\min\{N_1, T_1\}$ grows, then it can be verified that

$$\mathbb{E}[(\widehat{M}_{i,t} - M_{i,t}^*)^2] = (1 + o(1)) \gamma_{i,t}^* \lesssim \frac{\omega^2}{N_1 T} \|\mathbf{U}_{i,\cdot}^*\|_2^2 + \frac{\omega^2}{N T_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2. \quad (4.7a)$$

To provide intuition for the convergence rate, suppose that the local incoherence parameters μ_i and ν_t are viewed as quantities of constant order; in this case, we have

$$\mathbb{E}[(\widehat{M}_{i,t} - M_{i,t}^*)^2] \lesssim \frac{\omega^2 r}{NT \min\{N_1, T_1\}} \asymp \frac{r}{NT} \rho_{N,T}^2. \quad (4.7b)$$

In fact, even without the incoherence conditions [\(4.4\)](#) and with a relaxation of the noise condition [\(4.3\)](#), our analysis still provides bounds on the estimation error; see [Lemma 1](#) for details.

4.1.2 Matching lower bound

Is it possible to improve the variance term (4.5b) in our achievable result? In this section, we provide a negative answer to this question by showing that it matches with a Bayesian Cramér–Rao lower bound.

We decompose the unknown matrix as $\mathbf{M}^* = \mathbf{X}^* \mathbf{Y}^{*\top}$, where $\mathbf{X}^* := \mathbf{U}^* (\boldsymbol{\Sigma}^*)^{1/2}$ and $\mathbf{Y}^* := \mathbf{V}^* (\boldsymbol{\Sigma}^*)^{1/2}$ are $N \times r$ and $T \times r$ matrices, respectively. Now suppose that there is a genie that reveals all rows of \mathbf{X}^* *except* for the i -th one $\mathbf{x}^* := \mathbf{X}_{i,\cdot}^*$, and also all rows of \mathbf{Y}^* *except* for its t -th one $\mathbf{y}^* := \mathbf{Y}_{t,\cdot}^*$. Given this information, the statistician is left with the following problem:

- (a) The only unknown parameters are the r -dimensional row vectors $\mathbf{x}^* := \mathbf{X}_{i,\cdot}^*$ and $\mathbf{y}^* := \mathbf{Y}_{t,\cdot}^*$, and the goal is to estimate their inner product $(\mathbf{M}_d^*)_{i,t} \equiv \langle \mathbf{x}^*, \mathbf{y}^* \rangle$.
- (b) The remaining observations of any use are the $N_1 + T_1$ linear measurements

$$\{\mathbf{M}_{i,s} = \langle \mathbf{x}^*, \mathbf{Y}_{s,\cdot}^* \rangle + E_{i,s}\}_{s=1}^{T_1} \quad \text{and} \quad \{\mathbf{M}_{k,t} = \langle \mathbf{X}_{k,\cdot}^*, \mathbf{y}^* \rangle + E_{k,t}\}_{k=1}^{N_1}. \quad (4.8)$$

All other observations are uninformative given the information revealed by the genie.

Thus, with the aid from the genie, the causal panel data model reduces to a linear regression model with dimension $2r$.

We use this genie-aided problem to compute lower bounds for the estimation task—both the Cramér–Rao lower bound as well as a local minimax risk. The local minimax risk at scale $\varepsilon > 0$ is given by

$$R_{\text{loc}}(\varepsilon) := \inf_{\widehat{M}_{i,t}} \sup_{\substack{\mathbf{x} \in \mathcal{B}_\infty(\mathbf{x}^*, \varepsilon) \\ \mathbf{y} \in \mathcal{B}_\infty(\mathbf{y}^*, \varepsilon)}} \mathbb{E} \left[\left(\widehat{M}_{i,t} - \langle \mathbf{x}, \mathbf{y} \rangle \right)^2 \right]. \quad (4.9)$$

Here the infimum is taken over all estimators of the scalar $\langle \mathbf{x}, \mathbf{y} \rangle$ based on samples from the model (4.8), except that the true parameters are \mathbf{x} and \mathbf{y} instead of \mathbf{x}^* and \mathbf{y}^* .

Theorem 2 (Optimality). *In the genie-aided setting, the Cramér–Rao lower bound for estimating $(\mathbf{M}_d^*)_{i,t}$ is given by $\gamma_{i,t}^*$. More generally, under the sub-block condition (4.1), the local minimax risk at scale $\varepsilon_{N,T} = \frac{\sqrt{\sigma_r^*}}{\sqrt{\min\{N,T\}}}$ is lower bounded by*

$$R_{\text{loc}}(\varepsilon_{N,T}) \geq \left(1 - \frac{c_u \pi^2}{c_\ell^2} \rho_{N,T}^2 \right) \gamma_{i,t}^*, \quad (4.10)$$

where $\rho_{N,T}$ is the noise level previously defined (4.2).

See Section 5.3 for the proof.

Under the noise assumption (4.3), we have $\rho_{N,T}^2 \leq \frac{c_{\text{noise}}^2 \delta^2}{r + \log(N+T)}$, whence $\frac{R_{\text{loc}}(\varepsilon_{N,T})}{\gamma_{i,t}^*} \geq 1 - O(\delta^2)$ as δ goes to zero. Thus, we see that our estimator (cf. Algorithm 1) matches the local minimax risk aided by the genie, including a sharp pre-factor. Thus, our procedure is optimal in a strong instance-dependent sense.

4.1.3 Interpretation of structural conditions

Let us interpret the structural conditions—namely, the sub-block and local incoherence conditions—that underlie Theorem 1.

Sub-block condition: Since $\mathbf{M}_a^* = \mathbf{U}_1^* \boldsymbol{\Sigma}^* \mathbf{V}_1^*$, the existence of the constant $c_\ell > 0$ ensures that \mathbf{M}_a^* has full rank. This condition is necessary, because otherwise it would be impossible to recover the missing block \mathbf{M}_d^* even in the noiseless setting.² Otherwise, we observe that \mathbf{U}_1^* is an $N_1 \times T_1$ sub-matrix of the orthonormal matrix $\mathbf{U}^* \in \mathbb{R}^{N \times T}$. Since \mathbf{U}^* has orthonormal columns, we expect that the typical size of each entry is $1/\sqrt{N}$. Thus, the typical squared norm of each column of the sub-matrix \mathbf{U}_1^* is of the order N_1/N , which accounts for the normalization in the condition (4.1) on \mathbf{U}_1^* . Similar comments apply to the condition on \mathbf{V}_1^* .

Local incoherence conditions: Since the Frobenius norm of \mathbf{U}^* and \mathbf{V}^* are both \sqrt{r} , the “typical size” of their rows (in ℓ_2 norm) are $\sqrt{r/N}$ and $\sqrt{r/T}$ respectively. We can see that μ_i (resp. ν_i) evaluates the ratio between $\|\mathbf{U}_{i,\cdot}^*\|_2$ (resp. $\|\mathbf{V}_{i,\cdot}^*\|_2$) and its typical size. This definition connects to the (global) incoherence condition that is widely assumed in the low-rank matrix completion literature [CR09; Che15; CLC19], where the (global) incoherence parameters $\mu := \max_{1 \leq i \leq N} \mu_i$ and $\nu := \max_{1 \leq t \leq T} \nu_t$ defined therein cannot be too large in order for the recovery to be possible. In contrast, our theory does not assume any global incoherence condition to hold. If our goal is to estimate and infer the counterfactual outcome or treatment effect of unit i at time t , we just need the local incoherence parameters μ_i and ν_t to be reasonably well-behaved.

Our constraints involving the local incoherence conditions (4.4b) are reasonably mild. When $\min\{N_1, T_1\} \gg r$ and the noise condition (4.3) is satisfied, we can allow both $\|\mathbf{U}_{i,\cdot}^*\|_2$ and $\|\mathbf{V}_{i,\cdot}^*\|_2$ to be much larger than their typical size. The other condition (4.4c) requires one of $\|\mathbf{U}_{i,\cdot}^*\|_2$ and $\|\mathbf{V}_{i,\cdot}^*\|_2$ to be not vanishingly small compared to their typical size. In fact, even when (4.4c) does not hold, we can still characterize the entrywise distribution of $\widehat{\mathbf{M}}_d - \mathbf{M}_d^*$, although it is not approximately Gaussian; we refer interested readers to Lemma 8 for this general result.

4.2 Theory for staggered adoption design

In this section, we extend our theoretical analysis to the general (non-four-block) setting to which Algorithm 2 applies. Without loss of generality, we focus on estimating and inferring entries of an unobserved submatrix \mathbf{M}_{i_0, j_0}^* . In Section 3, we described how to construct a data matrix $\mathbf{M}^{(i_0, j_0)}$ for estimating \mathbf{M}_{i_0, j_0}^* that admits the four-block design; see equation (3.2) for details. Figure 5 provides a simple example so as to help visualize the set-up and to facilitate later discussion.

At the left of Figure 5, we depict a panel data problem with units separated into $k = 6$ subsets, as indicated by the large blocks. The i -th subset has N_i units that all received the treatment at time $1 + \sum_{j=1}^i T_j$; recall that a time larger than T means that the unit never received the treatment. All the colored blocks are observed, while the uncolored blocks are missing. Suppose we want to estimate the sub-matrix \mathbf{M}_{i_0, j_0}^* indexed by the pair $(i_0, j_0) = (5, 4)$, as indicated by a the question mark in the figure. The data matrix $\mathbf{M}_{(i_0, j_0)}^*$ consists of the $i_0 = 5$ by $j_0 = 4$ block matrix on the top left corner of \mathbf{M} . The gray blocks, although observed, are discarded to enforce the four-block structure. We use three colors (red, blue and yellow) to show the sub-matrices that play the role of \mathbf{M}_a^* , \mathbf{M}_b^* and \mathbf{M}_c^* in the four-block model. The red component consists of the $k_1 = 3$ by $k_2 = 2$ block matrix on the top left corner; in general, these two integers satisfy the relations $k_1 = k + 1 - j_0$ and $k_2 = k + 1 - i_0$.

²For example, if the row rank of \mathbf{M}_a^* is less than r , then we may find a row of \mathbf{M}_b^* that is not in the subspace spanned by the rows of \mathbf{M}_a^* . In this case, the corresponding row of \mathbf{M}_d^* is unidentifiable.

Now we define the dimension parameters \bar{N}_1 , \bar{N}_2 , \bar{T}_1 and \bar{T}_2 with respect to the four-block structure, given by

$$\bar{N}_1 := \sum_{i=1}^{k_1} N_i, \quad \bar{N}_2 := \sum_{i=k_1+1}^{i_0} N_i, \quad \bar{T}_1 := \sum_{j=1}^{k_2} T_j, \quad \text{and} \quad \bar{T}_2 := \sum_{j=k_2+1}^{j_0} T_j,$$

We also define $\bar{N} := \bar{N}_1 + \bar{N}_2$ and $\bar{T} := \bar{T}_1 + \bar{T}_2$. Let $\mathbf{U}_1^* \in \mathbb{R}^{\bar{N}_1 \times r}$ and $\mathbf{U}_2^* \in \mathbb{R}^{\bar{N}_2 \times r}$ be submatrices of $\mathbf{U}^* \in \mathbb{R}^{\bar{N} \times r}$, consisting of its first \bar{N}_1 rows and the next \bar{N}_2 rows respectively. Similarly, let $\mathbf{V}_1^* \in \mathbb{R}^{\bar{T}_1 \times r}$ and $\mathbf{V}_2^* \in \mathbb{R}^{\bar{T}_2 \times r}$ be submatrices of $\mathbf{V}^* \in \mathbb{R}^{\bar{T} \times r}$, consisting of its first \bar{T}_1 rows and the next \bar{T}_2 rows respectively. These dimension parameters are also illustrated in [Figure 5](#).




Figure 5: An illustration of the dimension parameters and the partition used in [Section 4.2](#).

In analogy with the four-block design, we first introduce the conditions required for establishing theoretical guarantees in the general staggered adoption design.

Sub-block conditioning: We assume that there are constants $0 < c_\ell \leq c_u < \infty$ such that

$$c_\ell \frac{\bar{N}_1}{N} \mathbf{I}_r \preceq \mathbf{U}_1^{*\top} \mathbf{U}_1^* \preceq c_u \frac{\bar{N}_1}{N} \mathbf{I}_r, \quad c_\ell \frac{\bar{N}}{N} \mathbf{I}_r \preceq \mathbf{U}_1^{*\top} \mathbf{U}_1^* + \mathbf{U}_2^{*\top} \mathbf{U}_2^* \preceq c_u \frac{\bar{N}}{N} \mathbf{I}_r, \quad (4.11a)$$

$$c_\ell \frac{\bar{T}_1}{T} \mathbf{I}_r \preceq \mathbf{V}_1^{*\top} \mathbf{V}_1^* \preceq c_u \frac{\bar{T}_1}{T} \mathbf{I}_r, \quad c_\ell \frac{\bar{T}}{T} \mathbf{I}_r \preceq \mathbf{V}_1^{*\top} \mathbf{V}_1^* + \mathbf{V}_2^{*\top} \mathbf{V}_2^* \preceq c_u \frac{\bar{T}}{T} \mathbf{I}_r. \quad (4.11b)$$

Recall the sub-block condition (4.1) for the four-block design and the discussion in [Section 4.1.3](#), here we are assuming in addition that $\mathbf{U}_1^{*\top} \mathbf{U}_1^* + \mathbf{U}_2^{*\top} \mathbf{U}_2^*$ and $\mathbf{V}_1^{*\top} \mathbf{V}_1^* + \mathbf{V}_2^{*\top} \mathbf{V}_2^*$ do not deviate too much from their typical size. This additional requirement connects the spectrum of the ground truth of $\mathbf{M}^{(i_0, j_0)}$ (which is a submatrix of \mathbf{M}^*) with that of the full ground truth \mathbf{M}^* , which allows us to present cleaner results that depend on the spectrum of \mathbf{M}^* instead of its submatrices.

Noise level: The noise level in this general setting takes the following form:

$$\bar{\rho}_{N,T} := \max\{\bar{\rho}_N, \bar{\rho}_T\}, \quad \text{where} \quad \bar{\rho}_N := \frac{\omega}{\sigma_r^*} \sqrt{\frac{1}{\bar{N}_1}} \quad \text{and} \quad \bar{\rho}_T := \frac{\omega}{\sigma_r^*} \sqrt{\frac{1}{\bar{T}_1}}. \quad (4.12)$$

For any prescribed target error level $\delta > 0$, we assume that it is upper bounded by

$$\bar{\rho}_{N,T} \sqrt{r + \log(N+T)} \leq c_{\text{noise}} \delta, \quad (4.13)$$

for some sufficiently small constant $c_{\text{noise}} > 0$.

Local incoherence: For each $i \in [N]$ and $t \in [T]$, the local incoherence parameters μ_i and ν_t are defined as in equation (4.4). For any given target error level $\delta > 0$, we assume that there is some sufficiently small constant $c_{\text{inc}} > 0$ such that,

$$\max \{ \mu_i \bar{\rho}_T, \nu_t \bar{\rho}_N \} \sqrt{r} \leq c_{\text{inc}} \delta \quad \text{and} \quad \min \left\{ \frac{\mu_i}{\sqrt{N_1}}, \frac{\nu_t}{\sqrt{T_1}} \right\} \sqrt{r \log(N+T)} \leq c_{\text{inc}} \delta \quad (4.14a)$$

and that at least one of the following conditions holds

$$\min \left\{ \frac{\bar{\rho}_T}{\mu_i}, \frac{\bar{\rho}_N}{\nu_t} \right\} \left\{ \sqrt{\log(N+T)} + \frac{\log(N+T)}{\sqrt{r}} \right\} \leq c_{\text{inc}} \delta. \quad (4.14b)$$

With this set-up, we let us describe how our theory applies to this more general setting, in particular by showing proximity of the rescaled error to a standard normal $G_{i,t} \sim \mathcal{N}(0, 1)$.

Theorem 3 (Non-asymptotics and distribution: General setting). *For a given $\delta > 0$, suppose that the noise condition (4.13) and the incoherence conditions (4.14) hold. For any index (i, t) belonging to any unobserved submatrix \mathbf{M}_{i_0, j_0}^* , with probability at least $1 - O((N+T)^{-10})$, Algorithm 2 produces an estimate $\widehat{\mathbf{M}}_{i_0, j_0}$ such that*

$$\left| \frac{1}{\sqrt{\gamma_{i,t}^*}} (\widehat{\mathbf{M}}_{i_0, j_0} - \mathbf{M}_{i_0, j_0}^*)_{i,t} - G_{i,t} \right| \leq \delta, \quad (4.15a)$$

where the variance is given by

$$\gamma_{i,t}^* = \frac{\omega^2}{NT} \left\{ \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_i^{*\top} + \mathbf{V}_t^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_t^{*\top} \right\}. \quad (4.15b)$$

See Appendix E for the proof.

Observe that this result is the natural generalization of our previous achievable result (Theorem 1) for the simpler four-block setting. This correspondence makes sense, since our algorithm for the general case involves “reducing” the problem to instances of the simpler four-block case. Of course, one might wonder whether or not this reduction—which involves discarding some data that might be relevant—leads to a sub-optimal guarantee.

Interestingly, our lower bound—establishing the optimality of the variance $\gamma_{i,t}^*$, as stated in Theorem 2—also extends to this multi-block setting. (We do not state this lower bound formally here, but note that our proof is actually given for the multi-block setting; see Section 5.3 for details.) As a consequence, under the conditions of Theorem 3, there is *no appreciable loss* of accuracy in solving the problem by successive reduction to the four-block setting.

4.3 Comparison with previous work

Having given precise statements of our results, let us now provide a detailed comparison with related results established in past work [Ath+21; BN21; Aga+23; CY23]. To simplify discussion, we ignore any log factors in making these comparisons, and use \lesssim to reflect inequalities that ignore constants and such log factors. For making comparisons, we record here the estimation error bound that can be deduced from Theorem 3: for any entry (i, t) , the error is upper bounded as

$$|\widehat{M}_{i,t} - M_{i,t}^*| \lesssim \omega \sqrt{\frac{1}{N_1 T}} \|\mathbf{U}_{i,\cdot}^*\|_2 + \omega \sqrt{\frac{1}{N \bar{T}_1}} \|\mathbf{V}_{t,\cdot}^*\|_2 \quad (4.16)$$

with high probability.

- Athey et al. [Ath+21] proposed to use convex relaxation (nuclear norm minimization) to estimate the missing potential outcomes. They showed that with high probability

$$\frac{1}{\sqrt{NT}} \|\widehat{\mathbf{M}}^{\text{ABD}} - \mathbf{M}^*\|_{\text{F}} \lesssim \frac{N^{1/4} \|\mathbf{M}^*\|_{\infty}}{\sqrt{N_1}} + \frac{\omega}{N_1 T} \sqrt{\frac{r}{\min\{N, T\}}}. \quad (4.17)$$

This is a Frobenius norm bound, which measures the average error over the full matrix and does not provide entrywise guarantees like (4.16). In addition, their bound (4.17) does not vanish as the noise level σ goes to zero, so that even when there is no noise, it does not ensure exact recovery.

- In recent work, Choi and Yuan [CY23] built upon the paper [Ath+21], providing an algorithm with superior estimation guarantees. In particular, they suggested a more sophisticated and multi-round use of convex relaxation, based on dividing the missing entries into smaller groups, and then solving a nuclear-norm regularized problem for each of these groups. They showed that with high probability, their estimator $\widehat{\mathbf{M}}^{\text{CY}}$ satisfies the entrywise ℓ_{∞} -bound

$$\|\widehat{\mathbf{M}}^{\text{CY}} - \mathbf{M}^*\|_{\infty} \lesssim \frac{\omega}{\sqrt{NT}} \sqrt{\frac{\kappa^5 \mu^2 r^3}{\min\{N_1, \bar{T}_1\}}}. \quad (4.18)$$

Here $\kappa = \sigma_1^*/\sigma_r^*$ is the condition number of \mathbf{M}^* , and μ is the *global* incoherence parameter (i.e., the maximum over all the local parameters μ_i and ν_t defined in equation (4.4)). In comparison, our bound (4.16) is a localized improvement of their bound (4.18); in particular, it has better dependency on the rank r , and no dependence on the condition number. In addition, we also provide data-driven confidence intervals for each entry of \mathbf{M}^* with optimal width. It is also worth mentioning that our spectral algorithm is more computationally efficient than the SDP-based algorithm in [CY23].

- Agarwal et al. [Aga+23] proposed a procedure called “synthetic nearest neighbors” (SNN), for which they proved entrywise error bound and established asymptotic normality. Their algorithm works for a wider range of missing mechanism and is more general than our algorithm. However, their error rate is not optimal under the staggered adoption design. For example, if $N = T$ and there is only one missing entry, their estimation error scales as $N^{-1/4}$, which is slower than the $N^{-1/2}$ rate for our algorithm and convex relaxation.
- Another line of work [BN21; CBN23] is based on assuming a factor structure for the full panel data matrix, and applying spectral algorithms that share similar spirit with ours. In general, the

analyses in these papers impose stronger assumptions than those used in our analysis. First, they model the factors and factor loadings—roughly speaking, these correspond to the rows of \mathbf{U}^* and \mathbf{V}^* —as being random, and required to satisfy certain asymptotic moment conditions. Second, they require that the observation sizes satisfy the constraint $\min\{\bar{N}_1, \bar{T}_1\} \gg \sqrt{\max\{N, T\}}$, as well as certain eigengap conditions. Neither of these conditions are imposed in our analysis. Third, their underlying noise condition (used in the asymptotic analysis) is $\sqrt{\min\{\bar{N}_1, \bar{T}_1\}}$ times more stringent than our noise condition (4.3). On the other hand, we note that their set-up is general enough to handle weakly correlated noise, which is not covered by our framework.

5 Proofs

In this section, we provide the proofs of certain results, including our achievable guarantee in the four-block case (Theorem 1) in Section 5.1; our data-dependent procedure for computing confidence intervals (Proposition 1) in Section 5.2; and our lower bounds (Theorem 2) in Section 5.3.

In all cases, we defer the proofs of various intermediate results, of a more technical nature, to the appendices. In addition, we defer the proof of Theorem 3 to Appendix E.

5.1 Proof of Theorem 1

Throughout this and other proofs, we use the shorthand notation $\xi_{N,T} := \log(N + T)$.

5.1.1 A master decomposition

Our proof is based on a key decomposition, which we state as a separate lemma in its own right. It involves the matrix

$$\mathbf{Z} := \mathbf{U}^*(\mathbf{U}_1^{*\top}\mathbf{U}_1^*)^{-1}\mathbf{U}_1^{*\top}\mathbf{E}_b + \mathbf{E}_c\mathbf{V}_1^*(\mathbf{V}_1^{*\top}\mathbf{V}_1^*)^{-1}\mathbf{V}^{*\top} + \mathbf{E}_c\mathbf{V}_1^*(\mathbf{V}_1^{*\top}\mathbf{V}_1^*)^{-1}(\boldsymbol{\Sigma}^*)^{-1}(\mathbf{U}_1^{*\top}\mathbf{U}_1^*)^{-1}\mathbf{U}_1^{*\top}\mathbf{E}_b, \quad (5.1)$$

as well as the error terms

$$\begin{aligned} b_1(i, t) &= \frac{1}{\sqrt{N_1 T}} \left(\frac{\omega^3}{\sigma_r^{*2} N_1} + \frac{\omega^2}{\sigma_r^* \sqrt{T_1}} \right) \|\mathbf{U}_{i,\cdot}^*\|_2 \sqrt{r + \xi_{N,T}}, \\ b_2(i, t) &= \frac{1}{\sqrt{N T_1}} \left(\frac{\omega^3}{\sigma_r^{*2} N_1} + \frac{\omega^2}{\sigma_r^* \sqrt{N}} + \frac{\omega^2}{\sigma_r^* \sqrt{T_1}} \right) \|\mathbf{V}_{t,\cdot}^*\|_2 \sqrt{r + \xi_{N,T}}, \\ b_3(i, t) &= \left(\rho_{N,T} + \omega \sqrt{\frac{r}{N_1 T}} + \omega \sqrt{\frac{1}{N_1 T_1} \xi_{N,T}} \right) \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2, \\ b_4(i, t) &= \frac{1}{\sqrt{N T}} \frac{1}{\sqrt{N_1 T_1}} \left(\frac{\omega^3}{\sigma_r^{*2} \sqrt{N}} + \frac{\omega^3}{\sigma_r^{*2} \sqrt{T_1}} + \frac{\omega^4}{\sigma_r^{*3} N_1} \right) (r + \xi_{N,T}). \end{aligned}$$

Lemma 1 (Master decomposition). *Suppose that the sub-block condition (4.1) and the noise condition (4.3) hold. Then we can write*

$$\widehat{\mathbf{M}}_d - \mathbf{M}_d^* = \mathbf{Z} + \boldsymbol{\Delta}, \quad (5.2a)$$

for a perturbation matrix Δ whose entries satisfy the entrywise bound

$$|\Delta_{i,t}| \leq C_4 \sum_{k=1}^4 b_k(i,t) \quad \text{for all entries } (i,t) \quad (5.2b)$$

with probability at least $1 - O((N+T)^{-9})$.

See [Section 5.1.3](#) for the proof.

5.1.2 Main argument for [Theorem 1](#)

Armed with this result, we are now equipped to prove [Theorem 1](#). We begin with the additive decomposition $Z_{i,t} = \alpha_{i,t} + \beta_{i,t} + \lambda_{i,t}$, where

$$\alpha_{i,t} = (\mathbf{E}_c)_{i,\cdot} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{*\top}, \quad (5.3a)$$

$$\beta_{i,t} = \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} (\mathbf{E}_b)_{\cdot,t}, \quad \text{and} \quad (5.3b)$$

$$\lambda_{i,t} = (\mathbf{E}_c)_{i,\cdot} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} (\mathbf{E}_b)_{\cdot,t}. \quad (5.3c)$$

Observe that $\alpha_{i,t}$ and $\beta_{i,t}$ are independent zero-mean Gaussian random variables with variances

$$\text{var}(\alpha_{i,t}) = \frac{\omega^2}{NT} \|\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\|_2^2 \stackrel{(i)}{\in} \left[\frac{1}{c_u} \frac{\omega^2}{N_1 T} \|\mathbf{U}_{i,\cdot}^*\|_2^2, \frac{1}{c_\ell} \frac{\omega^2}{N_1 T} \|\mathbf{U}_{i,\cdot}^*\|_2^2 \right], \quad \text{and} \quad (5.4a)$$

$$\text{var}(\beta_{i,t}) = \frac{\omega^2}{NT} \|\mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top}\|_2^2 \stackrel{(ii)}{\in} \left[\frac{1}{c_u} \frac{\omega^2}{NT_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2, \frac{1}{c_\ell} \frac{\omega^2}{NT_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2 \right], \quad (5.4b)$$

where (i) and (ii) both follow from [\(4.1\)](#). Then we can check that

$$\begin{aligned} \frac{b_1(i,t)}{\text{var}^{1/2}(\alpha_{i,t})} + \frac{b_2(i,t)}{\text{var}^{1/2}(\beta_{i,t})} &\leq 3\sqrt{c_u}(\rho_{N,T}^2 + \rho_{N,T})\sqrt{r + \xi_{N,T}} \stackrel{(i)}{\leq} \frac{\delta}{4}, \\ \frac{b_3(i,t)}{\text{var}^{1/2}(\alpha_{i,t} + \beta_{i,t})} &\leq \sqrt{c_u} \left(\sqrt{r}(\rho_N \nu_t + \rho_T \mu_i) + \min \left\{ \frac{\mu_i}{\sqrt{N_1}}, \frac{\nu_t}{\sqrt{T_1}} \right\} \sqrt{r \xi_{N,T}} \right) \stackrel{(ii)}{\leq} \frac{\delta}{4}, \quad \text{and} \\ \frac{b_4(i,t)}{\text{var}^{1/2}(\alpha_{i,t} + \beta_{i,t})} &\leq 2\sqrt{c_u}(\rho_{N,T}^2 + \rho_{N,T}) \min \left\{ \frac{\rho_T}{\mu_i}, \frac{\rho_N}{\nu_t} \right\} \left(\sqrt{r} + \frac{\xi_{N,T}}{\sqrt{r}} \right) \stackrel{(iii)}{\leq} \frac{\delta}{4}. \end{aligned} \quad (5.5)$$

Here step (i) follows from [\(4.3\)](#), step (ii) utilizes the bound [\(4.4b\)](#), whereas step (iii) follows from the bounds [\(4.3\)](#) and [\(4.4c\)](#). Conditioned on the random matrix \mathbf{E}_b , we have

$$\lambda_{i,t} \sim \mathcal{N}\left(0, \frac{\omega^2}{NT} \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} (\mathbf{E}_b)_{\cdot,t}\|_2^2\right).$$

Therefore, for any fixed (i,t) , with probability at least $1 - O((N+T)^{-10})$, we have

$$\begin{aligned} |\lambda_{i,t}| &\stackrel{(i)}{\leq} \frac{5\omega}{\sqrt{NT}} \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} (\mathbf{E}_b)_{\cdot,t}\|_2 \sqrt{\xi_{N,T}} \\ &\stackrel{(ii)}{\leq} \frac{5C_g \omega^2}{NT} \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\| \sqrt{(r + \xi_{N,T}) \xi_{N,T}} \\ &\stackrel{(iii)}{\leq} \frac{5C_g \omega^2}{c_\ell^2 \sigma_r^*} \sqrt{\frac{1}{NTN_1T_1} (r + \xi_{N,T}) \xi_{N,T}}. \end{aligned}$$

Here the constant 5 in step (i) comes from the fact that $\mathbb{P}(X \geq 5\sqrt{\xi_{N,T}}) = O((N+T)^{-10})$ for $X \sim \mathcal{N}(0,1)$, step (ii) uses the concentration bound

$$\|\zeta(\mathbf{E}_b)_{\cdot,t}\|_2 \leq C_g \sigma \|\zeta\| \sqrt{r + \xi_{N,T}}, \quad \text{where } \zeta = \mathbf{V}_1^*(\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top},$$

which is a consequence of [Lemma 19](#) in [Appendix G](#), whereas step (iii) follows from [\(4.1\)](#). Taking the bound on $|\lambda_{i,t}|$ and [\(4.4c\)](#) together yields

$$\frac{|\lambda_{i,t}|}{\text{var}^{1/2}(\alpha_{i,t} + \beta_{i,t})} \leq \frac{5C_g \sqrt{c_u}}{c_\ell^2} \min \left\{ \frac{\rho_T}{\mu_i}, \frac{\rho_N}{\nu_t} \right\} \sqrt{\xi_{N,T} + \xi_{N,T}^2/r} \leq \frac{\delta}{4}. \quad (5.6)$$

Let $G_{i,t} = \alpha_{i,t} + \beta_{i,t}$ and $\gamma_{i,t}^* = \text{var}(\alpha_{i,t} + \beta_{i,t})$, we can check that

$$\left| \frac{1}{(\gamma_{i,t}^*)^{1/2}} (\widehat{\mathbf{M}}_d - \mathbf{M}_d^*)_{i,t} - G_{i,t} \right| \stackrel{(i)}{\leq} \sum_{j=1}^4 \frac{b_j(i,t)}{\text{var}^{1/2}(\alpha_{i,t} + \beta_{i,t})} + \frac{|\lambda_{i,t}|}{\text{var}^{1/2}(\alpha_{i,t} + \beta_{i,t})} \stackrel{(ii)}{\leq} \delta$$

where step (i) follows from [Lemma 1](#) and step (ii) follows from equations [\(5.5\)](#) and [\(5.6\)](#).

5.1.3 Proof of [Lemma 1](#)

Our proof is based on decomposing the estimation error as

$$\begin{aligned} \mathbf{M}_d - \mathbf{M}_d^* &= \mathbf{U}_2(\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top \mathbf{M}_b - \mathbf{U}_2^*(\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{M}_b^* \\ &= \mathbf{U}_2(\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top (\mathbf{M}_b - \mathbf{M}_b^*) + [\mathbf{U}_2(\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top - \mathbf{U}_2^*(\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}] \mathbf{M}_b^* \\ &= \mathbf{A} + \mathbf{B}, \end{aligned} \quad (5.7)$$

where

$$\mathbf{A} = \mathbf{U}_2(\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top (\mathbf{M}_b - \mathbf{M}_b^*), \quad \text{and} \quad \mathbf{B} = [\mathbf{U}_2(\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top - \mathbf{U}_2^*(\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}] \mathbf{M}_b^*.$$

With these definitions in hand, we now state two auxiliary lemmas that bound these two terms.

Lemma 2. *Under the conditions of [Lemma 1](#), we have the decomposition*

$$\mathbf{A} = \mathbf{U}_2^*(\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_b + \mathbf{E}_c \mathbf{V}_1^*(\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_b + \boldsymbol{\xi}_A, \quad (5.8)$$

where $|(\boldsymbol{\xi}_A)_{i,t}| \leq C_A \sum_{k=1}^4 b_k(i,t)$ holds for each (i,t) with probability at least $1 - O((N+T)^{-9})$.

See [Appendix C.2](#) for the proof.

Lemma 3. *Under the conditions of [Lemma 1](#), there exists some universal constant $C_B > 0$ such that we can decompose*

$$\mathbf{B} = \mathbf{E}_c \mathbf{V}_1^*(\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_2^{*\top} + \boldsymbol{\xi}_B, \quad (5.9)$$

where $|(\boldsymbol{\xi}_B)_{i,t}| \leq C_B \sum_{k=2}^3 b_k(i,t)$ holds for each (i,t) with probability at least $1 - O((N+T)^{-9})$.

See [Appendix C.3](#) for the proof.

Combining equation [\(5.7\)](#) with these two lemmas yields $\mathbf{M}_d - \mathbf{M}_d^* = \mathbf{Z} + \boldsymbol{\Delta}$, where the random matrix \mathbf{Z} was previously defined [\(5.1\)](#), and $\boldsymbol{\Delta} = \boldsymbol{\xi}_A + \boldsymbol{\xi}_B$. The residual matrix $\boldsymbol{\Delta}$ satisfies, with probability exceeding $1 - O((N+T)^{-9})$, that $|\Delta_{i,t}| \leq C_4 \sum_{k=1}^4 b_k(i,t)$ holds for each (i,t) with some universal constant $C_4 = C_A + C_B$.

5.2 Proof of Proposition 1

Naturally, a central step in the proof is to bound the error incurred by using $\widehat{\gamma}_{i,t}$ as an estimate of $\gamma_{i,t}^*$. Let us summarize this auxiliary claim here: with probability at least $1 - O((N+T)^{-10})$, there exists some constant $C_\gamma > 0$ such that

$$|\widehat{\gamma}_{i,t} - \gamma_{i,t}^*| \leq C_\gamma \left(\frac{\delta}{\sqrt{\xi_{N,T}}} + \sqrt{\frac{(N_1+T)r}{N_1T}} \right) \gamma_{i,t}^*. \quad (5.10)$$

Here $\delta \in (0, 1)$ is the tolerance parameter in the statement of Proposition 1. We return to prove this result in Section 5.2.2.

5.2.1 Main argument

Taking the auxiliary claim (5.10) as given, let us prove the coverage guarantee stated in Proposition 1. Define the random variable

$$V_{i,t} := (\widehat{M}_{i,t} - M_{i,t}^*) / \widehat{\gamma}_{i,t}^{1/2} - G_{i,t},$$

where $G_{i,t}$ is the standard Gaussian variable defined in Theorem 1.

By applying Theorem 1, we find that with probability at least $1 - O((N+T)^{-10})$,

$$|V_{i,t}| = \left| \frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\gamma_{i,t}^*)^{1/2}} - G_{i,t} \right| + \left| \frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\gamma_{i,t}^*)^{1/2}} - \frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\widehat{\gamma}_{i,t})^{1/2}} \right| \leq \delta + \left| \frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\gamma_{i,t}^*)^{1/2}} \right| \left| 1 - \frac{(\gamma_{i,t}^*)^{1/2}}{(\widehat{\gamma}_{i,t})^{1/2}} \right|. \quad (5.11)$$

Next we observe that

$$\left| \frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\gamma_{i,t}^*)^{1/2}} \right| \leq \left| \frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\gamma_{i,t}^*)^{1/2}} - G_{i,t} \right| + |G_{i,t}| \stackrel{(i)}{\leq} \delta + 5\sqrt{\xi_{N,T}}$$

and

$$\left| 1 - \frac{(\gamma_{i,t}^*)^{1/2}}{(\widehat{\gamma}_{i,t})^{1/2}} \right| = \frac{|\gamma_{i,t}^* - \widehat{\gamma}_{i,t}|}{(\widehat{\gamma}_{i,t})^{1/2}[(\gamma_{i,t}^*)^{1/2} + (\widehat{\gamma}_{i,t})^{1/2}]} \stackrel{(ii)}{\leq} 2C_\gamma \frac{\delta}{\sqrt{\xi_{N,T}}} + 2C_\gamma \sqrt{\frac{(N_1+T)r}{N_1T}};$$

where claim (i) follows from Theorem 1, and step (ii) follows from the auxiliary claim (5.10) and its consequence $\widehat{\gamma}_{i,t} \geq \gamma_{i,t}^*/2$, provided that δ is sufficiently small and $\min\{N_1, T\}$ is sufficiently large.

Combining these facts with our earlier bound (5.11) and defining $C_v = 12C_\gamma$, we find that

$$|V_{i,t}| \leq \delta + (\delta + 5\sqrt{\xi_{N,T}}) \left(2C_\gamma \frac{\delta}{\sqrt{\xi_{N,T}}} + 2C_\gamma \sqrt{\frac{(N_1+T)r}{N_1T}} \right) \stackrel{(iii)}{\leq} C_v \delta, \quad (5.12)$$

where step (iii) holds provided that $\min\{N_1, T\} \geq \delta^{-2} r \xi_{N,T}$.

We now use this high probability bound on $V_{i,t}$ to establish our coverage guarantee. For any $x \in \mathbb{R}$, from the definition of $V_{i,t}$, we have

$$\begin{aligned} \mathbb{P}\left(\frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\widehat{\gamma}_{i,t})^{1/2}} \leq x \right) &\stackrel{(*)}{=} \mathbb{P}(G_{i,t} + V_{i,t} \leq x, |V_{i,t}| \leq C_v \delta) + \mathbb{P}(G_{i,t} + V_{i,t} \leq x, |V_{i,t}| > C_v \delta) \quad (5.13a) \\ &\leq \mathbb{P}(G_{i,t} \leq x + C_v \delta) + \mathbb{P}(|V_{i,t}| > C_v \delta) \\ &\stackrel{(a)}{=} \Phi(x + C_v \delta) + O((N+T)^{-10}) \\ &\stackrel{(b)}{\leq} \Phi(x) + O(\delta + (N+T)^{-10}) \quad (5.13b) \end{aligned}$$

where step (a) follows from the bound (5.12); and step (b) follows since the normal CDF Φ is a $1/\sqrt{2\pi}$ -Lipschitz function. Similarly, we have the lower bound

$$\begin{aligned} \mathbb{P}\left(\frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\widehat{\gamma}_{i,t})^{1/2}} \leq x\right) &\stackrel{(c)}{\geq} \mathbb{P}(G_{i,t} + V_{i,t} \leq x, |V_{i,t}| \leq C_v \delta) \\ &\geq \mathbb{P}(G_{i,t} \leq x - C_v \delta) \\ &= \Phi(x - C_v \delta) \\ &\stackrel{(d)}{\geq} \Phi(x) - O(\delta), \end{aligned} \tag{5.13c}$$

where step (c) follows from equality (\star) in equation (5.13a); and step (d) follows from the Lipschitz property of Φ .

Finally, by taking $x = \pm\Phi^{-1}(1 - \alpha/2)$ and combining the upper and lower bounds (5.13b) and (5.13c), we find that

$$\mathbb{P}(\text{CI}_{i,t}^{1-\alpha} \ni M_{i,t}^*) = \mathbb{P}\left(\frac{\widehat{M}_{i,t} - M_{i,t}^*}{(\widehat{\gamma}_{i,t})^{1/2}} \in [\pm\Phi^{-1}(1 - \alpha/2)]\right) = 1 - \alpha + O(\delta + (N + T)^{-10}),$$

as claimed in Proposition 1.

5.2.2 Proof of the auxiliary claim (5.10)

To simplify presentation, we introduce the shorthand $\sigma = \omega/\sqrt{NT}$, along with the associated estimate $\widehat{\sigma}^2 = \widehat{\omega}^2/NT$. Using this notation, we write

$$\begin{aligned} \widehat{\gamma}_{i,t} - \gamma_{i,t}^* &= \underbrace{(\widehat{\sigma}^2 - \sigma^2) \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_{i,\cdot}^{*\top} + (\widehat{\sigma}^2 - \sigma^2) \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{\top}}_{=: \beta_1} \\ &\quad + \underbrace{\widehat{\sigma}^2 (\mathbf{U}_{i,\cdot} (\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_{i,\cdot}^\top - \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_{i,\cdot}^{*\top}) + \widehat{\sigma}^2 (\mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{\top} - \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{\top})}_{=: \beta_2} \end{aligned}$$

Next we bound each of the terms β_1 and β_2 in turn.

Bounding β_1 : In order to bound β_1 , we need to control the estimation error associated with $\widehat{\sigma}^2$. The following result provides such a bound:

Lemma 4. *Defining the constant $C_\sigma = 2\sqrt{2C_b + 1} + C_{\text{upper}}^2 + 2C_b$, we have*

$$|\widehat{\sigma}^2 - \sigma^2| \leq \alpha_1 + \alpha_2 + \alpha_3 \leq C_\sigma \sigma^2 \sqrt{\frac{(N_1 + T)r}{N_1 T}}, \tag{5.14}$$

with probability at least $1 - O((N + T)^{-10})$.

See Section 5.2.3 for the proof.

We now apply the relation (5.14) to bound β_1 , thereby obtaining

$$\begin{aligned} |\beta_1| &\leq C_\sigma \sigma^2 \sqrt{\frac{(N_1 + T)r}{N_1 T}} (\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_{i,\cdot}^{*\top} + \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{\top}) \\ &= C_\sigma \sqrt{\frac{(N_1 + T)r}{N_1 T}} \gamma_{i,t}^*. \end{aligned} \tag{5.15a}$$

Bounding β_2 : In order to bound the term β_2 , we need the following lemma characterizing the error of the plug-in estimate.

Lemma 5. *Under the conditions of Proposition 1, with probability at least $1 - O((N + T)^{-10})$,*

$$\sigma^2 |\mathbf{U}_{i,\cdot} (\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_{i,\cdot}^\top - \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_{i,\cdot}^{*\top}| + \sigma^2 |\mathbf{V}_{t,\cdot} (\mathbf{V}_1^\top \mathbf{V}_1)^{-1} \mathbf{V}_{t,\cdot}^\top - \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{*\top}| \leq \frac{\delta \gamma_{i,t}^*}{\sqrt{\xi_{N,T}}}.$$

See Appendix D.2 for the proof.

When $\min\{N_1, T\} \geq r$, a direct consequence of (5.14) is that $\hat{\sigma}^2 \leq (\sqrt{2}C_\sigma + 1)\sigma^2$. Combining this bound with Lemma 5 yields

$$|\beta_2| \leq (\sqrt{2}C_\sigma + 1) \frac{\delta}{\sqrt{\xi_{N,T}}} \gamma_{i,t}^*, \quad (5.15b)$$

Collecting together the two bounds (5.15a) and (5.15b) on β_1 and β_2 , respectively, yields the upper bound

$$|\hat{\gamma}_{i,t} - \gamma_{i,t}^*| \leq |\beta_1| + |\beta_2| \leq \left((\sqrt{2}C_\sigma + 1) \frac{\delta}{\sqrt{\xi_{N,T}}} + C_\sigma \sqrt{\frac{(N_1 + T)r}{N_1 T}} \right) \gamma_{i,t}^*.$$

Let $C_\gamma = \sqrt{2}C_\sigma + 1$ to achieve the claimed bound (5.10).

5.2.3 Proof of Lemma 4

We first decompose $\hat{\sigma}^2$ into a sum of three terms as

$$\begin{aligned} \hat{\sigma}^2 &= \frac{1}{N_1 T} \|\mathbf{M}_{\text{upper}}^* + \mathbf{E}_{\text{upper}} - \widehat{\mathbf{M}}_{\text{upper}}\|_{\text{F}}^2 \\ &= \frac{1}{N_1 T} \|\mathbf{E}_{\text{upper}}\|_{\text{F}}^2 + \frac{1}{N_1 T} \|\mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}}\|_{\text{F}}^2 + \frac{2}{N_1 T} \langle \mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}}, \mathbf{E}_{\text{upper}} \rangle. \end{aligned}$$

As our analysis will show, the first quantity concentrates around σ^2 . Thus, it is natural to bound the estimation error associated with $\hat{\sigma}^2$ in the following way:

$$|\hat{\sigma}^2 - \sigma^2| \leq \underbrace{\left| \frac{1}{N_1 T} \|\mathbf{E}_{\text{upper}}\|_{\text{F}}^2 - \sigma^2 \right|}_{=:\alpha_1} + \underbrace{\frac{1}{N_1 T} \|\mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}}\|_{\text{F}}^2}_{=:\alpha_2} + \underbrace{\frac{2}{N_1 T} \|\mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}}\|_{\text{F}} \|\mathbf{E}_{\text{upper}}\|_{\text{F}}}_{=:\alpha_3}.$$

By applying Bernstein's inequality [Ver17, Theorem 2.8.1], we can conclude that for some constant $C_b > 0$, as long as $N_1 T \geq \xi_{N,T}$, we find that

$$\alpha_1 = \frac{1}{N_1 T} \left| \sum_{i=1}^{N_1} \sum_{j=1}^T (E_{i,j}^2 - \sigma^2) \right| \leq C_b \sigma^2 \sqrt{\frac{\xi_{N,T}}{N_1 T}} + C_b \sigma^2 \frac{\xi_{N,T}}{N_1 T} \leq 2C_b \sigma^2 \sqrt{\frac{\xi_{N,T}}{N_1 T}} \quad (5.16a)$$

with probability at least $1 - O((N + T)^{-10})$. As a consequence, we have

$$\|\mathbf{E}_{\text{upper}}\|_{\text{F}}^2 \leq N_1 T \sigma^2 + N_1 T \alpha_1 \leq (2C_b + 1) N_1 T \sigma^2, \quad (5.16b)$$

provided that $N_1 T \geq \xi_{N,T}$.

In order to bound the terms α_2 and α_3 , we make use of the following lemma.

Lemma 6. *Under the conditions of Proposition 1, we have*

$$\|\mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}}\|_{\text{F}} \leq C_{\text{upper}}\sigma\sqrt{(T + N_1)r}$$

with probability at least $1 - O((N + T)^{-10})$.

See Appendix D.1 for the proof.

Combining Lemma 6 with the bounds (5.16) yields

$$\alpha_2 \leq C_{\text{upper}}^2 \frac{\sigma^2 (T + N_1) r}{N_1 T} \quad \text{and} \quad \alpha_3 \leq 2\sqrt{2C_b + 1}C_{\text{upper}}\sigma^2 \sqrt{\frac{(N_1 + T)r}{N_1 T}}. \quad (5.17)$$

Putting together the upper bounds for α_1 , α_2 and α_3 yields the claimed bound (5.14).

5.3 Proof of lower bounds

In this section, we prove the Cramér–Rao and local minimax bounds stated in the main text. While our formal statement (Theorem 2) only covers the four-block case, here we actually provide a proof for the multi-block case, so that we match our general set of achievable results (Theorem 3).

Let us recall the set-up for our genie-aided problem associated with estimating entry $M_{i,t}^*$ of the matrix. The only unknown quantities are the vectors $\mathbf{x}^* := \mathbf{X}_{i,\cdot}^*$ and $\mathbf{y}^* := \mathbf{Y}_{i,\cdot}^*$, and our goal is to estimate the inner product $M_{i,t}^* = \langle \mathbf{x}^*, \mathbf{y}^* \rangle$ based on linear measurements of the form

$$\{\mathbf{M}_{i,s} = \langle \mathbf{x}^*, \mathbf{Y}_{s,\cdot}^* \rangle + E_{i,s}\}_{s=1}^{\bar{T}_1} \quad \text{and} \quad \{\mathbf{M}_{k,t} = \langle \mathbf{X}_{k,\cdot}^*, \mathbf{y}^* \rangle + E_{k,t}\}_{k=1}^{\bar{N}_1}.$$

Cramér–Rao lower bound: We first compute the Cramér–Rao lower bound (CRLB) for the problem. Using $\theta = (\mathbf{x}^*, \mathbf{y}^*)$ to denote the unknown parameters, our goal is to estimate the functional $\Psi(\theta) := \langle \mathbf{x}^*, \mathbf{y}^* \rangle$. The CRLB is given by $\nabla\Psi(\theta)^T \mathbf{I}^{-1}(\theta) \nabla\Psi(\theta)$, where $\mathbf{I}(\theta)$ is the Fisher information matrix, and the gradient takes the form $\nabla\Psi(\theta) = [\mathbf{y}^* \quad \mathbf{x}^*]^T$. Given that our measurements are linear with Gaussian observation noise with variance $\omega^2/(NT)$, the Fisher information matrix takes the form

$$\mathbf{I}(\theta) = \frac{NT}{\omega^2} \text{blkdiag} \left(\sum_{s=1}^{\bar{T}_1} \mathbf{Y}_{s,\cdot}^{*\top} \mathbf{Y}_{s,\cdot}^*, \quad \sum_{k=1}^{\bar{N}_1} \mathbf{X}_{k,\cdot}^{*\top} \mathbf{X}_{k,\cdot}^* \right).$$

Putting together the pieces, we find that the CRLB is given by

$$\begin{aligned} \text{CRLB}(M_{i,t}^*) &= \mathbf{Y}_{t,\cdot}^{*\top} \left(\frac{NT}{\omega^2} \sum_{s=1}^{\bar{T}_1} \mathbf{Y}_{s,\cdot}^{*\top} \mathbf{Y}_{s,\cdot}^* \right)^{-1} \mathbf{Y}_{t,\cdot}^* + \mathbf{X}_{i,\cdot}^{*\top} \left(\frac{NT}{\omega^2} \sum_{k=1}^{\bar{N}_1} \mathbf{X}_{k,\cdot}^{*\top} \mathbf{X}_{k,\cdot}^* \right)^{-1} \mathbf{X}_{i,\cdot}^* \\ &\stackrel{(i)}{=} \frac{\omega^2}{NT} \left\{ \mathbf{V}_{t,\cdot}^{*\top} \left(\sum_{s=1}^{\bar{T}_1} \mathbf{V}_{s,\cdot}^{*\top} \mathbf{V}_{s,\cdot}^* \right)^{-1} \mathbf{V}_{t,\cdot}^* + \mathbf{U}_{i,\cdot}^{*\top} \left(\sum_{k=1}^{\bar{N}_1} \mathbf{U}_{k,\cdot}^{*\top} \mathbf{U}_{k,\cdot}^* \right)^{-1} \mathbf{U}_{i,\cdot}^* \right\} \\ &= \underbrace{\frac{\omega^2}{NT} \left\{ \mathbf{V}_{t,\cdot}^{*\top} (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^* + \mathbf{U}_{i,\cdot}^{*\top} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_{i,\cdot}^* \right\}}_{\equiv \gamma_{i,t}^*} \end{aligned}$$

Here step (i) uses the fact that $\mathbf{X}^* = \mathbf{U}^*(\boldsymbol{\Sigma}^*)^{1/2}$ and $\mathbf{Y}^* = \mathbf{V}^*(\boldsymbol{\Sigma}^*)^{1/2}$, along with some algebra.

Computation of the local minimax bound: We now turn to computation of the local minimax lower bound, where we make use of a Bayesian form of the CRLB. We begin by observing that for any $\varepsilon > 0$, the local minimax risk for estimating $\mathbf{M}_{i,t}^*$ can be lower bounded as

$$R^*(\varepsilon) \geq \inf_{\widehat{M}_{i,t}} \mathbb{E}_{(\mathbf{x}, \mathbf{y}) \sim \pi} \mathbb{E} \left[(\widehat{M}_{i,t} - \langle \mathbf{x}, \mathbf{y} \rangle)^2 \right],$$

where π is any prior supported on the Cartesian product space. $\mathcal{B}_\infty(\mathbf{x}^*, \varepsilon) \times \mathcal{B}_\infty(\mathbf{y}^*, \varepsilon)$.

Following a standard avenue for computing Bayesian CRLBs [PW24], we first define the squared cosine density function $g(u) = \cos^2(\pi u/2)$ for $u \in [-1, 1]$. Using this building block, we then define the product prior distribution $\pi(\theta_1, \dots, \theta_{2r}) = \prod_{j=1}^{2r} \pi_j(\theta_j)$ with marginals

$$\pi_j(\theta) = \begin{cases} \frac{1}{\varepsilon} g\left(\frac{\theta - x_j^*}{\varepsilon}\right) & \text{for } \theta_j \in [x_j^* - \varepsilon, x_j^* + \varepsilon] \text{ and } j = 1, \dots, r, \text{ and} \\ \frac{1}{\varepsilon} g\left(\frac{\theta - y_{j-r}^*}{\varepsilon}\right) & \text{for } \theta_j \in [y_{j-r}^* - \varepsilon, y_{j-r}^* + \varepsilon] \text{ and } j = r + 1, \dots, 2r. \end{cases}$$

With this choice of prior, we can compute the lower bound

$$\begin{aligned} R^*(\varepsilon) &\stackrel{(i)}{\geq} \mathbf{Y}_{t,\cdot}^* \left(\frac{NT}{\omega^2} \sum_{s=1}^{\bar{T}_1} \mathbf{Y}_{s,\cdot}^{*\top} \mathbf{Y}_{s,\cdot}^* + \frac{\pi^2}{\varepsilon^2} \mathbf{I}_r \right)^{-1} \mathbf{Y}_{t,\cdot}^{*\top} + \mathbf{X}_{i,\cdot}^* \left(\frac{NT}{\omega^2} \sum_{k=1}^{\bar{N}_1} \mathbf{X}_{k,\cdot}^{*\top} \mathbf{X}_{k,\cdot}^* + \frac{\pi^2}{\varepsilon^2} \mathbf{I}_r \right)^{-1} \mathbf{X}_{i,\cdot}^{*\top} \\ &\stackrel{(ii)}{=} \underbrace{\frac{\omega^2}{NT} \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^* + \frac{\pi^2 \omega^2}{\varepsilon^2 NT} (\boldsymbol{\Sigma}^*)^{-1})^{-1} \mathbf{V}_{t,\cdot}^{*\top} + \frac{\omega^2}{NT} \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^* + \frac{\pi^2 \omega^2}{\varepsilon^2 NT} (\boldsymbol{\Sigma}^*)^{-1})^{-1} \mathbf{U}_{i,\cdot}^{*\top}}_{=: \tilde{\gamma}_{i,t}(\varepsilon)} \end{aligned}$$

where step (i) makes use of some standard computations with squared cosine priors (e.g., §29.2 in the book [PW24]); and step (ii) follows from the spectral decomposition of \mathbf{M}^* , along with some algebra.

Thus far, we proven that $R^*(\varepsilon) \geq \tilde{\gamma}_{i,t}(\varepsilon)$. It remains to show that with the choice $\varepsilon = \varepsilon_{N,T} \equiv \sqrt{\sigma_r^*} \max\{1/\sqrt{N}, 1/\sqrt{T}\}$ we have

$$\Delta_{i,t}^* := \gamma_{i,t}^* - \tilde{\gamma}_{i,t}(\varepsilon) \leq \gamma_{i,t}^* \frac{c_u \pi^2}{c_\ell} \rho_{N,T}^2, \quad (5.18)$$

from which it will follow that $R^*(\varepsilon) \geq \tilde{\gamma}_{i,t}(\varepsilon) = \gamma_{i,t}^* - \Delta_{i,t}^* \geq (1 - \frac{c_u \pi^2}{c_\ell} \rho_{N,T}^2) \gamma_{i,t}^*$, as claimed.

From the definitions of $\tilde{\gamma}_{i,t}(\varepsilon)$ and $\gamma_{i,t}^*$, we have

$$\begin{aligned} \Delta_{i,t}^* = \gamma_{i,t}^* - \tilde{\gamma}_{i,t}(\varepsilon) &= \frac{\pi^2 \omega^4}{\varepsilon^2 (NT)^2} \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{V}_1^{*\top} \mathbf{V}_1^* + \frac{\pi^2 \omega^2}{\varepsilon^2 NT} (\boldsymbol{\Sigma}^*)^{-1})^{-1} \mathbf{V}_{t,\cdot}^{*\top} \\ &\quad + \frac{\pi^2 \omega^4}{\varepsilon^2 (NT)^2} \mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^* + \frac{\pi^2 \omega^2}{\varepsilon^2 NT} (\boldsymbol{\Sigma}^*)^{-1})^{-1} \mathbf{U}_{i,\cdot}^{*\top} \\ &\leq \frac{\pi^2 \omega^4}{\varepsilon^2 (NT)^2} \left(\|(\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}\|^2 \|(\boldsymbol{\Sigma}^*)^{-1}\| \|\mathbf{V}_{t,\cdot}^*\|_2^2 + \|(\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1}\|^2 \|(\boldsymbol{\Sigma}^*)^{-1}\| \|\mathbf{U}_{i,\cdot}^*\|_2^2 \right). \end{aligned}$$

Now observe that $\|(\boldsymbol{\Sigma}^*)^{-1}\| \leq \frac{1}{\sigma_r^*}$, and that the sub-block condition (4.11) ensures that

$$\|(\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}\|^2 \leq \frac{T^2}{c_\ell^2 T_1^2} \quad \text{and} \quad \|(\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1}\|^2 \leq \frac{1}{c_\ell^2} \frac{N^2}{N_1^2}.$$

Combining these facts, we find that

$$\Delta_{i,t}^* \leq \frac{\pi^2 \omega^4}{c_\ell^2 \varepsilon^2 \sigma_r^* (NT)^2} \left(\frac{T^2}{T_1^2} \|\mathbf{V}_{t,\cdot}^*\|_2^2 + \frac{N^2}{N_1^2} \|\mathbf{U}_{i,\cdot}^*\|_2^2 \right). \quad (5.19a)$$

On the other hand, using the c_u -bound in the sub-block condition (4.11), we have

$$\frac{1}{c_u} \omega^2 \frac{1}{N_1 T} \|\mathbf{U}_{i,\cdot}^*\|_2^2 + \frac{1}{c_u} \omega^2 \frac{1}{NT_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2 \leq \gamma_{i,t}^*. \quad (5.19b)$$

Combining inequalities (5.19a) and (5.19b) yields the bound

$$\Delta_{i,t}^* \leq \frac{c_u \pi^2 \omega^2}{c_\ell^2 \varepsilon^2 \sigma_r^*} \max \left\{ \frac{1}{NT_1}, \frac{1}{N_1 T} \right\} \gamma_{i,t}^* \leq \frac{c_u \pi^2 \omega^2}{c_\ell^2 \sigma_r^{*,2}} \frac{1}{\min\{T_1, N_1\}} \underbrace{\frac{\sigma_r^*}{\varepsilon^2} \max\left\{ \frac{1}{N}, \frac{1}{T} \right\}}_{=1} \gamma_{i,t}^* \stackrel{(i)}{\leq} \frac{c_u \pi^2}{c_\ell^2} \rho_{N,T}^2 \gamma_{i,t}^*,$$

where step (i) follows from our choice of ε , and the definition of $\rho_{N,T}$. This establishes the auxiliary claim (5.18), and thus the claimed lower bound.

6 Discussion

In this paper, we proposed and analyzed a simple algorithm for causal inference with panel data under staggered adoption design. It is appealing from a computational point of view, since it is non-iterative, requiring only elementary matrix operations and singular value decomposition. At the same time, we demonstrated that its statistical properties are also attractive, in that it achieves estimation accuracy, in an elementwise sense, that matches non-asymptotic lower bounds applicable to any estimator. Moreover, we showed how this theory enables data-driven construction of confidence intervals for the missing potential outcomes. We note that our development relies on a more general inferential toolbox for the SVD algorithm in the matrix denoising model, which might also be useful in other problems.

Moving forward, let us outline some open questions, along with broader directions for future study. Our current theory relies on a noise condition (4.3) and an incoherence condition (4.4); it remains unclear whether or not these conditions are improvable, for instance in terms of dependence on the rank r . Moreover, it would be interesting to extend our guarantees to accommodate matrices that are only approximately low-rank, along with more general noise models. Finally, while the staggered adoption design arises frequently, it would be interesting to understand optimal procedures for more general mechanisms of missing data.

Acknowledgements

Y. Yan was supported by the Norbert Wiener Postdoctoral Fellowship from MIT. M. J. Wainwright was partially supported by ONR grant N00014-21-1-2842 and NSF grant DMS-2311072.

References

- [Aba21] A. Abadie. ‘‘Using synthetic controls: Feasibility, data requirements, and methodological aspects’’. In: *Journal of Economic Literature* 59.2 (2021), pp. 391–425 (cit. on p. 1).

- [Abb+20] E. Abbe, J. Fan, K. Wang, Y. Zhong, et al. “Entrywise eigenvector analysis of random matrices with low expected rank”. In: *Annals of Statistics* 48.3 (2020), pp. 1452–1474 (cit. on p. 4).
- [ADH15] A. Abadie, A. Diamond, and J. Hainmueller. “Comparative politics and the synthetic control method”. In: *American Journal of Political Science* 59.2 (2015), pp. 495–510 (cit. on p. 1).
- [AG03] A. Abadie and J. Gardeazabal. “The economic costs of conflict: A case study of the Basque Country”. In: *American economic review* 93.1 (2003), pp. 113–132 (cit. on p. 1).
- [Aga+23] A. Agarwal, M. Dahleh, D. Shah, and D. Shen. “Causal matrix completion”. In: *The Thirty Sixth Annual Conference on Learning Theory*. PMLR. 2023, pp. 3821–3826 (cit. on pp. 2, 4, 17).
- [AI22] S. Athey and G. W. Imbens. “Design-based analysis in difference-in-differences settings with staggered adoption”. In: *Journal of Econometrics* 226.1 (2022), pp. 62–79 (cit. on pp. 1, 5).
- [ASS20] A. Agarwal, D. Shah, and D. Shen. “Synthetic interventions”. In: *arXiv preprint arXiv:2006.07691* (2020) (cit. on p. 4).
- [Ath+21] S. Athey, M. Bayati, N. Doudchenko, G. Imbens, and K. Khosravi. “Matrix completion methods for causal panel data models”. In: *Journal of the American Statistical Association* 116.536 (2021), pp. 1716–1730 (cit. on pp. 2, 4, 5, 17).
- [BM03] S. Burer and R. D. Monteiro. “A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization”. In: *Mathematical Programming* 95.2 (2003), pp. 329–357 (cit. on p. 4).
- [BN21] J. Bai and S. Ng. “Matrix completion, counterfactuals, and factor analysis of missing data”. In: *Journal of the American Statistical Association* 116.536 (2021), pp. 1746–1763 (cit. on pp. 2, 4, 17).
- [Cai+21] C. Cai, G. Li, Y. Chi, H. V. Poor, and Y. Chen. “Subspace estimation from unbalanced and incomplete data matrices: $\ell_{2,\infty}$ statistical guarantees”. In: *The Annals of Statistics* 49.2 (2021), pp. 944–967 (cit. on p. 4).
- [CBN23] E. Cahan, J. Bai, and S. Ng. “Factor-based imputation of missing values and covariances in panel data of large dimensions”. In: *Journal of Econometrics* 233.1 (2023), pp. 113–131 (cit. on pp. 2, 4, 17).
- [Che15] Y. Chen. “Incoherence-optimal matrix completion”. In: *IEEE Transactions on Information Theory* 61.5 (2015), pp. 2909–2923 (cit. on p. 14).
- [Che+19] Y. Chen, J. Fan, C. Ma, and Y. Yan. “Inference and uncertainty quantification for noisy matrix completion”. In: *Proceedings of the National Academy of Sciences* 116.46 (2019), pp. 22931–22937 (cit. on p. 4).
- [Che+20] Y. Chen, Y. Chi, J. Fan, C. Ma, and Y. Yan. “Noisy matrix completion: Understanding statistical guarantees for convex relaxation via nonconvex optimization”. In: *SIAM Journal on Optimization* 30.4 (2020), pp. 3098–3121 (cit. on p. 4).

- [Che+21a] Y. Chen, Y. Chi, J. Fan, C. Ma, et al. “Spectral methods for data science: A statistical perspective”. In: *Foundations and Trends® in Machine Learning* 14.5 (2021), pp. 566–806 (cit. on pp. 4, 50, 55, 56, 64).
- [Che+21b] Y. Chen, J. Fan, C. Ma, and Y. Yan. “Bridging convex and nonconvex optimization in robust PCA: Noise, outliers and missing data”. In: *The Annals of Statistics* 49.5 (2021), pp. 2948–2971 (cit. on p. 4).
- [CKL23] J. Choi, H. Kwon, and Y. Liao. “Inference for Low-rank Models without Estimating the Rank”. In: *arXiv preprint arXiv:2311.16440* (2023) (cit. on p. 4).
- [CLC19] Y. Chi, Y. M. Lu, and Y. Chen. “Nonconvex optimization meets low-rank matrix factorization: An overview”. In: *IEEE Transactions on Signal Processing* 67.20 (2019), pp. 5239–5269 (cit. on pp. 4, 14).
- [CR09] E. Candès and B. Recht. “Exact Matrix Completion via Convex Optimization”. In: *Foundations of Computational Mathematics* 9.6 (2009), pp. 717–772 (cit. on pp. 2, 4, 14).
- [CT10] E. Candès and T. Tao. “The Power of Convex Relaxation: Near-Optimal Matrix Completion”. In: *IEEE Transactions on Information Theory* 56.5 (2010), pp. 2053–2080 (cit. on p. 4).
- [CY23] J. Choi and M. Yuan. “Matrix Completion When Missing Is Not at Random and Its Applications in Causal Panel Data Models”. In: *arXiv preprint arXiv:2308.02364* (2023) (cit. on pp. 2, 4, 17).
- [EK15] N. El Karoui. “On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators”. In: *Probability Theory and Related Fields* (2015), pp. 1–81 (cit. on p. 4).
- [FLP21] V. Farias, A. Li, and T. Peng. “Learning treatment effects in panels with general intervention patterns”. In: *Advances in Neural Information Processing Systems* 34 (2021), pp. 14001–14013 (cit. on p. 4).
- [HHY08] B.-N. Huang, M. J. Hwang, and C. W. Yang. “Causal relationship between energy consumption and GDP growth revisited: a dynamic panel data approach”. In: *Ecological economics* 67.1 (2008), pp. 41–54 (cit. on p. 1).
- [HI08] Y. Hu and S. Izumida. “Ownership concentration and corporate performance: A causal analysis with Japanese panel data”. In: *Corporate Governance: An International Review* 16.4 (2008), pp. 342–358 (cit. on p. 1).
- [IR15] G. W. Imbens and D. B. Rubin. *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press, 2015 (cit. on p. 1).
- [Klo14] O. Klopp. “Noisy low-rank matrix completion with general sampling distribution”. In: *Bernoulli* 20.1 (2014), pp. 282–303 (cit. on p. 4).
- [KMO10] R. H. Keshavan, A. Montanari, and S. Oh. “Matrix Completion From a Few Entries”. In: *IEEE Transactions on Information Theory* 56.6 (2010), pp. 2980–2998 (cit. on p. 4).
- [LBNE08] M. S. Lewis-Beck, R. Nadeau, and A. Elias. “Economics, party, and the vote: Causality issues and panel data”. In: *American Journal of Political Science* 52.1 (2008), pp. 84–95 (cit. on p. 1).

- [LR23] L. Lei and B. Ross. “Estimating Counterfactual Matrix Means with Short Panel Data”. In: *arXiv preprint arXiv:2312.07520* (2023) (cit. on p. 4).
- [Ma+20] C. Ma, K. Wang, Y. Chi, and Y. Chen. “Implicit Regularization in Nonconvex Statistical Estimation: Gradient Descent Converges Linearly for Phase Retrieval, Matrix Completion, and Blind Deconvolution”. In: *Foundations of Computational Mathematics* 20.3 (2020), pp. 451–632 (cit. on p. 4).
- [NW12] S. Negahban and M. Wainwright. “Restricted strong convexity and weighted matrix completion: Optimal bounds with noise”. In: *Journal of Machine Learning Research* (2012), pp. 1665–1697 (cit. on pp. 2, 4).
- [PW24] Y. Polyanskiy and Y. Wu. *Information Theory: From Coding to Learning*. Cambridge University Press, 2024 (cit. on p. 25).
- [RFP10] B. Recht, M. Fazel, and P. A. Parrilo. “Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization”. In: *SIAM review* 52.3 (2010), pp. 471–501 (cit. on p. 4).
- [RR83] P. R. Rosenbaum and D. B. Rubin. “The central role of the propensity score in observational studies for causal effects”. In: *Biometrika* 70.1 (1983), pp. 41–55 (cit. on p. 1).
- [SCC17] P. Sur, Y. Chen, and E. J. Candès. “The Likelihood Ratio Test in High-Dimensional Logistic Regression Is Asymptotically a Rescaled Chi-Square”. In: *arXiv:1706.01191, accepted to Probability Theory and Related Fields* (2017) (cit. on p. 4).
- [Sre04] N. Srebro. “Learning with matrix factorizations”. In: *Ph. D. thesis* (2004) (cit. on p. 4).
- [ST21] A. M. Shaikh and P. Toulis. “Randomization tests in observational studies with staggered adoption of treatment”. In: *Journal of the American Statistical Association* 116.536 (2021), pp. 1835–1848 (cit. on pp. 1, 5).
- [Ver17] R. Vershynin. *High Dimensional Probability*. Cambridge University Press, 2017 (cit. on pp. 23, 48, 68, 69).
- [Wed72] P.-Å. Wedin. “Perturbation bounds in connection with singular value decomposition”. In: *BIT Numerical Mathematics* 12.1 (1972), pp. 99–111 (cit. on pp. 50, 55).
- [XY21] D. Xia and M. Yuan. “Statistical inferences of linear forms for noisy matrix completion”. In: *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 83.1 (2021), pp. 58–77 (cit. on p. 4).
- [YCF21] Y. Yan, Y. Chen, and J. Fan. “Inference for heteroskedastic PCA with missing data”. In: *arXiv preprint arXiv:2107.12365* (2021) (cit. on p. 4).
- [ZC23] Y. Zhou and Y. Chen. “Deflated HeteroPCA: Overcoming the curse of ill-conditioning in heteroskedastic PCA”. In: *arXiv preprint arXiv:2303.06198* (2023) (cit. on p. 4).

A Details of simulation study

In this appendix, we describe the simulation study underlying the results shown in [Figure 2](#). We set the dimensions $N = T = 1000$, the rank $r = 5$, and consider the staggered adoption design

with $k = 3$ groups with dimension parameters $N_1 = T_1 = 400$, $N_2 = T_2 = 300$ and $N_3 = T_3 = 300$. We generate $\mathbf{M}^* = \mathbf{X}^* \mathbf{Y}^{*\top}$ where $\mathbf{X}^* \in \mathbb{R}^{N \times r}$ and $\mathbf{Y}^* \in \mathbb{R}^{T \times r}$ are random orthonormal matrices following the Haar distribution over the Grassmann manifolds, and set the noise level $\omega = 1$. For any entry (i, t) , we define the rescaled error

$$R_{i,t} = (\widehat{M}_{i,t} - M_{i,t}^*) / \sqrt{\widehat{\gamma}_{i,t}}.$$

Figure 2 gives plots of this quantity, where we select one entry $M_{800,800}$ from the block $\mathbf{M}_{3,3}^*$ and another entry $M_{800,500}$ from the block $\mathbf{M}_{3,2}^*$.

B Distributional theory without condition (4.4c)

In this section, we discuss how to characterize the output distribution of Algorithm 1 without imposing the lower bounds (4.4c) on the local incoherence parameters.

B.1 More careful control

Recall the decomposition from Lemma 1. Some computation shows that the entrywise variance of \mathbf{Z} takes the form

$$\begin{aligned} \text{var}(Z_{i,t}) &= \frac{\omega^2}{NT} \|\mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top}\|_2^2 + \frac{\omega^2}{NT} \|\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\|_2^2 \\ &\quad + \frac{\omega^4}{N^2 T^2} \text{tr}[\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top}]. \end{aligned} \quad (\text{B.1})$$

The following lemma provides a lower bound on the variance of $Z_{i,t}$.

Lemma 7. *Under the sub-block condition (4.1), the variance of $Z_{i,t}$ is lower bounded by*

$$\text{var}(Z_{i,t}) \geq \frac{1}{c_u} \frac{\omega^2}{NT_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2 + \frac{1}{c_u} \frac{\omega^2}{N_1 T} \|\mathbf{U}_{i,\cdot}^*\|_2^2 + \frac{1}{c_u^2} \frac{\omega^4}{\sigma_r^{*2}} \frac{1}{NTN_1 T_1}.$$

See Appendix B.2 for the proof.

Combining the upper bound on $|\Delta_{i,t}|$ from Lemma 1 and the lower bound on $\text{var}(Z_{i,t})$ from Lemma 7, we can check that the size of $|\Delta_{i,t}|$ in the decomposition in Lemma 1 is dominated by the standard deviation of $Z_{i,j}$. We summarize our conclusion in the following:

Lemma 8. *Under all conditions of Theorem 1 with the exception of the lower bound (4.4c), the decomposition (5.2a) satisfies $|\Delta_{i,t}| \leq \delta \sqrt{\text{var}(Z_{i,t})}$ with probability at least $1 - O((N + T)^{-10})$.*

The above lemma shows that the error $(\mathbf{M}_d - \mathbf{M}_d^*)_{i,t}$ is well-approximated by the variable $Z_{i,t}$, and its distribution is fully determined by equation (5.1). Comparing with Theorem 1, this lemma does not impose the lower bound (4.4c) on either of $\|\mathbf{U}_{i,\cdot}^*\|_2$ and $\|\mathbf{V}_{t,\cdot}^*\|_2$, hence is more general and can be useful especially when $\|\mathbf{U}_{i,\cdot}^*\|_2$ and $\|\mathbf{V}_{t,\cdot}^*\|_2$ are vanishingly small.

B.2 Proof of Lemma 7

Recall from equation (5.3) the decomposition $Z_{i,t} = \alpha_{i,t} + \beta_{i,t} + \lambda_{i,t}$, along with the variances $\alpha_{i,t}$ and $\beta_{i,t}$ given in equation (5.4). Given the sub-block condition (4.1), they are lower bounded by

$$\text{var}(\alpha_{i,t}) \geq \frac{1}{c_u} \frac{\omega^2}{NT_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2, \quad \text{var}(\beta_{i,t}) \geq \frac{1}{c_u} \frac{\omega^2}{N_1 T} \|\mathbf{U}_{i,\cdot}^*\|_2^2.$$

Then we lower bound the variance of $\lambda_{i,t}$. Recall that

$$\lambda_{i,t} = (\mathbf{E}_c)_{i,\cdot} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} (\mathbf{E}_b)_{\cdot,t}.$$

It is straightforward to check that $\lambda_{i,t}$ is mean zero. Since \mathbf{E}_c and \mathbf{E}_b are independent, we have

$$\begin{aligned} \text{var}(\lambda_{i,t}) &\stackrel{(i)}{=} \frac{\omega^2}{NT} \mathbb{E}[(\mathbf{E}_c)_{i,\cdot} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top} (\mathbf{E}_c)_{i,\cdot}^\top] \\ &\stackrel{(ii)}{=} \frac{\omega^4}{N^2 T^2} \text{tr}[\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top}] \\ &\stackrel{(iii)}{\geq} \frac{1}{c_u} \frac{N}{N_1} \frac{\omega^4}{N^2 T^2} \text{tr}[\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-2} (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top}] \\ &= \frac{1}{c_u} \frac{N}{N_1} \frac{\omega^4}{N^2 T^2} \text{tr}[(\boldsymbol{\Sigma}^*)^{-2} (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}] \stackrel{(iv)}{\geq} \frac{1}{c_u^2} \frac{\omega^4}{N^2 T^2} \frac{NT}{N_1 T_1} \text{tr}[(\boldsymbol{\Sigma}^*)^{-2}] \geq \frac{1}{c_u^2} \frac{\omega^4}{\sigma_r^{*2}} \frac{1}{NTN_1 T_1}. \end{aligned}$$

Here (i) uses the fact that $\mathbb{E}[(\mathbf{E}_b)_{\cdot,t} (\mathbf{E}_b)_{\cdot,t}^\top] = \sigma^2 \mathbf{I}_{N_1}$, (ii) utilizes $\mathbb{E}[(\mathbf{E}_c)_{i,\cdot}^\top (\mathbf{E}_c)_{i,\cdot}] = \sigma^2 \mathbf{I}_{T_1}$, while (iii) and (iv) both follow from (4.1). Since $\alpha_{i,t}$, $\beta_{i,t}$ and $\lambda_{i,t}$ are mutually uncorrelated, we conclude that

$$\text{var}(Z_{i,t}) = \text{var}(\alpha_{i,t}) + \text{var}(\beta_{i,t}) + \text{var}(\lambda_{i,t}) \geq \frac{1}{c_u} \frac{\omega^2}{NT_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2 + \frac{1}{c_u} \frac{\omega^2}{N_1 T} \|\mathbf{U}_{i,\cdot}^*\|_2^2 + \frac{1}{c_u^2} \frac{\omega^4}{\sigma_r^{*2}} \frac{1}{NTN_1 T_1}$$

as claimed.

C Proofs of Lemmas from Section 5.1.3

This section is devoted to the proofs of the lemmas stated in Section 5.1.3. Appendix C.1 provides some useful matrix/subspace perturbation results, obtained by applying general results for matrix denoising (cf. Appendix F) to the causal panel data model. We use these results to prove the lemmas stated in Section 5.1.3.

Throughout this section, in addition to the rescaled noise level ω defined in Section 2.1, we will also use the standard deviation of the noise added on each observed entry of \mathbf{M}^* , given by

$$\sigma := \omega / \sqrt{NT}. \tag{C.1}$$

C.1 A few perturbation bounds

In order to apply our general theory for matrix denoising model to \mathbf{M}_{left} and $\mathbf{M}_{\text{upper}}$, we first need to understand the spectra of the corresponding ground truth matrices $\mathbf{M}_{\text{left}}^*$ and $\mathbf{M}_{\text{upper}}^*$.

Lemma 9. Under the sub-block condition (4.1), we have

$$\sqrt{c_\ell \frac{T_1}{T}} \sigma_r^* \leq \sigma_r(\mathbf{M}_{\text{left}}^*) \leq \sqrt{c_u \frac{T_1}{T}} \sigma_r^*, \quad (\text{C.2a})$$

$$\sqrt{c_\ell \frac{N_1}{N}} \sigma_r^* \leq \sigma_r(\mathbf{M}_{\text{upper}}^*) \leq \sqrt{c_u \frac{N_1}{N}} \sigma_r^*. \quad (\text{C.2b})$$

See [Appendix C.4](#) for the proof.

Define the matrices $\mathbf{E}_{\text{left}} = \mathbf{M}_{\text{left}} - \mathbf{M}_{\text{left}}^*$ and $\mathbf{E}_{\text{upper}} = \mathbf{M}_{\text{upper}} - \mathbf{M}_{\text{upper}}^*$. Recall that the truncated rank- r SVD of \mathbf{M}_{left} was assumed to be $\mathbf{U}_{\text{left}} \boldsymbol{\Sigma}_{\text{left}} \mathbf{V}_{\text{left}}^\top$ in the main text. However, we simplify the notation in the proof by dropping the subscript: in particular, we let $\mathbf{U} \boldsymbol{\Sigma} \mathbf{V}^\top$ be the truncated rank- r SVD of \mathbf{M}_{left} . We also let $\mathbf{H} = \text{sgn}(\mathbf{U}^\top \mathbf{U}^*)$ be the optimal rotation matrix that aligns \mathbf{U} and \mathbf{U}^* . We have the following results for the subspace estimation error $\mathbf{U} \mathbf{H} - \mathbf{U}^*$.

Lemma 10. Under the conditions of [Lemma 1](#), we have the decomposition

$$\mathbf{U} \mathbf{H} - \mathbf{U}^* = \mathbf{Z} + \boldsymbol{\Psi}, \quad \text{where } \mathbf{Z} := \mathbf{E}_{\text{left}} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1},$$

and there exists universal constant $C_6 > 0$ such that the matrix $\boldsymbol{\Psi}$ satisfies the bounds

$$\|\boldsymbol{\Psi}_{i,\cdot}\|_2 \leq C_6 \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^{*2} T_1 / T} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{(\sigma_r^* \sqrt{T_1 / T})^3} + \left(\frac{\sigma^2 (N + T_1)}{\sigma_r^{*2} T_1 / T} + \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1 / T}} \right) \|\mathbf{U}_{i,\cdot}^*\|_2 \right] \quad (\text{C.3})$$

with probability at least $1 - O((N + T)^{-9})$, for all $i \in [N]$. As a consequence, with probability at least $1 - O((N + T)^{-9})$, the row-wise error bound

$$\|\mathbf{U}_{i,\cdot} \mathbf{H} - \mathbf{U}_{i,\cdot}^*\|_2 \leq C_6 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1 / T}} + C_6 \frac{\sigma^2 (N + T_1)}{\sigma_r^{*2} T_1 / T} \|\mathbf{U}_{i,\cdot}^*\|_2 \quad (\text{C.4})$$

holds for all $i \in [N]$, and the submatrix \mathbf{U}_1 satisfies the following spectral norm error bound

$$\|\mathbf{U}_1 \mathbf{H} - \mathbf{U}_1^*\| \leq C_6 \frac{\sigma^2 T}{\sigma_r^{*2}} \sqrt{\frac{N_1}{N}} + C_6 \frac{\sigma \sqrt{N_1}}{\sigma_r^* \sqrt{T_1 / T}}. \quad (\text{C.5})$$

As a byproduct, the spectrum of \mathbf{U}_1 is upper and lower bounded by

$$\frac{c_\ell}{2} \frac{N_1}{N} \mathbf{I}_r \preceq \mathbf{U}_1^\top \mathbf{U}_1 \preceq 2c_u \frac{N_1}{N} \mathbf{I}_r. \quad (\text{C.6})$$

See [Appendix C.5](#) for the proof.

We also need the following lemma that studies the proximity of $\mathbf{U} \boldsymbol{\Sigma}$ and $\mathbf{U}^* \boldsymbol{\Sigma}^*$. While it looks similar to the previous one, it is crucial to establish our result that is condition number free (i.e. does not depend on the condition number of $\boldsymbol{\Sigma}^*$).

Lemma 11. Under the conditions of [Lemma 1](#), we have the decomposition

$$\mathbf{U} \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}^* \boldsymbol{\Sigma}^* = \mathbf{E}_{\text{left}} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} + \boldsymbol{\Delta} = \mathbf{Z} \boldsymbol{\Sigma}^* + \boldsymbol{\Delta},$$

where $\mathbf{R} \in \mathbb{R}^{r \times r}$ is some invertible matrix and $\mathbf{\Delta}$ is some residual matrix satisfying: there exists some universal constant $C_7 > 0$ such that with probability at least $1 - O((N+T)^{-9})$, for all $i \in [N]$,

$$\|\mathbf{\Delta}_{i,\cdot}\|_2 \leq C_7 \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^* T_1 / T} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} (T_1 / T)^{3/2}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1 / T} \|\mathbf{U}_{i,\cdot}^*\|_2 \right]. \quad (\text{C.7})$$

As a consequence, with probability at least $1 - O((N+T)^{-9})$, the row-wise error bound

$$\|\mathbf{U}_{i,\cdot} \mathbf{\Sigma} \mathbf{R} - \mathbf{U}_{i,\cdot}^* \mathbf{\Sigma}^*\|_2 \leq C_7 \left[\frac{\sigma \sqrt{r + \xi_{N,T}}}{\sqrt{T_1 / T}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1 / T} \|\mathbf{U}_{i,\cdot}^*\|_2 \right] \quad (\text{C.8})$$

holds for all $i \in [N]$, and the submatrix $\mathbf{U}_1 \mathbf{\Sigma}$ satisfies the following spectral norm error bound

$$\|\mathbf{U}_1 \mathbf{\Sigma} \mathbf{R} - \mathbf{U}_1^* \mathbf{\Sigma}^*\| \leq C_7 \left[\frac{\sigma \sqrt{N_1}}{\sqrt{T_1 / T}} + \frac{\sigma^2 T}{\sigma_r^*} \sqrt{\frac{N_1}{N}} \right]. \quad (\text{C.9})$$

We also have the following spectral norm bound on the submatrix $\mathbf{\Delta}_1 := \mathbf{\Delta}_{1:N_1}$,

$$\|\mathbf{\Delta}_1\| \leq C_7 \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1 / T} \sqrt{\frac{N_1}{N}}. \quad (\text{C.10})$$

See [Appendix C.6](#) for the proof.

The next lemma characterizes the error associated with estimating \mathbf{M}_b^* .

Lemma 12. *Under the conditions of [Lemma 1](#), we can decompose*

$$\mathbf{M}_b - \mathbf{M}_b^* = \mathbf{W} + \mathbf{\Phi}, \quad \text{where } \mathbf{W} := \mathbf{E}_a \mathbf{V}_1^* \mathbf{V}_2^{*\top} + \mathbf{E}_b \mathbf{V}_2^* \mathbf{V}_2^{*\top} + \mathbf{U}_1^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_b.$$

Here $\mathbf{\Phi}$ is some residual matrix satisfying: there exists some universal constant $C_8 > 0$ such that with probability at least $1 - O((N+T)^{-9})$, for all $t \in [T]$,

$$\|\mathbf{\Phi}_{\cdot,t}\|_2 \leq C_8 \left[\frac{\sigma^2 \sqrt{N} (r + \xi_{N,T})}{\sigma_r^*} + \frac{\sigma^3 T \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} N_1 / N} + \left(\frac{\sigma^2 (N_1 + T)}{\sigma_r^* \sqrt{N_1 / N}} + \sigma \sqrt{r + \xi_{N,T}} \right) \|\mathbf{V}_{t,\cdot}^*\|_2 \right].$$

As a consequence, with probability at least $1 - O((N+T)^{-9})$, the column-wise error bound

$$\|(\mathbf{M}_b - \mathbf{M}_b^*)_{\cdot,t}\|_2 \leq C_8 \sigma \sqrt{r + \xi_{N,T}} + C_8 \left(\frac{\sigma^2 T}{\sigma_r^* \sqrt{N_1 / N}} + \sigma \sqrt{N_1} \right) \|\mathbf{V}_{t,\cdot}^*\|_2$$

holds for all $t \in [T]$.

See [Appendix C.7](#) for the proof.

Finally, we collect some results in the next lemma that will be useful in later proof.

Lemma 13. *Under the conditions of [Lemma 1](#), there exists universal constant $C_9 > 0$ such that with probability at least $1 - O((N+T)^{-10})$, we have*

$$\|\mathbf{U}_1^{*\top} (\mathbf{U}_1 \mathbf{H} - \mathbf{U}_1^*)\| \leq C_9 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1 / T}} \sqrt{\frac{N_1}{N}} + C_9 \frac{\sigma^2 (N + T_1) N_1}{\sigma_r^{*2} T_1 / T} \frac{N_1}{N} := \delta_{\text{prod}}. \quad (\text{C.11})$$

As a direct consequence, the following two bounds also hold:

$$\|[(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} - (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1}\| \leq \delta_{\text{prod}} \frac{N^2}{N_1^2}, \quad (\text{C.12})$$

$$\|[(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} (\mathbf{U}_1\mathbf{H})^\top - (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top}\| \leq C_9 \frac{\sigma^2 T}{\sigma_r^{\star 2}} \sqrt{\frac{N}{N_1}} + C_9 \frac{\sigma \sqrt{N_1}}{\sigma_r^* \sqrt{T_1/T}} \frac{N}{N_1}. \quad (\text{C.13})$$

See [Appendix C.8](#) for the proof.

C.2 Proof of [Lemma 2](#)

It is convenient to define the matrix

$$\mathbf{L} := \mathbf{U}_2\mathbf{H} [(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} (\mathbf{U}_1\mathbf{H})^\top - \mathbf{U}_2^* (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top}. \quad (\text{C.14})$$

Recall the decomposition $\mathbf{U}_2\mathbf{H} = \mathbf{U}_2^* + \mathbf{Z}_2 + \mathbf{\Psi}_2$ from [Lemma 10](#), we can write

$$\mathbf{L} = \mathbf{Z}_2 (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} + \mathbf{\Delta}_L \quad (\text{C.15})$$

where we define

$$\mathbf{\Delta}_L := \mathbf{\Psi}_2 (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} + \mathbf{U}_2\mathbf{H} \{ [(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} (\mathbf{U}_1\mathbf{H})^\top - (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \}.$$

The matrix $\mathbf{\Delta}_L$ can be bounded as follows: for each $i \in [N_2]$, we have

$$\begin{aligned} \|(\mathbf{\Delta}_L)_{i,\cdot}\|_2 &\leq \|(\mathbf{\Psi}_2)_{i,\cdot}\|_2 \sigma_r^{-1} (\mathbf{U}_1^*) + \|\mathbf{U}_{i,\cdot}\|_2 \| [(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} (\mathbf{U}_1\mathbf{H})^\top - (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \| \\ &\stackrel{(i)}{\leq} \frac{C_6}{c_\ell} \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^{\star 2} T_1/T} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{(\sigma_r^* \sqrt{T_1/T})^3} + \left(\frac{\sigma^2 (N + T_1)}{\sigma_r^{\star 2} T_1/T} + \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \right) \|\mathbf{U}_{i,\cdot}^*\|_2 \right] \sqrt{\frac{N}{N_1}} \\ &\quad + (2 \|\mathbf{U}_{i,\cdot}^*\|_2 + C_6 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}}) \cdot C_9 \left(\frac{\sigma^2 T}{\sigma_r^{\star 2}} \sqrt{\frac{N}{N_1}} + \frac{\sigma \sqrt{N_1}}{\sigma_r^* \sqrt{T_1/T}} \frac{N}{N_1} \right) \quad (\text{C.16}) \\ &\stackrel{(ii)}{\leq} C_L \left(\frac{\sigma^2 \sqrt{r + \xi_{N,T}}}{\sigma_r^{\star 2} \sqrt{T_1/T}} \sqrt{\frac{NT}{N_1}} + \frac{\sigma^2}{\sigma_r^{\star 2}} \frac{NT}{T_1} \sqrt{\frac{r + \xi_{N,T}}{N_1}} \right) + C_L \|\mathbf{U}_{i,\cdot}^*\|_2 \left(\frac{\sigma^2 T}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right) \sqrt{\frac{N}{N_1}}, \end{aligned}$$

where we take $C_L = 2C_6/c_\ell + 2C_6C_9 + 4C_9$. Here (i) follows from [\(4.1\)](#), relations [\(C.3\)](#), [\(C.13\)](#), and a direct consequence of [\(C.4\)](#):

$$\|\mathbf{U}_{i,\cdot}\|_2 \leq \|\mathbf{U}_{i,\cdot}^*\|_2 + \|\mathbf{U}_{i,\cdot}\mathbf{H} - \mathbf{U}_{i,\cdot}^*\| \stackrel{(iv)}{\leq} 2 \|\mathbf{U}_{i,\cdot}^*\|_2 + C_6 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}}; \quad (\text{C.17})$$

steps (ii) and (iii) holds provided that $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ and $\sigma \sqrt{T} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small $c_{\text{noise}} > 0$. We decompose

$$\begin{aligned} \mathbf{A} &= \mathbf{L} (\mathbf{M}_b - \mathbf{M}_b^*) + \mathbf{U}_2^* (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} (\mathbf{M}_b - \mathbf{M}_b^*) \\ &= \mathbf{Z}_2 (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} (\mathbf{W} + \mathbf{\Phi}) + \mathbf{\Delta}_L (\mathbf{M}_b - \mathbf{M}_b^*) + \mathbf{U}_2^* (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} (\mathbf{W} + \mathbf{\Phi}) \\ &= \underbrace{\mathbf{U}_2^* (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \mathbf{E}_b + \mathbf{E}_c \mathbf{V}_1^* (\mathbf{V}_1^{\star\top} \mathbf{V}_1^\star)^{-1} (\mathbf{\Sigma}^*)^{-1} (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \mathbf{E}_b}_{=: \mathbf{A}_1} \\ &\quad + \underbrace{\mathbf{E}_c \mathbf{V}_1^* (\mathbf{V}_1^{\star\top} \mathbf{V}_1^\star)^{-1} (\mathbf{\Sigma}^*)^{-1} (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \mathbf{E}_{\text{upper}} \mathbf{V}^* \mathbf{V}_2^{\star\top}}_{=: \mathbf{A}_2} + \underbrace{\mathbf{U}_2^* (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \mathbf{E}_{\text{upper}} \mathbf{V}^* \mathbf{V}_2^{\star\top}}_{=: \mathbf{A}_3} \\ &\quad + \underbrace{\mathbf{\Delta}_L (\mathbf{M}_b - \mathbf{M}_b^*)}_{=: \mathbf{A}_4} + \underbrace{(\mathbf{U}_2^* + \mathbf{Z}_2) (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \mathbf{\Phi}}_{=: \mathbf{A}_5}, \end{aligned}$$

where the second relation follows from [Lemma 12](#) and [\(C.15\)](#). The matrix \mathbf{A}_1 appears in our desired decomposition [\(5.8\)](#). Then we bound the entrywise magnitude of the other four matrices.

We start with \mathbf{A}_2 . Notice that \mathbf{E}_c and $\mathbf{E}_{\text{upper}}$ are independent, then conditional on $\mathbf{E}_{\text{upper}}$,

$$(\mathbf{A}_2)_{i,t} | \mathbf{E}_{\text{upper}} \sim \mathcal{N}\left(0, \sigma^2 \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_{\text{upper}} \mathbf{V}^* \mathbf{V}_{t,\cdot}^{*\top}\|_2^2\right).$$

Therefore, with probability at least $1 - O((N + T)^{-10})$, we have

$$\begin{aligned} |(\mathbf{A}_2)_{i,t}| &\leq 5\sigma \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_{\text{upper}} \mathbf{V}^* \mathbf{V}_{t,\cdot}^{*\top}\|_2 \sqrt{\xi_{N,T}} \\ &\stackrel{(i)}{\leq} 5C_g \sigma^2 \sqrt{r + \xi_{N,T}} \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\| \|\mathbf{V}^* \mathbf{V}_{t,\cdot}^{*\top}\| \sqrt{\xi_{N,T}} \\ &\stackrel{(ii)}{\leq} \tilde{C}_2 \frac{\sigma^2}{\sigma_r^*} \sqrt{\frac{NT}{N_1 T_1}} \xi_{N,T} (r + \xi_{N,T}) \|\mathbf{V}_{j,\cdot}^*\|_2, \end{aligned}$$

where $\tilde{C}_2 = 5C_g/c_\ell$. Here (i) follows from [Lemma 19](#), and (ii) holds due to [\(4.1\)](#). Regarding \mathbf{A}_3 , one can check that

$$(\mathbf{A}_3)_{i,t} \sim \mathcal{N}\left(0, \sigma^2 \|\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\|_2^2 \|\mathbf{V}_{t,\cdot}^* \mathbf{V}^{*\top}\|_2^2\right),$$

hence with probability at least $1 - O((N + T)^{-10})$,

$$|(\mathbf{A}_3)_{i,t}| \leq 5\sigma \|\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\|_2 \|\mathbf{V}_{t,\cdot}^* \mathbf{V}^{*\top}\|_2 \sqrt{\xi_{N,T}} \leq \tilde{C}_3 \sigma \sqrt{\frac{N}{N_1}} \xi_{N,T} \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2,$$

where $\tilde{C}_3 = 5/\sqrt{c_\ell}$, and we use [\(4.1\)](#) in the last step. Then we bound \mathbf{A}_4 , where we use the Cauchy–Schwarz inequality to achieve

$$\begin{aligned} |(\mathbf{A}_4)_{i,t}| &\leq \|(\boldsymbol{\Delta}_L)_{i,\cdot}\|_2 \|(\mathbf{M}_b - \mathbf{M}_b^*)_{\cdot,t}\|_2 \\ &\leq \tilde{C}_4 \left(\frac{\sigma^2 \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} \sqrt{T_1/T}} \sqrt{\frac{NT}{N_1}} + \frac{\sigma^2}{\sigma_r^{*2}} \frac{NT}{T_1} \sqrt{\frac{r + \xi_{N,T}}{N_1}} \right) \left[\sigma \sqrt{r + \xi_{N,T}} + \left(\frac{\sigma^2 T}{\sigma_r^* \sqrt{N_1/N}} + \sigma \sqrt{N_1} \right) \|\mathbf{V}_{t,\cdot}^*\|_2 \right] \\ &\quad + \tilde{C}_4 \|\mathbf{U}_{i,\cdot}^*\|_2 \left(\frac{\sigma^2 T}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right) \sqrt{\frac{N}{N_1}} \left[\sigma \sqrt{r + \xi_{N,T}} + \left(\frac{\sigma^2 T}{\sigma_r^* \sqrt{N_1/N}} + \sigma \sqrt{N_1} \right) \|\mathbf{V}_{t,\cdot}^*\|_2 \right] \\ &\leq \tilde{C}_4 \left(\frac{\sigma^3 T}{\sigma_r^{*2}} \sqrt{\frac{N}{N_1 T_1}} + \frac{\sigma^3 \sqrt{N_1}}{\sigma_r^{*2}} \frac{NT}{N_1 T_1} \right) (r + \xi_{N,T}) + \tilde{C}_4 \left[\frac{\sigma^2 \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{N_1/N}} \sqrt{\frac{NT}{T_1}} + \frac{\sigma^3 T \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} \sqrt{N_1/N}} \right] \|\mathbf{U}_{i,\cdot}^*\|_2 \\ &\quad + \tilde{C}_4 \left(\frac{\sigma^2}{\sigma_r^*} \frac{NT}{N_1} + \sigma \sqrt{N} \right) \left[\frac{\sigma^2 \sqrt{(N + T_1)(r + \xi_{N,T})}}{\sigma_r^{*2} T_1/T} \|\mathbf{V}_{t,\cdot}^*\|_2 + \left(\frac{\sigma^2 T}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right) \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2 \right]. \end{aligned}$$

where $\tilde{C}_4 = C_L C_8$. Here we use the bounds [\(C.16\)](#) and [Lemma 12](#). Finally we use [Lemmas 10](#)

and 12 to reach

$$\begin{aligned}
|(\mathbf{A}_5)_{i,t}| &\leq (\|\mathbf{U}_{i,\cdot}^*\|_2 + \|\mathbf{E}_{i,\cdot} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1}\|_2) \sigma_r^{-1} (\mathbf{U}_1^*) \|\Phi_{\cdot,t}\|_2 \\
&\stackrel{(i)}{\leq} \left(\|\mathbf{U}_{i,\cdot}^*\|_2 + \frac{C_g}{\sqrt{c_\ell}} \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^*} \sqrt{\frac{T}{T_1}} \right) \frac{1}{\sqrt{c_\ell}} \sqrt{\frac{N}{N_1}} \|\Phi_{\cdot,t}\|_2 \\
&\stackrel{(ii)}{\leq} \tilde{C}_5 \left(\frac{\sigma^3 \sqrt{N}}{\sigma_r^{*2}} + \frac{\sigma^4}{\sigma_r^{*3}} \frac{NT}{N_1} \right) \sqrt{\frac{NT}{N_1 T_1}} (r + \xi_{N,T}) + \tilde{C}_5 \left[\frac{\sigma^2 \sqrt{N(r + \xi_{N,T})}}{\sigma_r^* \sqrt{N_1/N}} + \frac{\sigma^3 T \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} (N_1/N)^{3/2}} \right] \|\mathbf{U}_{i,\cdot}^*\|_2 \\
&\quad + \tilde{C}_5 \left[\frac{\sigma^3 (N_1 + T) \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} \sqrt{N_1/N}} + \frac{\sigma^2 (r + \xi_{N,T})}{\sigma_r^*} \right] \sqrt{\frac{NT}{N_1 T_1}} \|\mathbf{V}_{t,\cdot}^*\|_2 \\
&\quad + \tilde{C}_5 \left[\frac{\sigma^2 (N_1 + T)}{\sigma_r^* N_1/N} + \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sqrt{N_1/N}} \right] \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2,
\end{aligned} \tag{C.18}$$

where $\tilde{C}_5 = 4/\sqrt{c_\ell} + 4C_8 C_g / c_\ell$. Here (i) holds due to an application of Lemma 19 and (4.1):

$$\|\mathbf{E}_{i,\cdot} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1}\|_2 \leq C_g \sigma \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1}\| \sqrt{r + \xi_{N,T}} \leq \frac{C_g}{\sqrt{c_\ell}} \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^*} \sqrt{\frac{T}{T_1}},$$

while (ii) follows from Lemma 12 and holds true when $c_{\text{noise}} > 0$ is sufficiently small.

Taking the above bounds on \mathbf{A}_2 to \mathbf{A}_5 collectively, we conclude that $\mathbf{A} = \mathbf{A}_1 + \boldsymbol{\xi}_A$, where

$$\begin{aligned}
|(\boldsymbol{\xi}_A)_{i,t}| &\leq |(\mathbf{A}_2)_{i,t}| + |(\mathbf{A}_3)_{i,t}| + |(\mathbf{A}_4)_{i,t}| + |(\mathbf{A}_5)_{i,t}| \\
&\leq C_A \frac{\sigma^3 T}{\sigma_r^{*2}} \sqrt{\frac{N}{N_1 T_1}} (r + \xi_{N,T}) + C_A \frac{\sigma^3 \sqrt{N_1}}{\sigma_r^{*2}} \frac{NT}{N_1 T_1} (r + \xi_{N,T}) + C_A \frac{\sigma^4}{\sigma_r^{*3}} \sqrt{\frac{NT}{N_1 T_1}} \frac{NT}{N_1} (r + \xi_{N,T}) \\
&\quad + C_A \left[\frac{\sigma^2 \sqrt{N_1} (r + \xi_{N,T})}{\sigma_r^* \sqrt{T_1/T}} \frac{N}{N_1} + \frac{\sigma^3 T \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} (N_1/N)^{3/2}} \right] \|\mathbf{U}_{i,\cdot}^*\|_2 \\
&\quad + C_A \left[\frac{\sigma^2}{\sigma_r^*} \sqrt{\frac{NT(r + \xi_{N,T})}{N_1 T_1}} + \frac{\sigma^2 \sqrt{(N + T_1)}}{\sigma_r^{*2} T_1/T} \left(\frac{\sigma^2 NT}{\sigma_r^* N_1} + \sigma \sqrt{N} \right) + \frac{\sigma^3 NT}{\sigma_r^{*2} N_1} \sqrt{\frac{T}{T_1}} \right] \|\mathbf{V}_{t,\cdot}^*\|_2 \sqrt{r + \xi_{N,T}} \\
&\quad + C_A \left[\sigma \sqrt{\frac{N}{N_1}} (r + \xi_{N,T}) + \frac{\sigma^2 NT}{\sigma_r^* N_1} + \left(\frac{\sigma^2 T}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right) \sigma \sqrt{N} \right] \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2.
\end{aligned}$$

Here we take $C_A = 2 \sum_{k=2}^5 \tilde{C}_k$ and the above relation holds as long as $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ and $\sigma \sqrt{T} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small constant $c_{\text{noise}} > 0$.

C.3 Proof of Lemma 3

We begin with the decomposition

$$\begin{aligned}
\mathbf{B} &= \mathbf{U}_2 (\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_1^\top (\mathbf{U}_1 \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_1^* \boldsymbol{\Sigma}^*) \mathbf{V}_2^{*\top} + (\mathbf{U}_2 \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_2^* \boldsymbol{\Sigma}^*) \mathbf{V}_2^{*\top} \\
&\stackrel{(i)}{=} \mathbf{U}_2^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} (\mathbf{U}_1 \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_1^* \boldsymbol{\Sigma}^*) \mathbf{V}_2^{*\top} + \mathbf{L} (\mathbf{U}_1 \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_1^* \boldsymbol{\Sigma}^*) \mathbf{V}_2^{*\top} + (\mathbf{U}_2 \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_2^* \boldsymbol{\Sigma}^*) \mathbf{V}_2^{*\top} \\
&\stackrel{(ii)}{=} \underbrace{\mathbf{U}_2^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_a \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_2^{*\top}}_{=: \mathbf{B}_1} + \underbrace{\mathbf{L} (\mathbf{U}_1 \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_1^* \boldsymbol{\Sigma}^*) \mathbf{V}_2^{*\top}}_{=: \mathbf{B}_2} \\
&\quad + \underbrace{(\mathbf{U}_2^* + \mathbf{Z}_2) (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \boldsymbol{\Delta}_1 \mathbf{V}_2^{*\top}}_{=: \mathbf{B}_3} + \underbrace{\mathbf{E}_c \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_2^*}_{=: \mathbf{B}_4} + \underbrace{\boldsymbol{\Delta}_2 \mathbf{V}_2^*}_{=: \mathbf{B}_5}.
\end{aligned}$$

Here step (i) follows from the definition (C.14) of \mathbf{L} , while step (ii) follows from Lemma 11.

We first bound \mathbf{B}_1 , whose entries follow the Gaussian distribution

$$(\mathbf{B}_1)_{i,t} \sim \mathcal{N}\left(0, \sigma^2 \|\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\|_2^2 \|\mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top}\|_2^2\right).$$

Therefore we can use (4.1) to show that with probability at least $1 - O((N+T)^{-10})$,

$$|(\mathbf{B}_1)_{i,t}| \leq 5\sigma \|\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\|_2 \|\mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_1^{*\top}\|_2 \sqrt{\xi_{N,T}} \leq \tilde{C}_1 \sigma \sqrt{\frac{NT}{N_1 T_1}} \xi_{N,T} \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2,$$

where $\tilde{C}_1 = 5/c_\ell$. Regarding \mathbf{C}_2 , we first derive a bound for $\|\mathbf{L}_{i,\cdot}\|_2$. Starting from (C.15), we have

$$\begin{aligned} \|\mathbf{L}_{i,\cdot}\|_2 &\leq \|\mathbf{U}_{i,\cdot} \mathbf{H} - \mathbf{U}_{i,\cdot}^*\|_2 \sigma_r^{-1} (\mathbf{U}_1^*) + \|\mathbf{U}_{i,\cdot}\|_2 \left\| [(\mathbf{U}_1 \mathbf{H})^\top \mathbf{U}_1 \mathbf{H}]^{-1} (\mathbf{U}_1 \mathbf{H})^\top - (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \right\| \\ &\stackrel{(i)}{\leq} C_6 \left(\frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2 (N+T_1)}{\sigma_r^{*2} T_1/T} \|\mathbf{U}_{i,\cdot}^*\|_2 \right) \cdot c_\ell^{-1/2} \sqrt{\frac{N}{N_1}} \\ &\quad + \left(2\|\mathbf{U}_{i,\cdot}^*\|_2 + C_6 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \right) \cdot C_9 \left(\frac{\sigma^2 T}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right) \sqrt{\frac{N}{N_1}} \\ &\stackrel{(ii)}{\leq} C'_L \|\mathbf{U}_{i,\cdot}^*\|_2 \left(\frac{\sigma^2 T}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right) \sqrt{\frac{N}{N_1}} + C'_L \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \sqrt{\frac{N}{N_1}}, \end{aligned} \quad (\text{C.19})$$

where we take $C'_L = 2C_6/\sqrt{c_\ell} + 4C_9 + 2C_6 C_9$. Here (i) follows from (C.3), (C.17), (4.1) and (C.13), while (ii) holds provided that $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ and $\sigma \sqrt{T} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small $c_{\text{noise}} > 0$. Apply the bounds (C.19) and (C.9) to achieve

$$\begin{aligned} |(\mathbf{B}_2)_{i,t}| &\leq \|\mathbf{L}_{i,\cdot}\|_2 \|\mathbf{U}_1 \Sigma \mathbf{R} - \mathbf{U}_1^* \Sigma^*\| \|\mathbf{V}_{t,\cdot}^*\|_2 \\ &\leq \tilde{C}_2 \left(\frac{\sigma^2 T}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right)^2 \sigma_r^* \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2 + \tilde{C}_2 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sqrt{T_1/T}} \left(\frac{\sigma^2 T}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{\frac{NT}{T_1}} \right) \|\mathbf{V}_{t,\cdot}^*\|_2, \end{aligned}$$

where $\tilde{C}_2 = C'_L C_7$. Then we move on to bounding \mathbf{B}_3 , where we have

$$\begin{aligned} |(\mathbf{B}_3)_{i,t}| &\leq \|\mathbf{U}_{i,\cdot}^* + \mathbf{Z}_{i,\cdot}\|_2 \sigma_r^{-1} (\mathbf{U}_1^*) \|\Delta_1\| \|\mathbf{V}_{t,\cdot}^*\|_2 \\ &\stackrel{(i)}{\leq} \left(\|\mathbf{U}_{i,\cdot}^*\|_2 + \frac{C_g}{\sqrt{c_\ell}} \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^*} \sqrt{\frac{T}{T_1}} \right) \cdot c_\ell^{-1/2} \sqrt{\frac{N}{N_1}} \cdot C_7 \frac{\sigma^2 (N+T_1)}{\sigma_r^* T_1/T} \sqrt{\frac{N_1}{N}} \|\mathbf{V}_{t,\cdot}^*\|_2 \\ &\stackrel{(ii)}{\leq} \tilde{C}_3 \frac{\sigma^2 (N+T_1)}{\sigma_r^* T_1/T} \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2 + \tilde{C}_3 \frac{\sigma^3 (N+T_1)}{\sigma_r^{*2} (T_1/T)^{3/2}} \sqrt{r + \xi_{N,T}} \|\mathbf{V}_{t,\cdot}^*\|_2. \end{aligned}$$

Here (i) follows from (C.10), (4.1), and also uses a similar upper bound for $\|(\mathbf{U}_2^* + \mathbf{Z}_2)_{i,\cdot}\|_2$ as in (C.18); while (ii) holds true if we take $\tilde{C}_3 = C_g C_7 / c_\ell + C_7 / \sqrt{c_\ell}$. The matrix \mathbf{B}_4 appears in our desired decomposition (5.9). Finally we invoke Lemma 11 to show that

$$|(\mathbf{B}_5)_{i,t}| \leq C_7 \left(\frac{\sigma^2 \sqrt{T_1 (r + \xi_{N,T})}}{\sigma_r^* T_1/T} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} (T_1/T)^{3/2}} \right) \|\mathbf{V}_{t,\cdot}^*\|_2 + C_7 \frac{\sigma^2 (N+T_1)}{\sigma_r^* T_1/T} \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2.$$

Taking the bounds on $\mathbf{B}_1, \mathbf{B}_2, \mathbf{B}_3$ and \mathbf{B}_5 collectively yields $\mathbf{B} = \mathbf{B}_4 + \boldsymbol{\xi}_B$ where $\boldsymbol{\xi}_B = \mathbf{B}_1 + \mathbf{B}_2 + \mathbf{B}_3 + \mathbf{B}_5$. The residual matrix $\boldsymbol{\xi}_B$ satisfies

$$\begin{aligned} |(\boldsymbol{\xi}_B)_{i,t}| &\leq |(\mathbf{B}_1)_{i,t}| + |(\mathbf{B}_2)_{i,t}| + |(\mathbf{B}_3)_{i,t}| + |(\mathbf{B}_5)_{i,t}| \\ &\leq C_B \frac{\sigma\sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \left(\sigma\sqrt{T} + \sigma\sqrt{\frac{NT}{T_1}} \right) \|\mathbf{V}_{t,\cdot}^*\|_2 + C_B \left(\frac{\sigma^2(N + T_1)}{\sigma_r^* T_1/T} + \sigma\sqrt{\frac{NT\xi_{N,T}}{N_1 T_1}} \right) \|\mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{V}_{t,\cdot}^*\|_2, \end{aligned}$$

where we take $C_B = 2\tilde{C}_1 + 2\tilde{C}_2 + 2\tilde{C}_3 + 2C_7$. The above relation holds provided that $\sigma\sqrt{NT/T_1} \leq c_{\text{noise}}\sigma_r^*$ and $\sigma\sqrt{T} \leq c_{\text{noise}}\sigma_r^*$ for some sufficiently small $c_{\text{noise}} > 0$.

C.4 Proof of Lemma 9

It suffices to upper and lower bound the r -th eigenvalue of $\mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top} = \mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top} \mathbf{V}_1^* \boldsymbol{\Sigma}^* \mathbf{U}^{*\top}$. On the one hand, by the max-min characterization for eigenvalues, we know that

$$\begin{aligned} \lambda_r(\mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top}) &= \max_{S \subset \mathbb{R}^N: \dim(S)=r} \min_{\mathbf{x} \in S, \|\mathbf{x}\|_2=1} \mathbf{x}^\top \mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top} \mathbf{x} \stackrel{(i)}{\geq} \min_{\boldsymbol{\beta} \in \mathbb{R}^r: \|\boldsymbol{\beta}\|_2=1} \boldsymbol{\beta}^\top \mathbf{U}^{*\top} \mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top} \mathbf{U}^* \boldsymbol{\beta} \\ &= \min_{\boldsymbol{\beta} \in \mathbb{R}^r: \|\boldsymbol{\beta}\|_2=1} \boldsymbol{\beta}^\top \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top} \mathbf{V}_1^* \boldsymbol{\Sigma}^* \boldsymbol{\beta} \stackrel{(ii)}{\geq} c_\ell \frac{T_1}{T} \min_{\boldsymbol{\beta} \in \mathbb{R}^r: \|\boldsymbol{\beta}\|_2=1} \|\boldsymbol{\Sigma}^* \boldsymbol{\beta}\|_2^2 = c_\ell \frac{T_1}{T} \sigma_r^{*2}. \quad (\text{C.20}) \end{aligned}$$

Here step (i) holds by taking the subspace S in the first line to be the subspace spanned by the columns of \mathbf{U}^* , which allows one to parameterize any unit vector $\mathbf{x} \in S$ with $\mathbf{U}^* \boldsymbol{\beta}$ for some unit vector $\boldsymbol{\beta} \in \mathbb{R}^r$; and step (ii) follows again from (4.1). On the other hand, by the min-max characterization for eigenvalues, we know that

$$\lambda_r(\mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top}) = \min_{S \subset \mathbb{R}^N: \dim(S)=N-r+1} \max_{\mathbf{x} \in S, \|\mathbf{x}\|_2=1} \mathbf{x}^\top \mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top} \mathbf{x}.$$

By taking S to be the subspace spanned by the orthogonal complement of \mathbf{U}^* and the singular vector $\mathbf{U}_{\cdot,r}^*$ associated with the minimum singular value σ_r^* , namely

$$S = \text{span} \{ \mathbf{U}_{\cdot,1}^*, \mathbf{U}_{\cdot,r}^* \},$$

we can express any unit vector \mathbf{x} in S as

$$\mathbf{x} = (\mathbf{I} - \mathbf{U}^* \mathbf{U}^{*\top}) \boldsymbol{\beta} + \alpha \mathbf{U}_{\cdot,r}^*$$

for some vector $\boldsymbol{\beta} \in \mathbb{R}^N$ and scalar $\alpha \in \mathbb{R}$ that satisfy $\|(\mathbf{I} - \mathbf{U}^* \mathbf{U}^{*\top}) \boldsymbol{\beta}\|_2^2 + \alpha^2 = 1$. We can also compute

$$\begin{aligned} \mathbf{x}^\top \mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top} \mathbf{x} &= [(\mathbf{I} - \mathbf{U}^* \mathbf{U}^{*\top}) \boldsymbol{\beta} + \alpha \mathbf{U}_{\cdot,r}^*]^\top \mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top} \mathbf{V}_1^* \boldsymbol{\Sigma}^* \mathbf{U}^{*\top} [(\mathbf{I} - \mathbf{U}^* \mathbf{U}^{*\top}) \boldsymbol{\beta} + \alpha \mathbf{U}_{\cdot,r}^*]^\top \\ &= \alpha^2 \mathbf{U}_{\cdot,r}^{*\top} \mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top} \mathbf{V}_1^* \boldsymbol{\Sigma}^* \mathbf{U}^{*\top} \mathbf{U}_{\cdot,r}^* = \alpha^2 \sigma_r^{*2} \mathbf{e}_r^\top \mathbf{V}_1^{*\top} \mathbf{V}_1^* \mathbf{e}_r \leq \alpha^2 c_u \frac{T_1}{T} \sigma_r^{*2}. \end{aligned}$$

Here \mathbf{e}_r is the r -th canonical basis vector in \mathbb{R}^r , and we use (4.1) in the last relation. Then we can bound

$$\lambda_r(\mathbf{M}_{\text{left}}^* \mathbf{M}_{\text{left}}^{*\top}) \leq \max_{\alpha \in [-1,1]} \alpha^2 c_u \frac{T_1}{T} \sigma_r^{*2} = c_u \frac{T_1}{T} \sigma_r^{*2}. \quad (\text{C.21})$$

Taking equations (C.20) and (C.21) together gives the desired result (C.2a). The remaining relation (C.2b) can be proved in a similar way.

C.5 Proof of Lemma 10

We apply Proposition 2 to the observation $\mathbf{M}_{\text{left}} = \mathbf{M}_{\text{left}}^* + \mathbf{E}_{\text{left}}$, where we assume that the SVD of $\mathbf{M}_{\text{left}}^*$ is given by $\mathbf{U}_{\text{left}}^* \boldsymbol{\Sigma}_{\text{left}}^* \mathbf{V}_{\text{left}}^{*\top}$. In order to ensure that the condition of Proposition 2 is satisfied, we check that

$$\sigma \sqrt{\max\{N, T_1\}} \leq c_{\text{noise}} \sqrt{T_1/T} \stackrel{(i)}{\leq} c_n \sqrt{c_\ell T_1/T} \stackrel{(ii)}{\leq} c_n \sigma_r(\mathbf{M}_{\text{left}}^*).$$

Here (i) holds as long as $c_{\text{noise}} \leq c_n \sqrt{c}$, while (ii) follows from Lemma 9. We take $\mathcal{I} = \{i\}$ for any $i \in [N]$. Then (F.2a) tells us that, with probability at least $1 - O((N+T)^{-9})$,

$$\mathbf{U} \text{sgn}(\mathbf{U}^\top \mathbf{U}_{\text{left}}^*) - \mathbf{U}_{\text{left}}^* = \mathbf{E}_{\text{left}} \mathbf{V}_{\text{left}}^* (\boldsymbol{\Sigma}_{\text{left}}^*)^{-1} + \boldsymbol{\Psi}_{\text{left}} \quad (\text{C.22})$$

where for each $i \in [N]$,

$$\begin{aligned} \|(\boldsymbol{\Psi}_{\text{left}})_{i,\cdot}\|_2 &\leq c_1 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r(\mathbf{M}_{\text{left}}^*)} \left(\frac{\sigma \sqrt{T_1}}{\sigma_r(\mathbf{M}_{\text{left}}^*)} + \frac{\sigma^2 N}{\sigma_r^2(\mathbf{M}_{\text{left}}^*)} \right) + c_1 \left(\frac{\sigma^2 (N + T_1)}{\sigma_r^2(\mathbf{M}_{\text{left}}^*)} + \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r(\mathbf{M}_{\text{left}}^*)} \right) \|(\mathbf{U}_{\text{left}}^*)_{i,\cdot}\|_2 \\ &\leq c_1 c_\ell^{-3/2} \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^{*2} T_1/T} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{(\sigma_r^* \sqrt{T_1/T})^3} + \left(\frac{\sigma^2 (N + T_1)}{\sigma_r^{*2} T_1/T} + \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \right) \|U_{i,\cdot}^*\|_2 \right]. \end{aligned}$$

Here we makes use of Lemma 9. Since $\mathbf{U}_{\text{left}}^*$ and \mathbf{U}^* have the same column subspace, there exists a rotation matrix $\mathbf{O}_{\text{left}} \in \mathcal{O}^{r \times r}$ such that $\mathbf{U}_{\text{left}}^* \mathbf{O}_{\text{left}} = \mathbf{U}^*$. In addition, since $\mathbf{V}_{\text{left}}^*$ and \mathbf{V}_1^* have the same column subspace, there exists an invertible matrix $\mathbf{R}_{\text{left}} \in \mathbb{R}^{r \times r}$ such that $\mathbf{V}_{\text{left}}^* \mathbf{R}_{\text{left}} = \mathbf{V}_1^*$. Note that \mathbf{R}_{left} is in general not a rotation matrix because \mathbf{V}_1^* does not necessarily have orthonormal columns. We can express $\mathbf{M}_{\text{left}}^*$ in two ways, either $\mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top}$ or $\mathbf{U}_{\text{left}}^* \boldsymbol{\Sigma}_{\text{left}}^* \mathbf{V}_{\text{left}}^{*\top}$, which further gives

$$\boldsymbol{\Sigma}_{\text{left}}^* = \mathbf{U}_{\text{left}}^{*\top} \mathbf{M}_{\text{left}}^* \mathbf{V}_{\text{left}}^* = \mathbf{U}_{\text{left}}^{*\top} \mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^* = \mathbf{O}_{\text{left}} \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*. \quad (\text{C.23})$$

Notice that $\mathbf{H} = \text{sgn}(\mathbf{U}^\top \mathbf{U}^*) = \text{sgn}(\mathbf{U}^\top \mathbf{U}_{\text{left}}^*) \mathbf{O}_{\text{left}}$. Then we have

$$\begin{aligned} \mathbf{U} \mathbf{H} - \mathbf{U}^* &= \mathbf{U} \text{sgn}(\mathbf{U}^\top \mathbf{U}_{\text{left}}^*) \mathbf{O}_{\text{left}} - \mathbf{U}^* = \mathbf{E}_{\text{left}} \mathbf{V}_{\text{left}}^* (\boldsymbol{\Sigma}_{\text{left}}^*)^{-1} \mathbf{O}_{\text{left}} + \boldsymbol{\Psi}_{\text{left}} \mathbf{O}_{\text{left}} \\ &\stackrel{(i)}{=} \mathbf{E}_{\text{left}} \mathbf{V}_{\text{left}}^* (\mathbf{O}_{\text{left}} \boldsymbol{\Sigma}^* \mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1} \mathbf{O}_{\text{left}} + \boldsymbol{\Psi}_{\text{left}} \mathbf{O}_{\text{left}} \\ &\stackrel{(ii)}{=} \mathbf{E}_{\text{left}} \mathbf{V}_{\text{left}}^* (\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} + \boldsymbol{\Psi}_{\text{left}} \mathbf{O}_{\text{left}} \\ &= \mathbf{E}_{\text{left}} \mathbf{V}_1^* \mathbf{R}_{\text{left}}^{-1} (\mathbf{V}_1^{*\top} \mathbf{V}_1^* \mathbf{R}_{\text{left}}^{-1})^{-1} (\boldsymbol{\Sigma}^*)^{-1} + \boldsymbol{\Psi}_{\text{left}} \mathbf{O}_{\text{left}} \\ &= \mathbf{E}_{\text{left}} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} + \boldsymbol{\Psi}_{\text{left}} \mathbf{O}_{\text{left}}. \end{aligned}$$

Here (i) follows from (C.22), and (ii) follows from (C.23). Note that for each $i \in [N]$, we have $\|(\boldsymbol{\Psi}_{\text{left}})_{i,\cdot} \mathbf{O}_{\text{left}}\|_2 = \|(\boldsymbol{\Psi}_{\text{left}})_{i,\cdot}\|_2$, therefore we have established the desired bound (C.3) by letting

$$\mathbf{Z} := \mathbf{E}_{\text{left}} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} (\boldsymbol{\Sigma}^*)^{-1} \quad \text{and} \quad \boldsymbol{\Psi} := \boldsymbol{\Psi}_{\text{left}} \mathbf{O}_{\text{left}},$$

and by taking $C_6 \geq c_1 c_\ell^{-3/2}$. Next, (F.3a) tells us that with probability at least $1 - O((N+T)^{-9})$, for any $i \in [N]$,

$$\begin{aligned} \|U_{i,\cdot} \mathbf{H} - U_{i,\cdot}^*\|_2 &= \|[U \text{sgn}(\mathbf{U}^\top \mathbf{U}_{\text{left}}^*) - \mathbf{U}_{\text{left}}^*]_{i,\cdot}\|_2 \leq c_1 \frac{\sigma}{\sigma_r(\mathbf{M}_{\text{left}}^*)} \sqrt{r + \xi_{N,T}} + c_1 \frac{\sigma^2 (N + T_1)}{\sigma_r^2(\mathbf{M}_{\text{left}}^*)} \|U_{i,\cdot}^*\|_2 \\ &\leq c_1 c_\ell^{-1} \left[\frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^{*2} T_1/T} \|U_{i,\cdot}^*\|_2 \right]. \end{aligned}$$

In addition, we may apply (F.3a) with $\mathcal{I} = [N_1]$ to obtain that

$$\begin{aligned} \|\mathbf{U}_1 \mathbf{H} - \mathbf{U}_1^*\| &\leq c_1 \left[\frac{\sigma^2 (N + T_1)}{\sigma_r^2 (\mathbf{M}_{\text{left}}^*)} \|\mathbf{U}_1^*\| + \frac{\sigma \sqrt{N_1 + r + \xi_{N,T}}}{\sigma_r (\mathbf{M}_{\text{left}}^*)} \right] \stackrel{(i)}{\leq} c_1 c_\ell^{-1} \left[\frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1 / T} \|\mathbf{U}_1^*\| + \frac{\sigma \sqrt{N_1}}{\sigma_r^* \sqrt{T_1 / T}} \right] \\ &\stackrel{(ii)}{\leq} c_1 c_\ell^{-1} \left[\frac{\sigma^2 T}{\sigma_r^*} \sqrt{\frac{N_1}{N}} + \frac{\sigma \sqrt{N_1}}{\sigma_r^* \sqrt{T_1 / T}} \right] \stackrel{(iii)}{\leq} \frac{\sqrt{c_\ell}}{5} \sqrt{\frac{N_1}{N}} \stackrel{(iv)}{\leq} \frac{\sqrt{c_\ell}}{5} \sigma_r (\mathbf{U}_1^*). \end{aligned}$$

Here (i) makes use of Lemma 9, (ii) and (iii) follow from (4.1) and hold as long as $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ and $\sigma \sqrt{T} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small constant $c_{\text{noise}} > 0$, while (iv) follows from (4.1). This establishes (C.5) by setting $C_6 \geq c_1 c^{-1}$. By Weyl's inequality, we can deduce that

$$\sqrt{\frac{c_\ell}{2} \frac{N_1}{N}} \leq \sigma_r (\mathbf{U}_1^*) - \|\mathbf{U}_1 \mathbf{H} - \mathbf{U}_1^*\| \leq \sigma_r (\mathbf{U}_1) \leq \sigma_1 (\mathbf{U}_1) \leq \sigma_1 (\mathbf{U}_1^*) + \|\mathbf{U}_1 \mathbf{H} - \mathbf{U}_1^*\| \leq \sqrt{2c_u} \frac{N_1}{N}.$$

C.6 Proof of Lemma 11

We apply Proposition 3 to the observation $\mathbf{M}_{\text{left}} = \mathbf{M}_{\text{left}}^* + \mathbf{E}_{\text{left}}$ and use the notation introduced in the last section for the proof of Lemma 10. Similar to the proof of Lemma 10, we can check that the conditions of Proposition 3 are satisfied. We take $\mathcal{I} = \{i\}$ for any $i \in [N]$. Then the bound (F.5a) guarantees us that, with probability at least $1 - O((N + T)^{-9})$,

$$\mathbf{U} \Sigma \text{sgn}(\mathbf{U}^\top \mathbf{U}_{\text{left}}^*) - \mathbf{U}_{\text{left}}^* \Sigma_{\text{left}}^* = \mathbf{E}_{\text{left}} \mathbf{V}_{\text{left}}^* + \Delta_{\text{left}}$$

where for each $i \in [N]$

$$\begin{aligned} \|(\Delta_{\text{left}})_{i,\cdot}\|_2 &\leq c_2 \frac{\sigma^2 (N + T_1)}{\sigma_r (\mathbf{M}_{\text{left}}^*)} \|(\mathbf{U}_{\text{left}}^*)_{i,\cdot}\|_2 + c_2 \sigma \sqrt{r + \xi_{N,T}} \left[\frac{\sigma \sqrt{T_1}}{\sigma_r (\mathbf{M}_{\text{left}}^*)} + \frac{\sigma^2 N}{\sigma_r^2 (\mathbf{M}_{\text{left}}^*)} \right] \\ &\leq c_2 c_\ell^{-1} \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^* \sqrt{T_1 / T}} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{\sigma_r^* T_1 / T} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* \sqrt{T_1 / T}} \|\mathbf{U}_{i,\cdot}^*\|_2 \right]. \end{aligned} \quad (\text{C.24})$$

The matrix $\mathbf{U}_{\text{left}}^* \Sigma_{\text{left}}^*$ can be expressed as

$$\mathbf{U}_{\text{left}}^* \Sigma_{\text{left}}^* = \mathbf{U}^* \mathbf{O}_{\text{left}}^\top \mathbf{O}_{\text{left}} \Sigma_{\text{left}}^* \mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^* = \mathbf{U}^* \Sigma_{\text{left}}^* \mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*,$$

where \mathbf{O}_{left} was defined in Appendix C.5. If we let $\mathbf{R} = \text{sgn}(\mathbf{U}^\top \mathbf{U}_{\text{left}}^*) (\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1}$, then

$$\begin{aligned} \mathbf{U} \Sigma \mathbf{R} - \mathbf{U}^* \Sigma^* &= \mathbf{E}_{\text{left}} \mathbf{V}_{\text{left}}^* (\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1} + \Delta_{\text{left}} (\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1} \\ &= \mathbf{E}_{\text{left}} \mathbf{V}_1^* \mathbf{R}_{\text{left}}^{-1} (\mathbf{V}_1^{*\top} \mathbf{V}_1^* \mathbf{R}_{\text{left}}^{-1})^{-1} + \Delta_{\text{left}} (\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1} \\ &= \mathbf{E}_{\text{left}} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} + \Delta_{\text{left}} (\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1}. \end{aligned} \quad (\text{C.25})$$

Let $\Delta = \Delta_{\text{left}} (\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1}$, and we have

$$\begin{aligned} \|\Delta_{i,\cdot}\|_2 &\leq \|(\Delta_{\text{left}})_{i,\cdot}\|_2 \|(\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1}\| \leq \|(\Delta_{\text{left}})_{i,\cdot}\|_2 \sigma_r^{-1} (\mathbf{V}_1^*) \\ &\stackrel{(i)}{\leq} c_2 c_\ell^{-3/2} \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^* \sqrt{T_1 / T}} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{\sigma_r^* T_1 / T} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* \sqrt{T_1 / T}} \|\mathbf{U}_{i,\cdot}^*\|_2 \right] \sqrt{\frac{T}{T_1}} \\ &\leq c_2 c_\ell^{-3/2} \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^* T_1 / T} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{\sigma_r^* (T_1 / T)^{3/2}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1 / T} \|\mathbf{U}_{i,\cdot}^*\|_2 \right], \end{aligned} \quad (\text{C.26})$$

where step (i) follows from equation (C.24) and the sub-block condition (4.1). In addition, with probability at least $1 - O((N + T)^{-10})$, we have

$$\|[\mathbf{E}_{\text{left}} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}]_{i,\cdot}\|_2 \stackrel{\text{(ii)}}{\leq} C_g \sigma \sqrt{r + \xi_{N,T}} \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}\| \stackrel{\text{(iii)}}{\leq} c_\ell^{-1/2} C_g \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sqrt{T_1/T}}, \quad (\text{C.27})$$

where step (ii) utilizes Lemma 19; and step (iii) follows from the sub-block condition (4.1). Putting together the relations (C.26) and (C.27) yields

$$\begin{aligned} \|\mathbf{U}_{i,\cdot} \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_{i,\cdot}^* \boldsymbol{\Sigma}^*\|_2 &\leq \|[\mathbf{E}_{\text{left}} \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}]_{i,\cdot}\|_2 + \|\boldsymbol{\Delta}_{i,\cdot}\|_2 \\ &\leq c_\ell^{-1/2} C_g \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sqrt{T_1/T}} + c_2 c_\ell^{-3/2} \left[\frac{\sigma^2 \sqrt{T_1} (r + \xi_{N,T})}{\sigma_r^* T_1/T} + \frac{\sigma^3 N \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} (T_1/T)^{3/2}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1/T} \|\mathbf{U}_{i,\cdot}^*\|_2 \right] \\ &\leq C_7 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sqrt{T_1/T}} + C_7 \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1/T} \|\mathbf{U}_{i,\cdot}^*\|_2, \end{aligned}$$

as long as $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ and $\sigma \sqrt{T} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small constant $c_{\text{noise}} > 0$, and $C_7 \geq 2c_\ell^{-1/2} C_g + 2c_2 c_\ell^{-3/2}$. In addition, we can apply the bound (F.5a) with $\mathcal{I} = [N_1]$ to obtain

$$\begin{aligned} \|(\boldsymbol{\Delta}_{\text{left}})_{\mathcal{I},\cdot}\| &\leq c_2 \sigma \sqrt{N_1} \left[\frac{\sigma \sqrt{N_1 + T_1}}{\sigma_r(\mathbf{M}_{\text{left}}^*)} + \frac{\sigma^2 N}{\sigma_r^2(\mathbf{M}_{\text{left}}^*)} \right] + c_2 \frac{\sigma^2 (N + T_1)}{\sigma_r(\mathbf{M}_{\text{left}}^*)} \|(\mathbf{U}_{\text{left}}^*)_{\mathcal{I},\cdot}\|_2 \\ &\leq c_2 c_\ell^{-1} \left[\sigma \sqrt{N_1} \left(\frac{\sigma \sqrt{N_1 + T_1}}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2 N}{\sigma_r^{*2} T_1/T} \right) + \frac{\sigma^2 (N + T_1)}{\sigma_r^* \sqrt{T_1/T}} \|\mathbf{U}_1^*\| \right]. \end{aligned} \quad (\text{C.28})$$

Therefore, we have

$$\begin{aligned} \|\boldsymbol{\Delta}_1\| &\leq \|(\boldsymbol{\Delta}_{\text{left}})_{\mathcal{I},\cdot}\| \|(\mathbf{V}_1^{*\top} \mathbf{V}_{\text{left}}^*)^{-1}\| \\ &\stackrel{\text{(i)}}{\leq} c_2 c_\ell^{-3/2} \left[\sigma \sqrt{N_1} \left(\frac{\sigma \sqrt{N_1 + T_1}}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2 N}{\sigma_r^{*2} T_1/T} \right) \sqrt{\frac{T}{T_1}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* \sqrt{T_1/T}} \|\mathbf{U}_1^*\| \right] \sqrt{\frac{T}{T_1}} \\ &\stackrel{\text{(ii)}}{\leq} 2c_2 c_\ell^{-3/2} \left[\frac{\sigma^2 \sqrt{N_1} (N_1 + T_1)}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1/T} \sqrt{\frac{N_1}{N}} \right] \stackrel{\text{(iii)}}{\leq} C_7 \frac{\sigma^2 (N + T_1)}{\sigma_r^* T_1/T} \sqrt{\frac{N_1}{N}}. \end{aligned} \quad (\text{C.29})$$

Here step (i) follows from the relations (C.28) and (4.1); step (ii) follows from the sub-block condition (4.1) and holds provided that $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small $c_{\text{noise}} > 0$, and (iii) follows from the AM-GM inequality and holds by taking $C_7 \geq 4c_2 c_\ell^{-3/2}$. We also invoke Lemma 19 to guarantee that with probability at least $1 - O((N + T)^{-10})$,

$$\|\mathbf{E}_a \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}\| \leq \frac{1}{2} C_g \sigma \sqrt{N_1 + r + \xi_{N,T}} \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}\| \leq c_\ell^{-1} C_g \frac{\sigma \sqrt{N_1}}{\sqrt{T_1/T}}. \quad (\text{C.30})$$

Collecting together claims (C.25), (C.29) and (C.30), we conclude that

$$\|\mathbf{U}_1 \boldsymbol{\Sigma} \mathbf{R} - \mathbf{U}_1^* \boldsymbol{\Sigma}^*\| \leq \|\mathbf{E}_a \mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1}\| + \|\boldsymbol{\Delta}_1\| \leq C_7 \frac{\sigma \sqrt{N_1}}{\sqrt{T_1/T}} + C_7 \frac{\sigma^2 T}{\sigma_r^*} \sqrt{\frac{N_1}{N}}$$

provided that $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small constant $c_{\text{noise}} > 0$ (Here we take $C_7 \geq 2c_\ell^{-1} C_g + 4c_2 c_\ell^{-3/2}$.)

C.7 Proof of Lemma 12

We apply Corollary 1 to the observation $\mathbf{M}_{\text{upper}} = \mathbf{M}_{\text{upper}}^* + \mathbf{E}_{\text{upper}}$, where we assume that the SVD of $\mathbf{M}_{\text{upper}}^*$ is given by $\mathbf{U}_{\text{upper}}^* \boldsymbol{\Sigma}_{\text{upper}}^* \mathbf{V}_{\text{upper}}^{*\top}$. In order to ensure that the condition of Corollary 1 is satisfied, we check that

$$\sigma \sqrt{\max\{N_1, T\}} \leq c_{\text{noise}} \sqrt{N_1/N} \stackrel{(i)}{\leq} c_n \sqrt{c_\ell N_1/N} \stackrel{(ii)}{\leq} c_n \sigma_r(\mathbf{M}_{\text{upper}}^*).$$

Here step (i) holds as long as $c_{\text{noise}} \leq c_n \sqrt{c_\ell}$, while step (ii) follows from Lemma 9. We take $\mathcal{I} = [N_1]$ and $\mathcal{J} = \{t\}$ for any $t \in [T]$. Then the relation (F.7) guarantees that with probability at least $1 - O((N+T)^{-9})$,

$$\mathbf{U}_{\text{upper}} \boldsymbol{\Sigma}_{\text{upper}} \mathbf{V}_{\text{upper}}^\top - \mathbf{M}_{\text{upper}}^* = \mathbf{E}_{\text{upper}} \mathbf{V}_{\text{upper}}^* \mathbf{V}_{\text{upper}}^{*\top} + \mathbf{U}_{\text{upper}}^* \mathbf{U}_{\text{upper}}^{*\top} \mathbf{E}_{\text{upper}} + \boldsymbol{\Phi}_{\text{upper}}$$

where for each $t \in [T]$

$$\begin{aligned} \|(\boldsymbol{\Phi}_{\text{upper}})_{\cdot,t}\|_2 &\stackrel{(i)}{\leq} 4c_3 \left[\frac{\sigma^2 (N_1 + T)}{\sigma_r(\mathbf{M}_{\text{upper}}^*)} + \sigma \sqrt{r + \xi_{N,T}} \right] \|(\mathbf{V}_{\text{upper}}^*)_{t,\cdot}\|_2 + 4c_3 \sigma \sqrt{r + \xi_{N,T}} \left[\frac{\sigma \sqrt{N_1}}{\sigma_r(\mathbf{M}_{\text{upper}}^*)} + \frac{\sigma^2 T}{\sigma_r^2(\mathbf{M}_{\text{upper}}^*)} \right] \\ &\stackrel{(ii)}{\leq} 4c_3 c_\ell^{-1} \left[\frac{\sigma^2 (N_1 + T)}{\sigma_r^* \sqrt{N_1/N}} + \sigma \sqrt{r + \xi_{N,T}} \right] \|\mathbf{V}_{t,\cdot}^*\|_2 + 4c_3 c_\ell^{-1} \sigma \sqrt{r + \xi_{N,T}} \left[\frac{\sigma \sqrt{N}}{\sigma_r^*} + \frac{\sigma^2 T}{\sigma_r^{*2} N_1/N} \right] \\ &\leq C_8 \left[\frac{\sigma^2 \sqrt{N} (r + \xi_{N,T})}{\sigma_r^*} + \frac{\sigma^3 T \sqrt{r + \xi_{N,T}}}{\sigma_r^{*2} N_1/N} + \left(\frac{\sigma^2 (N_1 + T)}{\sigma_r^* \sqrt{N_1/N}} + \sigma \sqrt{r + \xi_{N,T}} \right) \|\mathbf{V}_{t,\cdot}^*\|_2 \right]. \end{aligned}$$

Here step (i) holds provided that c_{noise} is sufficiently small; step (ii) follows from Lemma 9; and step (iii) holds by taking $C_8 \geq 4c_3 c_\ell^{-1}$. Note that $\mathbf{U}_{\text{upper}}^* \mathbf{U}_{\text{upper}}^{*\top} = \mathbf{U}_1^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}$ and $\mathbf{V}_{\text{upper}}^* \mathbf{V}_{\text{upper}}^{*\top} = \mathbf{V}^* \mathbf{V}^{*\top}$, therefore

$$\mathbf{U}_{\text{upper}} \boldsymbol{\Sigma}_{\text{upper}} \mathbf{V}_{\text{upper}}^\top - \mathbf{M}_{\text{upper}}^* = \mathbf{E}_{\text{upper}} \mathbf{V}^* \mathbf{V}^{*\top} + \mathbf{U}_1^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_{\text{upper}} + \boldsymbol{\Phi}_{\text{upper}}.$$

Finally, since we are only interested the error on the subblock \mathbf{M}_b^* , we can let

$$\mathbf{W} := \mathbf{E}_a \mathbf{V}_1^* \mathbf{V}_2^{*\top} + \mathbf{E}_b \mathbf{V}_2^* \mathbf{V}_2^{*\top} + \mathbf{U}_1^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top} \mathbf{E}_b \quad \text{and} \quad \boldsymbol{\Phi} := (\boldsymbol{\Phi}_{\text{upper}})_{\cdot,(T_1+1):T}$$

to obtain the desired bound. In addition, we can invoke equation (F.8) to show that with probability at least $1 - O((N+T)^{-9})$, for each $t \in [T]$

$$\begin{aligned} \|(\mathbf{M}_b - \mathbf{M}_b^*)_{\cdot,t}\|_2 &\leq 2c_3 \left[\sigma \sqrt{r + \xi_{N,T}} + \sigma \sqrt{N_1} \|\mathbf{V}_{t,\cdot}^*\|_2 + \frac{\sigma^2 (N_1 + T)}{\sigma_r(\mathbf{M}_{\text{upper}}^*)} \|\mathbf{V}_{t,\cdot}^*\|_2 + \frac{\sigma^2 \sqrt{N_1} (r + \xi_{N,T})}{\sigma_r(\mathbf{M}_{\text{upper}}^*)} \right] \\ &\stackrel{(a)}{\geq} C_8 \sigma \sqrt{r + \xi_{N,T}} + C_8 \left[\frac{\sigma^2 T}{\sigma_r^* \sqrt{N_1/N}} + \sigma \sqrt{N_1} \right] \|\mathbf{V}_{t,\cdot}^*\|_2. \end{aligned}$$

Here step (a) follows from Lemma 9, and holds as long as $C_8 \geq 4c_3 c_\ell^{-1/2}$.

C.8 Proof of Lemma 13

We apply Corollary 2 with $\mathcal{I} = [N_1]$ to show that with probability at least $1 - O((N + T)^{-10})$

$$\begin{aligned}
\|U_1^{\star\top}(\mathbf{U}_1\mathbf{H} - U_1^{\star})\| &\leq c_4 \left[\frac{\sigma\sqrt{r + \xi_{N,T}}}{\sigma_r(\mathbf{M}_{\text{left}}^{\star})} + \frac{\sigma^2\sqrt{(N_1 + T_1)(T_1)}}{\sigma_r^2(\mathbf{M}_{\text{left}}^{\star})} + \frac{\sigma^3 N\sqrt{N_1}}{\sigma_r^3(\mathbf{M}_{\text{left}}^{\star})} \right] \|U_1^{\star}\| + c_4 \frac{\sigma^2(N + T_1)}{\sigma_r^2(\mathbf{M}_{\text{left}}^{\star})} \|U_1^{\star}\|^2 \\
&\stackrel{(i)}{\leq} c_4 c_\ell^{-1} \left[\frac{\sigma\sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2\sqrt{(N_1 + T_1)T_1}}{\sigma_r^{*2} T_1/T} + \frac{\sigma^3 N\sqrt{N_1}}{(\sigma_r^* \sqrt{T_1/T})^3} \right] \sqrt{\frac{N_1}{N}} + c_4 c_\ell^{-1} \frac{\sigma^2(N + T_1) N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N} \\
&\stackrel{(ii)}{\leq} 4c_4 c_\ell^{-1} \frac{\sigma\sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \sqrt{\frac{N_1}{N}} + 4c_4 c_\ell^{-1} \frac{\sigma^2(N + T_1) N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N}, \tag{C.31}
\end{aligned}$$

Here step (i) follows from Lemma 9, whereas step (ii) holds provided that $\sigma\sqrt{NT/T_1} \leq c_{\text{noise}}\sigma_r^*$ and $\sigma\sqrt{T} \leq c_{\text{noise}}\sigma_r^*$ for some sufficiently small constant $c_{\text{noise}} > 0$. It is worth mentioning that, we also use the AM-GM inequality in the relation (C.31), which guarantees that

$$\frac{\sigma^2\sqrt{N_1 T_1}}{\sigma_r^{*2} T_1/T} \sqrt{\frac{N_1}{N}} = \frac{\sigma^2\sqrt{NT_1} N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N} \leq \frac{1}{2} \frac{\sigma^2(N + T_1) N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N}.$$

The inequality (C.31) allows us to control some other quantities. For example, we can decompose

$$(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H} - U_1^{\star\top} U_1^{\star} = (\mathbf{U}_1\mathbf{H} - U_1^{\star})^\top U_1^{\star} + U_1^{\star\top} (\mathbf{U}_1\mathbf{H} - U_1^{\star}) + (\mathbf{U}_1\mathbf{H} - U_1^{\star})^\top (\mathbf{U}_1\mathbf{H} - U_1^{\star}),$$

which leads to the bound

$$\begin{aligned}
\|(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H} - U_1^{\star\top} U_1^{\star}\| &\leq 2\|U_1^{\star\top}(\mathbf{U}_1\mathbf{H} - U_1^{\star})\| + \|\mathbf{U}_1\mathbf{H} - U_1^{\star}\|^2 \\
&\stackrel{(a)}{\leq} 8c_4 c_\ell^{-1} \frac{\sigma\sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \sqrt{\frac{N_1}{N}} + 8c_4 c_\ell^{-1} \frac{\sigma^2(N + T_1) N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N} + C_6^2 \left[\frac{\sigma^2 T}{\sigma_r^{*2}} \sqrt{\frac{N_1}{N}} + \frac{\sigma\sqrt{N_1}}{\sigma_r^* \sqrt{T_1/T}} \right]^2 \\
&\stackrel{(b)}{\leq} (16c_4 c_\ell^{-1} + 2C_6^2) \left[\frac{\sigma\sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \sqrt{\frac{N_1}{N}} + \frac{\sigma^2(N + T_1) N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N} \right]. \tag{C.32}
\end{aligned}$$

Here step (a) follows from the relations (C.31) and (C.5), while step (b) holds provided that c_{noise} is sufficiently small. In addition, in view of the decomposition

$$[(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} - (U_1^{\star\top} U_1^{\star})^{-1} = (U_1^{\top} U_1)^{-1} [(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H} - U_1^{\star\top} U_1^{\star}] (U_1^{\star\top} U_1^{\star})^{-1},$$

we can show that

$$\begin{aligned}
\|[(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} - (U_1^{\star\top} U_1^{\star})^{-1}\| &\leq \sigma_r^{-2}(\mathbf{U}_1) \sigma_r^{-2}(U_1^{\star}) \left\| (\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H} - U_1^{\star\top} U_1^{\star} \right\| \\
&\leq 2c_\ell^{-2} (16c_4 c_\ell^{-1} + 2C_6^2) \left(\frac{N}{N_1} \right)^2 \left[\frac{\sigma\sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \sqrt{\frac{N_1}{N}} + \frac{\sigma^2(N + T_1) N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N} \right]. \tag{C.33}
\end{aligned}$$

Here the last step follows from equations (C.6), (C.6) and (C.32). Starting from

$$\begin{aligned}
&[(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} (\mathbf{U}_1\mathbf{H})^\top - (U_1^{\star\top} U_1^{\star})^{-1} U_1^{\star\top} \\
&= \{[(\mathbf{U}_1\mathbf{H})^\top \mathbf{U}_1\mathbf{H}]^{-1} - (U_1^{\star\top} U_1^{\star})^{-1}\} (\mathbf{U}_1\mathbf{H})^\top + (U_1^{\star\top} U_1^{\star})^{-1} (\mathbf{U}_1\mathbf{H} - U_1^{\star})^\top,
\end{aligned}$$

we can also deduce that

$$\begin{aligned}
& \left\| [(\mathbf{U}_1 \mathbf{H})^\top \mathbf{U}_1 \mathbf{H}]^{-1} (\mathbf{U}_1 \mathbf{H})^\top - (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_1^{\star\top} \right\| \\
& \leq \left\| [(\mathbf{U}_1 \mathbf{H})^\top \mathbf{U}_1 \mathbf{H}]^{-1} - (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \right\| \|\mathbf{U}_1\| + \sigma_r^{-2} (\mathbf{U}_1^\star) \|\mathbf{U}_1 \mathbf{H} - \mathbf{U}_1^\star\| \\
& \stackrel{(i)}{\leq} 2c_\ell^{-2} (16c_4 c_\ell^{-1} + 2C_6^2) \left(\frac{N}{N_1} \right)^2 \left[\frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \sqrt{\frac{N_1}{N}} + \frac{\sigma^2 (N + T_1) N_1}{\sigma_r^{*2} T_1/T} \frac{N_1}{N} \right] \sqrt{c_u \frac{N_1}{N}} \\
& \quad + c_\ell^{-1} C_6 \frac{N}{N_1} \left[\frac{\sigma^2 T}{\sigma_r^{*2}} \sqrt{\frac{N_1}{N}} + \frac{\sigma \sqrt{N_1}}{\sigma_r^* \sqrt{T_1/T}} \right] \\
& \stackrel{(ii)}{\leq} 2c_\ell^{-1} C_6 \frac{\sigma^2 T}{\sigma_r^{*2}} \sqrt{\frac{N}{N_1}} + 2c_\ell^{-1} C_6 \frac{\sigma \sqrt{N_1}}{\sigma_r^* \sqrt{T_1/T}} \frac{N}{N_1}.
\end{aligned}$$

Here step (i) follows from the bounds (C.6), (C.33) and (C.5), while step (ii) holds provided that $\sigma \sqrt{NT/T_1} \leq c_{\text{noise}} \sigma_r^*$ for some sufficiently small constant $c_{\text{noise}} > 0$. Taking $C_9 = 2c_\ell^{-2} (16c_4 c_\ell^{-1} + 2C_6^2)$ completes the proof.

D Proof of Lemmas from Section 5.2

D.1 Proof of Lemma 6

Similar to the proof of Lemma 12, we apply Corollary 1 to the observation $\mathbf{M}_{\text{upper}} = \mathbf{M}_{\text{upper}}^* + \mathbf{E}_{\text{upper}}$. The condition of Corollary 1 was checked in the proof of Lemma 12. Then with probability exceeding $1 - O((N + T)^{-10})$, we have

$$\begin{aligned}
\left\| \mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}} \right\| & \stackrel{(i)}{\leq} 2c_3 \sigma \sqrt{T} + 2c_3 \sigma \sqrt{N_1} + c_3 \frac{\sigma^2 (N_1 + T)}{\sigma_r^* (\mathbf{M}_{\text{upper}}^*)} + c_3 \frac{\sigma^2}{\sigma_r^*} \sqrt{N_1 T} \\
& \stackrel{(ii)}{\leq} 4c_3 \sigma \sqrt{N_1 + T} + c_3 \frac{\sigma^2 (N_1 + T)}{\sigma_r^* \sqrt{N_1/N}} + c_3 \frac{\sigma^2}{\sigma_r^*} \sqrt{N_1 T} \stackrel{(iii)}{\leq} 5c_3 \sigma \sqrt{N_1 + T}.
\end{aligned}$$

Here step (i) is obtained by taking $\mathcal{I} = [N_1]$ and $\mathcal{J} = [T]$ in (F.8) in Corollary 1; step (ii) uses the sub-block condition (4.1); step (iii) follows from the noise condition (4.3). Finally, we have

$$\left\| \mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}} \right\|_{\text{F}} \leq \sqrt{r} \left\| \mathbf{M}_{\text{upper}}^* - \widehat{\mathbf{M}}_{\text{upper}} \right\| \leq 5c_3 \sigma \sqrt{(N_1 + T)r}.$$

The desired bound can be obtained by taking $C_{\text{upper}} = 5c_3$.

D.2 Proof of Lemma 5

We first decompose $\Delta_U := \sigma^2 |U_{i,\cdot} (\mathbf{U}_1^\top \mathbf{U}_1)^{-1} \mathbf{U}_{i,\cdot}^\top - U_{i,\cdot}^* (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_{i,\cdot}^{\star\top}|$ into three terms

$$\begin{aligned}
\Delta_U & = \sigma^2 |U_{i,\cdot} \mathbf{H} [(\mathbf{U}_1 \mathbf{H})^\top \mathbf{U}_1 \mathbf{H}]^{-1} (\mathbf{U}_{i,\cdot} \mathbf{H})^\top - U_{i,\cdot}^* (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_{i,\cdot}^{\star\top}| \\
& \leq \underbrace{2\sigma^2 |(U_{i,\cdot} \mathbf{H} - U_{i,\cdot}^*) (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \mathbf{U}_{i,\cdot}^{\star\top}|}_{=:\alpha_1} + \underbrace{\sigma^2 |U_{i,\cdot} \mathbf{H} \{ [(\mathbf{U}_1 \mathbf{H})^\top \mathbf{U}_1 \mathbf{H}]^{-1} - (\mathbf{U}_1^{\star\top} \mathbf{U}_1^\star)^{-1} \} \mathbf{U}_{i,\cdot}^{\star\top}|}_{=:\alpha_2} \\
& \quad + \underbrace{\sigma^2 |(U_{i,\cdot} \mathbf{H} - U_{i,\cdot}^*) [(\mathbf{U}_1 \mathbf{H})^\top \mathbf{U}_1 \mathbf{H}]^{-1} (\mathbf{U}_{i,\cdot} \mathbf{H} - U_{i,\cdot}^*)^\top|}_{=:\alpha_3}.
\end{aligned}$$

On the other hand, we can lower bound $\gamma_{i,t}^*$ by

$$\gamma_{i,t}^* \stackrel{(i)}{\geq} \frac{1}{c_\ell} \sigma^2 \frac{N}{N_1} \|\mathbf{U}_{i,\cdot}^*\|_2^2 + \frac{1}{c_\ell} \sigma^2 \frac{T}{T_1} \|\mathbf{V}_{t,\cdot}^*\|_2^2 = \frac{1}{c_\ell} \sigma^2 \frac{\mu_i^2 r}{N_1} + \frac{1}{c_\ell} \sigma^2 \frac{\nu_t^2 r}{T_1} \stackrel{(ii)}{\geq} \frac{1}{c_{\text{inc}}^2 c_\ell \delta^2} \frac{\sigma^4}{\sigma_r^{*2}} \frac{NT}{N_1 T_1} \xi_{N,T}^2. \quad (\text{D.1a})$$

Here step (i) follows from we the sub-block condition (4.1), step (ii) follows from the incoherence condition (4.4c). By the AM-GM inequality, the above bounds further give

$$\gamma_{i,t}^* \geq \frac{\xi_{N,T}}{2\sqrt{2}c_\ell c_{\text{inc}} \delta} \frac{\sigma^3}{\sigma_r^*} \frac{N}{N_1} \sqrt{\frac{T}{T_1}} \|\mathbf{U}_{i,\cdot}^*\|_2 + \frac{\xi_{N,T}}{2\sqrt{2}c_\ell c_{\text{inc}} \delta} \frac{\sigma^3}{\sigma_r^*} \frac{T}{T_1} \sqrt{\frac{N}{N_1}} \|\mathbf{V}_{t,\cdot}^*\|_2. \quad (\text{D.1b})$$

Then we bound α_1 , α_2 and α_3 separately. The first term α_1 can be bounded by

$$\begin{aligned} \alpha_1 &\stackrel{(a)}{\leq} \frac{1}{c_\ell} \sigma^2 \frac{N}{N_1} \|\mathbf{U}_{i,\cdot} \mathbf{H} - \mathbf{U}_{i,\cdot}^*\|_2 \|\mathbf{U}_{i,\cdot}^*\|_2 \stackrel{(b)}{\leq} \frac{C_6}{c_\ell} \sigma^2 \frac{N}{N_1} \left(\frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^{*2} T_1/T} \|\mathbf{U}_{i,\cdot}^*\|_2 \right) \|\mathbf{U}_{i,\cdot}^*\|_2 \\ &\stackrel{(c)}{\leq} \frac{\delta}{12\sqrt{\xi_{N,T}}} \left(\frac{\xi_{N,T}}{2\sqrt{2}c_\ell c_{\text{inc}} \delta} \frac{\sigma^3}{\sigma_r^*} \frac{N}{N_1} \sqrt{\frac{T}{T_1}} \|\mathbf{U}_{i,\cdot}^*\|_2 + \frac{1}{c_\ell} \sigma^2 \frac{N}{N_1} \|\mathbf{U}_{i,\cdot}^*\|_2^2 \right) \stackrel{(d)}{\leq} \frac{\delta}{6\sqrt{\xi_{N,T}}} \gamma_{i,t}^* \end{aligned}$$

Here step (a) follows from the sub-block condition (4.1), step (b) follows from the bound (C.4) in Lemma 10, step (c) follows from the noise condition (4.3) and holds provided that c_{inc} is sufficiently small, and step (d) uses (D.1a) and (D.1b). For the second term α_2 , we have

$$\begin{aligned} \alpha_2 &\leq \sigma^2 \left[(\mathbf{U}_1 \mathbf{H})^\top \mathbf{U}_1 \mathbf{H} \right]^{-1} - (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \|\mathbf{U}_{i,\cdot}\|_2 \|\mathbf{U}_{i,\cdot}^*\|_2 \\ &\stackrel{(i)}{\leq} C_9 \left(\frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \sqrt{\frac{N_1}{N}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^{*2} T_1/T} \frac{N_1}{N} \right) \frac{N^2}{N_1^2} \sigma^2 \|\mathbf{U}_{i,\cdot}^*\|_2 \left(2\|\mathbf{U}_{i,\cdot}^*\|_2 + C_6 \frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} \right) \\ &\stackrel{(ii)}{\leq} \frac{\delta}{12\sqrt{\xi_{N,T}}} \left(\frac{\xi_{N,T}}{2\sqrt{2}c_\ell c_{\text{inc}} \delta} \frac{\sigma^3}{\sigma_r^*} \frac{N}{N_1} \sqrt{\frac{T}{T_1}} \|\mathbf{U}_{i,\cdot}^*\|_2 + \frac{1}{c_\ell} \sigma^2 \frac{N}{N_1} \|\mathbf{U}_{i,\cdot}^*\|_2^2 \right) \stackrel{(iii)}{\leq} \frac{\delta}{6\sqrt{\xi_{N,T}}} \gamma_{i,t}^* \end{aligned}$$

Here step (i) follows from (C.12) in Lemma 13 and (C.17), step (ii) holds under the noise condition (4.3), and step (iii) follows from uses (D.1a) and (D.1b). The last term α_3 is bounded by

$$\begin{aligned} \alpha_3 &\stackrel{(a)}{\leq} \frac{1}{c_\ell} \sigma^2 \frac{N}{N_1} \|\mathbf{U}_{i,\cdot} \mathbf{H} - \mathbf{U}_{i,\cdot}^*\|_2^2 \stackrel{(b)}{\leq} \frac{C_6^2}{c_\ell} \sigma^2 \frac{N}{N_1} \left(\frac{\sigma \sqrt{r + \xi_{N,T}}}{\sigma_r^* \sqrt{T_1/T}} + \frac{\sigma^2 (N + T_1)}{\sigma_r^{*2} T_1/T} \|\mathbf{U}_{i,\cdot}^*\|_2 \right)^2 \\ &\stackrel{(c)}{\leq} \frac{\delta}{12\sqrt{\xi_{N,T}}} \left(\frac{1}{c_{\text{inc}}^2 c_\ell \delta^2} \frac{\sigma^4}{\sigma_r^{*2}} \frac{NT}{N_1 T_1} \xi_{N,T}^2 + \frac{1}{c_\ell} \sigma^2 \frac{N}{N_1} \|\mathbf{U}_{i,\cdot}^*\|_2^2 \right) \stackrel{(d)}{\leq} \frac{\delta}{6\sqrt{\xi_{N,T}}} \gamma_{i,t}^* \end{aligned}$$

Here step (a) follows from the sub-block condition (4.1), step (b) follows from the bound (C.4) in Lemma 10, step (c) follows from the noise condition (4.3) and holds provided that c_{inc} is sufficiently small, and step (d) uses (D.1a). Taking above bounds on α_1 , α_2 and α_3 collectively yields

$$\Delta_U \leq \alpha_1 + \alpha_2 + \alpha_3 \leq \frac{\delta}{2\sqrt{\xi_{N,T}}} \gamma_{i,t}^*.$$

A similar argument can be used to show that $\Delta_V := \sigma^2 |\mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^\top - \mathbf{V}_{t,\cdot}^* (\mathbf{V}_1^{\top} \mathbf{V}_1)^{-1} \mathbf{V}_{t,\cdot}^\top|$ admits the same upper bound. Putting the bounds for Δ_U and Δ_V together gives the claimed bound.

E Proof of Theorem 3

So as to simplify notation, we introduce the shorthand $\bar{\mathbf{M}}^* := \mathbf{M}_{1:\bar{N},1:\bar{T}}^*$. Note that $\bar{\mathbf{M}}^*$ can be viewed as the ground truth of the four-block data matrix $\mathbf{M}^{(i_0, j_0)}$. The following lemma characterizes its spectrum.

Lemma 14. *Under the sub-block condition (4.11), we have*

$$c_\ell \sqrt{\frac{\bar{N}\bar{T}}{NT}} \sigma_r^* \leq \sigma_r(\bar{\mathbf{M}}^*) \leq \sigma_r(\mathbf{M}^*) \leq c_u \sqrt{\frac{\bar{N}\bar{T}}{NT}} \sigma_r^*.$$

See [Appendix E.1](#) for the proof.

Beginning with the singular value decomposition $\bar{\mathbf{M}}^* = \bar{\mathbf{U}}^* \bar{\mathbf{\Sigma}}^* \bar{\mathbf{V}}^{*\top}$, we can partition the matrices $\bar{\mathbf{U}}^*$ and $\bar{\mathbf{V}}^*$ into two blocks

$$\bar{\mathbf{U}}^* = \begin{bmatrix} \bar{\mathbf{U}}_1^* \\ \bar{\mathbf{U}}_2^* \end{bmatrix} \quad \text{and} \quad \bar{\mathbf{V}}^* = \begin{bmatrix} \bar{\mathbf{V}}_1^* \\ \bar{\mathbf{V}}_2^* \end{bmatrix},$$

where $\bar{\mathbf{U}}_1^* \in \mathbb{R}^{\bar{N}_1 \times r}$, $\bar{\mathbf{U}}_2^* \in \mathbb{R}^{\bar{N}_2 \times r}$, $\bar{\mathbf{V}}_1^* \in \mathbb{R}^{\bar{T}_1 \times r}$ and $\bar{\mathbf{V}}_2^* \in \mathbb{R}^{\bar{T}_2 \times r}$. Note that $\bar{\mathbf{M}}^*$ can also be written as $\mathbf{U}_{1:\bar{N}}^* \mathbf{\Sigma}^* (\mathbf{V}_{1:\bar{T}}^*)^\top$. We know that there exists two invertible matrices $\mathbf{R}_U, \mathbf{R}_V \in \mathbb{R}^{r \times r}$ such that $\mathbf{U}_{1:\bar{N}}^* = \bar{\mathbf{U}}^* \mathbf{R}_U$ and $\mathbf{V}_{1:\bar{T}}^* = \bar{\mathbf{V}}^* \mathbf{R}_V$. Since $\bar{\mathbf{U}}^*$ has orthonormal columns, we have $\mathbf{R}_U = \bar{\mathbf{U}}^{*\top} \mathbf{U}_{1:\bar{N}}^*$, and therefore

$$\mathbf{R}_U \mathbf{R}_U^\top = \bar{\mathbf{U}}^{*\top} \mathbf{U}_{1:\bar{N}}^* \mathbf{U}_{1:\bar{N}}^{*\top} \bar{\mathbf{U}}^* = \bar{\mathbf{U}}^{*\top} (\mathbf{U}_1^{*\top} \mathbf{U}_1^* + \mathbf{U}_2^{*\top} \mathbf{U}_2^*) \bar{\mathbf{U}}^*.$$

In view of the sub-block condition (4.11), we know that

$$c_\ell \frac{\bar{N}}{N} \mathbf{I}_r = c_\ell \frac{\bar{N}}{N} \bar{\mathbf{U}}^{*\top} \bar{\mathbf{U}}^* \preceq \mathbf{R}_U \mathbf{R}_U^\top \preceq c_u \frac{\bar{N}}{N} \bar{\mathbf{U}}^{*\top} \bar{\mathbf{U}}^* = c_u \frac{\bar{N}}{N} \mathbf{I}_r.$$

From equation (4.11) and the fact that $\mathbf{U}_1^* = \bar{\mathbf{U}}_1^* \mathbf{R}_U$, we can check that

$$\begin{aligned} \bar{\mathbf{U}}_1^{*\top} \bar{\mathbf{U}}_1^* &= (\mathbf{R}_U)^{-\top} \mathbf{U}_1^{*\top} \mathbf{U}_1^* (\mathbf{R}_U)^{-1} \succeq c_\ell \frac{\bar{N}_1}{N} (\mathbf{R}_U)^{-\top} (\mathbf{R}_U)^{-1} = c_\ell \frac{\bar{N}_1}{N} (\mathbf{R}_U \mathbf{R}_U^\top)^{-1} \succeq \frac{c_\ell}{c_u} \frac{\bar{N}_1}{N} \mathbf{I}_r, \\ \bar{\mathbf{U}}_1^{*\top} \bar{\mathbf{U}}_1^* &= (\mathbf{R}_U)^{-\top} \mathbf{U}_1^{*\top} \mathbf{U}_1^* (\mathbf{R}_U)^{-1} \preceq c_u \frac{\bar{N}_1}{N} (\mathbf{R}_U)^{-\top} (\mathbf{R}_U)^{-1} = c_u \frac{\bar{N}_1}{N} (\mathbf{R}_U \mathbf{R}_U^\top)^{-1} \preceq \frac{c_u}{c_\ell} \frac{\bar{N}_1}{N} \mathbf{I}_r \end{aligned}$$

hold. A similar argument can be used to verify that

$$c_\ell \frac{\bar{T}}{T} \mathbf{I}_r \preceq \mathbf{R}_V \mathbf{R}_V^\top \preceq c_u \frac{\bar{T}}{T} \mathbf{I}_r \quad \text{and} \quad \frac{c_\ell}{c_u} \frac{\bar{T}_1}{T} \mathbf{I}_r \preceq \bar{\mathbf{V}}_1^{*\top} \bar{\mathbf{V}}_1^* \preceq \frac{c_u}{c_\ell} \frac{\bar{T}_1}{T} \mathbf{I}_r.$$

Therefore, the sub-block condition (4.1) holds for $\bar{\mathbf{M}}^*$. We can also check that the assumptions of [Theorem 3](#), together with [Lemma 14](#) guarantees that all conditions of [Theorem 1](#) hold. Then we can apply [Theorem 1](#) to the four-block data matrix $\mathbf{M}^{(i_0, j_0)}$ to show that we can decompose

$$(\widehat{\mathbf{M}}_{i_0, j_0} - \mathbf{M}_{i_0, j_0}^*)_{i,t} = g_{i,t} + \Delta_{i,t},$$

where $g_{i,t}$ is a mean-zero Gaussian random variable with variance

$$\begin{aligned} \text{var}(g_{i,t}) &= \frac{\omega^2}{NT} \|\bar{\mathbf{U}}_{i,\cdot}^* (\bar{\mathbf{U}}_1^{*\top} \bar{\mathbf{U}}_1^*)^{-1} \bar{\mathbf{U}}_1^{*\top}\|_2^2 + \frac{\omega^2}{NT} \|\bar{\mathbf{V}}_1^* (\bar{\mathbf{V}}_1^{*\top} \bar{\mathbf{V}}_1^*)^{-1} \bar{\mathbf{V}}_{t,\cdot}^{*\top}\|_2^2 \\ &= \frac{\omega^2}{NT} \|\mathbf{U}_{i,\cdot}^* \mathbf{R}_U^{-1} (\mathbf{R}_U^{-\top} \mathbf{U}_1^{*\top} \mathbf{U}_1^* \mathbf{R}_U^{-1})^{-1} \mathbf{R}_U^{-\top} \mathbf{U}_1^{*\top}\|_2^2 + \frac{\omega^2}{NT} \|\mathbf{V}_1^* \mathbf{R}_V^{-1} (\mathbf{R}_V^{-\top} \mathbf{V}_1^{*\top} \mathbf{V}_1^* \mathbf{R}_V^{-1})^{-1} \mathbf{R}_V^{-\top} \mathbf{V}_{t,\cdot}^{*\top}\|_2^2 \\ &= \frac{\omega^2}{NT} \|\mathbf{U}_{i,\cdot}^* (\mathbf{U}_1^{*\top} \mathbf{U}_1^*)^{-1} \mathbf{U}_1^{*\top}\|_2^2 + \frac{\omega^2}{NT} \|\mathbf{V}_1^* (\mathbf{V}_1^{*\top} \mathbf{V}_1^*)^{-1} \mathbf{V}_{t,\cdot}^{*\top}\|_2^2 = \gamma_{i,t}^*, \end{aligned}$$

and $|\Delta_{i,t}| \leq \delta \gamma_{i,t}^{*1/2}$ holds with probability at least $1 - O((N+T)^{-10})$.

E.1 Proof of Lemma 14

We can use Lemma 9 to establish Lemma 14. In view of the sub-block condition (4.11), we have

$$c_\ell \frac{\bar{T}}{T} \mathbf{I}_r \preceq \mathbf{V}_{1:\bar{T},\cdot}^{*\top} \mathbf{V}_{1:\bar{T},\cdot}^* = \mathbf{V}_1^{*\top} \mathbf{V}_1^* + \mathbf{V}_2^{*\top} \mathbf{V}_2^* \preceq c_u \frac{\bar{T}}{T} \mathbf{I}_r.$$

Consequently, we can apply the bound (C.2a) to \mathbf{M}^* so as to obtain

$$\sqrt{c_\ell \frac{\bar{T}}{T}} \sigma_r^* \leq \sigma_r(\mathbf{M}_{\cdot,1:\bar{T}}^*) \leq \sqrt{c_u \frac{\bar{T}}{T}} \sigma_r^*.$$

We form the SVD $\mathbf{M}_{\cdot,1:\bar{T}}^* = \mathbf{U}_{\text{left}}^* \boldsymbol{\Sigma}_{\text{left}}^* \mathbf{V}_{\text{left}}^{*\top}$. Since the matrix $\mathbf{M}_{\cdot,1:\bar{T}}^*$ can also be written as $\mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}_{1:\bar{T},\cdot}^{*\top}$, there exists some rotation matrix \mathbf{H} such that $\mathbf{U}_{\text{left}}^* = \mathbf{U}^* \mathbf{H}$. If we write

$$\mathbf{U}_{\text{left}}^* = \begin{bmatrix} \mathbf{U}_{\text{left},1}^* \\ \mathbf{U}_{\text{left},2}^* \end{bmatrix}$$

where $\mathbf{U}_{\text{left},1}^* \in \mathbb{R}^{\bar{N} \times r}$ and $\mathbf{U}_{\text{left},2}^* \in \mathbb{R}^{(N-\bar{N}) \times r}$. Thus, we can write

$$\mathbf{U}_{\text{left},1}^{*\top} \mathbf{U}_{\text{left},1}^* = \mathbf{H}^\top \mathbf{U}_{1:\bar{N},\cdot}^{*\top} \mathbf{U}_{1:\bar{N},\cdot}^* \mathbf{H}, \quad \mathbf{H} = \mathbf{H}^\top (\mathbf{U}_1^{*\top} \mathbf{U}_1^* + \mathbf{U}_2^{*\top} \mathbf{U}_2^*) \mathbf{H}.$$

In view of the sub-block condition (4.11), we know that

$$c_\ell \frac{\bar{N}}{N} \mathbf{I}_r \preceq \mathbf{U}_{\text{left},1}^{*\top} \mathbf{U}_{\text{left},1}^* \preceq c_u \frac{\bar{N}}{N} \mathbf{I}_r.$$

Therefore, we can apply the bound (C.2b) on $\mathbf{M}_{\cdot,1:\bar{T}}^*$ to obtain

$$c_\ell \sqrt{\frac{\bar{N}\bar{T}}{NT}} \sigma_r^* \leq \sqrt{c_\ell \frac{\bar{N}}{N}} \sigma_r(\mathbf{M}_{\cdot,1:\bar{T}}^*) \leq \sigma_r(\mathbf{M}_{1:\bar{N},1:\bar{T}}^*) \leq \sqrt{c_u \frac{\bar{N}}{N}} \sigma_r(\mathbf{M}_{\cdot,1:\bar{T}}^*) \leq c_u \sqrt{\frac{\bar{N}\bar{T}}{NT}} \sigma_r^*.$$

F Auxiliary results on matrix denoising

In this appendix, we collect together various results on matrix denoising that are used in proving Lemma 1. Let $\mathbf{M}^* \in \mathbb{R}^{n_1 \times n_2}$ be a rank r matrix with SVD $\mathbf{M}^* = \mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}^{*\top}$, where $\mathbf{U}^* \in \mathbb{R}^{n_1 \times r}$ and $\mathbf{V}^* \in \mathbb{R}^{n_2 \times r}$ have orthonormal columns, and $\boldsymbol{\Sigma}^* = \text{diag}(\sigma_1^*, \dots, \sigma_r^*)$ is a diagonal matrix with

$\sigma_1^* \geq \sigma_2^* \geq \dots \geq \sigma_r^*$. For simplicity, let $n := \max\{n_1, n_2\}$. Let \mathbf{E} be a noise matrix whose entries are independent sub-Gaussian random variables with sub-Gaussian norm at most σ , i.e. $\|E_{i,j}\|_{\psi_2} \leq \sigma$ (see [Ver17, Definition 2.5.6]). Suppose that we observe

$$\mathbf{M} = \mathbf{M}^* + \mathbf{E}$$

and our goal is to estimate the unknown low-rank matrix \mathbf{M}^* and its singular subspaces \mathbf{U}^* and \mathbf{V}^* . Perhaps the most natural approach is to compute the (truncated) rank- r SVD $\mathbf{U}\Sigma\mathbf{V}^\top$ of \mathbf{M} , and use \mathbf{U} , \mathbf{V} and $\mathbf{U}\Sigma\mathbf{V}^\top$ to estimate \mathbf{U}^* , \mathbf{V}^* and \mathbf{M}^* respectively.

Our first result concerns subspace estimation error. Due to rotational ambiguity of linear subspaces, namely for any rotation matrix $\mathbf{O} \in \mathcal{O}^{r \times r}$, $\mathbf{U}\mathbf{O}$ (resp. $\mathbf{V}\mathbf{O}$) represents the same subspace as \mathbf{U} (resp. \mathbf{V}), we define the optimal rotation matrices \mathbf{R}_U that best aligns \mathbf{U} with \mathbf{U}^* in the Euclidean sense:

$$\mathbf{R}_U := \arg \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|\mathbf{U}\mathbf{R} - \mathbf{U}^*\|_F.$$

We know that \mathbf{R}_U is the matrix sign of $\mathbf{U}^\top \mathbf{U}^*$, which is given by $\mathbf{X}\mathbf{Y}^\top$ where $\mathbf{X}\mathbf{Y}\mathbf{Z}^\top$ is the SVD of $\mathbf{U}^\top \mathbf{U}^*$. Similarly we define

$$\mathbf{R}_V := \arg \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|\mathbf{V}\mathbf{R} - \mathbf{V}^*\|_F.$$

The following proposition gives first-order expansions of $\mathbf{U}\mathbf{R}_U - \mathbf{U}^*$ and $\mathbf{V}\mathbf{R}_V - \mathbf{V}^*$ with respect to the random perturbation matrix \mathbf{E} .

Proposition 2. *Suppose that $\sigma\sqrt{n} \leq c_n\sigma_r^*$ for some sufficiently small constant $c_n > 0$. Then we can decompose*

$$\mathbf{U}\mathbf{R}_U - \mathbf{U}^* = \mathbf{E}\mathbf{V}^*(\Sigma^*)^{-1} + \Psi_U, \quad \text{and} \quad (\text{F.1a})$$

$$\mathbf{V}\mathbf{R}_V - \mathbf{V}^* = \mathbf{E}^\top \mathbf{U}^*(\Sigma^*)^{-1} + \Psi_V, \quad (\text{F.1b})$$

where Ψ_U and Ψ_V are two matrices that satisfy: there exists some universal constant $c_1 > 0$ such that for any given $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$,

$$\|(\Psi_U)_{\mathcal{I},\cdot}\| \leq c_1 \frac{\sigma\sqrt{I+r+\log n}}{\sigma_r^*} \left[\frac{\sigma\sqrt{I+n_2+\log n}}{\sigma_r^*} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right] + c_1 \left(\frac{\sigma^2 n}{\sigma_r^{*2}} + \frac{\sigma\sqrt{r+\log n}}{\sigma_r^*} \right) \|\mathbf{U}_{\mathcal{I},\cdot}^*\|, \quad (\text{F.2a})$$

$$\|(\Psi_V)_{\mathcal{J},\cdot}\| \leq c_1 \frac{\sigma\sqrt{J+r+\log n}}{\sigma_r^*} \left[\frac{\sigma\sqrt{J+n_1+\log n}}{\sigma_r^*} + \frac{\sigma^2 n_2}{\sigma_r^{*2}} \right] + c_1 \left(\frac{\sigma^2 n}{\sigma_r^{*2}} + \frac{\sigma\sqrt{r+\log n}}{\sigma_r^*} \right) \|\mathbf{V}_{\mathcal{J},\cdot}^*\|, \quad (\text{F.2b})$$

where $I = |\mathcal{I}|$ and $J = |\mathcal{J}|$. In addition, with probability exceeding $1 - O(n^{-10})$,

$$\|\mathbf{U}_{\mathcal{I},\cdot}\mathbf{R}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| \leq c_1 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_1 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n}, \quad (\text{F.3a})$$

$$\|\mathbf{V}_{\mathcal{J},\cdot}\mathbf{R}_V - \mathbf{V}_{\mathcal{J},\cdot}^*\| \leq c_1 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{V}_{\mathcal{J},\cdot}^*\| + c_1 \frac{\sigma}{\sigma_r^*} \sqrt{J+r+\log n}. \quad (\text{F.3b})$$

In fact, the proof of [Proposition 2](#) also produces another expansion for $U\Sigma R_U - U^*\Sigma^*$ and $V\Sigma R_V - V^*\Sigma^*$, which is given below.

Proposition 3. *Under the condition of [Proposition 2](#), we can decompose*

$$U\Sigma R_U - U^*\Sigma^* = EV^* + \Delta_U, \quad \text{and} \quad (\text{F.4a})$$

$$V\Sigma R_V - V^*\Sigma^* = E^\top U^* + \Delta_V, \quad (\text{F.4b})$$

where Δ_U and Δ_V are two matrices that satisfy: there exists some universal constant $c_2 > 0$ such that for any given $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$,

$$\|(\Delta_U)_{\mathcal{I},\cdot}\| \leq c_2\sigma\sqrt{I+r+\log n} \left[\frac{\sigma}{\sigma_r^*} \sqrt{I+n_2+\log n} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right] + c_2 \frac{\sigma^2 n}{\sigma_r^*} \|U_{\mathcal{I},\cdot}^*\|, \quad (\text{F.5a})$$

$$\|(\Delta_V)_{\mathcal{J},\cdot}\| \leq c_2\sigma\sqrt{J+r+\log n} \left[\frac{\sigma}{\sigma_r^*} \sqrt{J+n_1+\log n} + \frac{\sigma^2 n_2}{\sigma_r^{*2}} \right] + c_2 \frac{\sigma^2 n}{\sigma_r^*} \|V_{\mathcal{J},\cdot}^*\|, \quad (\text{F.5b})$$

where $I = |\mathcal{I}|$ and $J = |\mathcal{J}|$. In addition, with probability exceeding $1 - O(n^{-10})$,

$$\|U\Sigma R_U - U^*\Sigma^*\| \leq c_2 \frac{\sigma^2 n}{\sigma_r^*} \|U_{\mathcal{I},\cdot}^*\| + c_2\sigma\sqrt{I+r+\log n}, \quad (\text{F.6a})$$

$$\|V\Sigma R_V - V^*\Sigma^*\| \leq c_2 \frac{\sigma^2 n}{\sigma_r^*} \|V_{\mathcal{J},\cdot}^*\| + c_2\sigma\sqrt{J+r+\log n}. \quad (\text{F.6b})$$

[Proposition 2](#) mainly concerns characterizing the subspace perturbation error. In fact, we can readily characterize the estimation error for the underlying low-rank matrix M^* , which is presented as follows.

Corollary 1. *Under the condition of [Proposition 2](#), we can decompose*

$$U\Sigma V^\top - M^* = U^*U^{*\top}E + EV^*V^{*\top} + \Phi$$

where Φ is some matrix that satisfies: there exists some universal constant $c_3 > 0$ such that for any given $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$, we have

$$\begin{aligned} \|\Phi_{\mathcal{I},\mathcal{J}}\| &\leq c_3 \frac{\sigma^2}{\sigma_r^*} \sqrt{(I+r+\log n)(J+r+\log n)} + c_3\sigma\sqrt{J+r+\log n} \left(\frac{\sigma\sqrt{J+n_1}}{\sigma_r^*} + \frac{\sigma^2 n_2}{\sigma_r^{*2}} \right) \|U_{\mathcal{I},\cdot}^*\| \\ &+ c_3\sigma\sqrt{I+r+\log n} \left(\frac{\sigma\sqrt{I+n_2}}{\sigma_r^*} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right) \|V_{\mathcal{J},\cdot}^*\| + c_3 \left(\frac{\sigma^2 n}{\sigma_r^*} + \sigma\sqrt{r+\log n} \right) \|U_{\mathcal{I},\cdot}^*\| \|V_{\mathcal{J},\cdot}^*\| \end{aligned} \quad (\text{F.7})$$

where $I = |\mathcal{I}|$ and $J = |\mathcal{J}|$. As a direct consequence, with probability exceeding $1 - O(n^{-10})$,

$$\begin{aligned} \| (U\Sigma V^\top - M^*)_{\mathcal{I},\mathcal{J}} \| &\leq c_3\sigma\sqrt{J+r+\log n} \|U_{\mathcal{I},\cdot}^*\| + c_3\sigma\sqrt{I+r+\log n} \|V_{\mathcal{J},\cdot}^*\| \\ &+ c_3 \frac{\sigma^2 n}{\sigma_r^*} \|U_{\mathcal{I},\cdot}^*\| \|V_{\mathcal{J},\cdot}^*\| + c_3 \frac{\sigma^2}{\sigma_r^*} \sqrt{(I+r+\log n)(J+r+\log n)}. \end{aligned} \quad (\text{F.8})$$

See [Appendix F.3.1](#) for the proof.

Last but not least, the following result will be useful when being applied to the causal panel data model.

Corollary 2. *Under the condition of Proposition 2, there exists some universal constant $c_4 > 0$ such that with probability exceeding $1 - O(n^{-10})$,*

$$\|\mathbf{U}_{\mathcal{I},\cdot}^{\star\top}(\mathbf{U}_{\mathcal{I},\cdot}\mathbf{R}_U - \mathbf{U}_{\mathcal{I},\cdot}^{\star})\| \leq c_4 \left[\frac{\sigma\sqrt{r + \log n}}{\sigma_r^*} + \frac{\sigma^2\sqrt{n_2(n_2 + I)}}{\sigma_r^{*2}} + \frac{\sigma^3 n_1}{\sigma_r^{*3}}\sqrt{I} \right] \|\mathbf{U}_{\mathcal{I},\cdot}^{\star}\| + c_4 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^{\star}\|^2.$$

See Appendix F.3.2 for the proof.

F.1 Proofs of Propositions 2 and 3

F.1.1 Step 1: Establishing preliminary lemmas

We first define two matrices

$$\mathbf{H}_U := \mathbf{U}^\top \mathbf{U}^* \quad \text{and} \quad \mathbf{H}_V := \mathbf{V}^\top \mathbf{V}^*.$$

Lemma 15 below shows that \mathbf{H}_U (resp. \mathbf{H}_V) is a good proxy of the rotation matrix \mathbf{R}_U (resp. \mathbf{R}_V), which depends on $\mathbf{U}^\top \mathbf{U}^*$ (resp. $\mathbf{V}^\top \mathbf{V}^*$) in a highly nonlinear way. As we will see from the proof, \mathbf{H}_U and \mathbf{H}_V are much easier to deal with. We start with applying the celebrated Wedin's sin Θ Theorem [Wed72; Che+21a], which gives us the following perturbation bounds.

Lemma 15. *Under the condition of Proposition 2, there exists some universal constant $c_5 > 0$ such that with probability exceeding $1 - O(n^{-10})$,*

$$\begin{aligned} \max \{ \|\mathbf{U}\mathbf{R}_U - \mathbf{U}^*\|, \|\mathbf{V}\mathbf{R}_V - \mathbf{V}^*\| \} &\leq c_5 \frac{\sigma}{\sigma_r^*} \sqrt{n}, \quad \text{and} \\ \max \{ \|\Sigma(\mathbf{R}_U - \mathbf{H}_U)\|, \|\Sigma(\mathbf{R}_V - \mathbf{H}_V)\| \} &\leq c_5 \frac{\sigma^2 n}{\sigma_r^*}. \end{aligned}$$

As an immediate consequence, we have

$$\frac{1}{2} \leq \sigma_i(\mathbf{H}_U) \leq 2 \quad \text{and} \quad \frac{1}{2} \leq \sigma_i(\mathbf{H}_V) \leq 2$$

for all $1 \leq i \leq r$.

See Appendix F.2.1 for the proof.

From Weyl's inequality and Lemma 19, we know that $\|\Sigma - \Sigma^*\| \leq \|\mathbf{E}\| \lesssim \sigma\sqrt{n}$. The next lemma shows that if we properly rotate Σ using \mathbf{R}_U and \mathbf{R}_V (or \mathbf{H}_U and \mathbf{H}_V), it can become even closer to Σ^* .

Lemma 16. *Under the condition of Proposition 2, there exists some universal constant $c_6 > 0$ such that with probability exceeding $1 - O(n^{-10})$,*

$$\|\mathbf{R}_U^\top \Sigma \mathbf{R}_V - \Sigma^*\| \leq c_6 \frac{\sigma^2 n}{\sigma_r^*} + c_6 \sigma \sqrt{r + \log n} \quad \text{and} \quad \|\mathbf{H}_U^\top \Sigma \mathbf{H}_V - \Sigma^*\| \leq c_6 \frac{(\sigma\sqrt{n})^3}{\sigma_r^{*2}} + c_6 \sigma \sqrt{r + \log n}.$$

See Appendix F.2.2 for the proof.

The next lemma characterizes the spectral norm bound for submatrices that are formed by a subset \mathcal{I} (resp. \mathcal{J}) of rows of $\mathbf{U}\Sigma\mathbf{H}_V - \mathbf{M}\mathbf{V}^*$ (resp. $\mathbf{V}\Sigma\mathbf{H}_U - \mathbf{M}^\top \mathbf{U}^*$), which will play a crucial role in our analysis.

Lemma 17. *Under the condition of Proposition 2, there exists some universal constant $c_7 > 0$ such that for any given $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$, we have*

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot} \Sigma \mathbf{H}_V - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| &\leq c_7 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} + c_7 \frac{\sigma^2 n_2}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|, \\ \|\mathbf{V}_{\mathcal{J},\cdot} \Sigma \mathbf{H}_U - \mathbf{M}_{\mathcal{J},\cdot}^\top \mathbf{U}^*\| &\leq c_7 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{V}_{\mathcal{J},\cdot}^*\| + c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(J+r+\log n)} + c_7 \frac{\sigma^2 n_1}{\sigma_r^*} \|\mathbf{V}_{\mathcal{J},\cdot} \mathbf{H}_V - \mathbf{V}_{\mathcal{J},\cdot}^*\|, \end{aligned}$$

where $I = |\mathcal{I}|$ and $J = |\mathcal{J}|$.

See Appendix F.2.3 for the proof.

Lemma 16 and Lemma 17 allow us to derive the following estimation error bounds.

Lemma 18. *Under the condition of Proposition 2, there exists some universal constant $c_8 > 0$ such that for any given $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$*

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| &\leq c_8 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n}, \\ \|\mathbf{V}_{\mathcal{J},\cdot} \mathbf{H}_V - \mathbf{V}_{\mathcal{J},\cdot}^*\| &\leq c_8 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{V}_{\mathcal{J},\cdot}^*\| + c_8 \frac{\sigma}{\sigma_r^*} \sqrt{J+r+\log n}. \end{aligned}$$

See Appendix F.2.4 for the proof.

F.1.2 Step 2: Deriving a crude bound

In this section, we will prove a crude bound using the lemmas in the last section. More specifically, instead of showing (F.1) with guarantees in (F.2), we first establish weaker guarantees as follows:

$$\|(\Psi_U)_{\mathcal{I},\cdot}\| \leq c_9 \frac{\sigma^2}{\sigma_r^{*2}} \sqrt{n(I+r+\log n)} + c_9 \left(\frac{\sigma^2 n}{\sigma_r^{*2}} + \frac{\sigma \sqrt{r+\log n}}{\sigma_r^*} \right) \|\mathbf{U}_{\mathcal{I},\cdot}^*\|, \quad \text{and} \quad (\text{F.9a})$$

$$\|(\Psi_V)_{\mathcal{J},\cdot}\| \leq c_9 \frac{\sigma^2}{\sigma_r^{*2}} \sqrt{n(J+r+\log n)} + c_9 \left(\frac{\sigma^2 n}{\sigma_r^{*2}} + \frac{\sigma \sqrt{r+\log n}}{\sigma_r^*} \right) \|\mathbf{V}_{\mathcal{J},\cdot}^*\| \quad (\text{F.9b})$$

for some universal constant $c_9 > 0$. The rest of this section is devoted to proving the bounds (F.9).

We first use the triangle inequality to achieve

$$\|\mathbf{U}_{\mathcal{I},\cdot} \Sigma \mathbf{R}_V - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| \leq \underbrace{\|\mathbf{U}_{\mathcal{I},\cdot} \Sigma \mathbf{H}_V - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\|}_{=:\alpha_1} + \underbrace{\|\mathbf{U}_{\mathcal{I},\cdot} \Sigma (\mathbf{H}_V - \mathbf{R}_V)\|}_{=:\alpha_2}.$$

We start by controlling α_1 . Taking Lemmas 17 and 18 collectively, we know that with probability exceeding $1 - O(n^{-10})$,

$$\begin{aligned} \alpha_1 &\leq c_7 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} + c_7 \frac{\sigma^2 n_2}{\sigma_r^*} \left(c_8 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} \right) \\ &\leq (c_7 + c_7 c_8 c_n^2) \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + (c_7 + c_7 c_n) \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} \\ &\leq 2c_7 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} \end{aligned} \quad (\text{F.10})$$

provided that $c_n > 0$ is sufficiently small. Next, we control α_2 , towards which we have

$$\alpha_2 \leq \|\mathbf{U}_{\mathcal{I},\cdot}\| \|\boldsymbol{\Sigma}(\mathbf{H}_V - \mathbf{R}_V)\| \leq c_5 \frac{\sigma^2 n}{\sigma_r^*} (3 \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n}). \quad (\text{F.11})$$

Here the last relation follows from [Lemma 15](#) and the following bound on $\|\mathbf{U}_{\mathcal{I},\cdot}\|$:

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot}\| &\stackrel{(i)}{\leq} 2 \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U\| \leq 2 \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| + 2 \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \\ &\stackrel{(ii)}{\leq} 2c_8 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} + 2 \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \\ &\stackrel{(iii)}{\leq} 3 \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n}, \end{aligned} \quad (\text{F.12})$$

where (i) follows from [Lemma 15](#), (ii) makes use of [Lemma 18](#), and (iii) holds true provided that $c_n > 0$ is sufficiently small. Putting [\(F.10\)](#) and [\(F.11\)](#) together yields

$$\|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{R}_V - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| \leq \alpha_1 + \alpha_2 \leq (2c_7 + 3c_5) \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 3c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} \quad (\text{F.13})$$

provided that $c_n > 0$ is sufficiently small. Follow the same analysis, we can also prove a similar bound on $\|\mathbf{V}_{\mathcal{J},\cdot} \boldsymbol{\Sigma} \mathbf{R}_U - \mathbf{M}_{\cdot,\mathcal{J}}^\top \mathbf{U}^*\|$. These allow us to write

$$\mathbf{U} \boldsymbol{\Sigma} \mathbf{R}_V = \mathbf{M} \mathbf{V}^* + \boldsymbol{\Delta}_U \quad \text{and} \quad \mathbf{V} \boldsymbol{\Sigma} \mathbf{R}_U = \mathbf{M}^\top \mathbf{U}^* + \boldsymbol{\Delta}_V, \quad (\text{F.14})$$

where for any $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$ we have

$$\|(\boldsymbol{\Delta}_U)_{\mathcal{I},\cdot}\| \leq (2c_7 + 3c_5) \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 3c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)}, \quad (\text{F.15a})$$

$$\|(\boldsymbol{\Delta}_V)_{\mathcal{J},\cdot}\| \leq (2c_7 + 3c_5) \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{V}_{\mathcal{J},\cdot}^*\| + 3c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(J+r+\log n)}. \quad (\text{F.15b})$$

Besides, in light of [Lemma 16](#) we can write

$$\boldsymbol{\Sigma}^* = \mathbf{R}_U^\top \boldsymbol{\Sigma} \mathbf{R}_V + \boldsymbol{\Delta}_\Sigma \quad \text{where} \quad \|\boldsymbol{\Delta}_\Sigma\| \leq c_6 \frac{\sigma^2 n}{\sigma_r^*} + c_6 \sigma \sqrt{r+\log n}. \quad (\text{F.16})$$

The expansions [\(F.14\)](#) and [\(F.16\)](#) yield

$$\begin{aligned} \mathbf{U} \mathbf{R}_U \boldsymbol{\Sigma}^* &= \mathbf{U} \boldsymbol{\Sigma} \mathbf{R}_V + \mathbf{U} (\mathbf{R}_U \boldsymbol{\Sigma}^* - \boldsymbol{\Sigma} \mathbf{R}_V) = \mathbf{M} \mathbf{V}^* + \boldsymbol{\Delta}_U + \mathbf{U} \mathbf{R}_U (\boldsymbol{\Sigma}^* - \mathbf{R}_U^\top \boldsymbol{\Sigma} \mathbf{R}_V) \\ &= (\mathbf{M}^* + \mathbf{E}) \mathbf{V}^* + \boldsymbol{\Delta}_U + \mathbf{U} \mathbf{R}_U \boldsymbol{\Delta}_\Sigma = \mathbf{U}^* \boldsymbol{\Sigma}^* + \mathbf{E} \mathbf{V}^* + \boldsymbol{\Delta}_U + \mathbf{U} \mathbf{R}_U \boldsymbol{\Delta}_\Sigma, \end{aligned}$$

which further leads to

$$\mathbf{U} \mathbf{R}_U = \mathbf{U}^* + \mathbf{E} \mathbf{V}^* (\boldsymbol{\Sigma}^*)^{-1} + \underbrace{\boldsymbol{\Delta}_U (\boldsymbol{\Sigma}^*)^{-1} + \mathbf{U} \mathbf{R}_U \boldsymbol{\Delta}_\Sigma (\boldsymbol{\Sigma}^*)^{-1}}_{=:\boldsymbol{\Psi}_U}. \quad (\text{F.17})$$

For any $\mathcal{I} \subseteq [n_1]$, we can bound

$$\begin{aligned}
\|(\Psi_U)_{\mathcal{I},\cdot}\| &\leq \frac{1}{\sigma_r^*} \|(\Delta_U)_{\mathcal{I},\cdot}\| + \frac{1}{\sigma_r^*} \|U_{\mathcal{I},\cdot} R_U\| \|\Delta_\Sigma\| \\
&\stackrel{(i)}{\leq} (2c_7 + 3c_5) \frac{\sigma^2 n}{\sigma_r^{*2}} \|U_{\mathcal{I},\cdot}^*\| + 3c_7 \frac{\sigma^2 \sqrt{n(I+r+\log n)}}{\sigma_r^{*2}} \\
&\quad + (3\|U_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma \sqrt{I+r+\log n}}{\sigma_r^*}) (c_6 \frac{\sigma^2 n}{\sigma_r^{*2}} + c_6 \frac{\sigma \sqrt{r+\log n}}{\sigma_r^*}) \\
&\stackrel{(ii)}{\leq} c_9 \frac{\sigma^2}{\sigma_r^{*2}} \sqrt{n(I+r+\log n)} + c_9 \left(\frac{\sigma^2 n}{\sigma_r^{*2}} + \frac{\sigma \sqrt{r+\log n}}{\sigma_r^*} \right) \|U_{\mathcal{I},\cdot}^*\|. \tag{F.18}
\end{aligned}$$

Here (i) follows from (F.15a), (F.12) and (F.16), while (ii) holds true if we take $c_9 = 3c_7 + 3c_5 + 3c_6$ and $c_n > 0$ is sufficiently small. The other bound (F.9b) can be verified in the similar way.

F.1.3 Step 3: Refining the bounds

In this section we will improve the bounds (F.9) to (F.2). Note that the suboptimality of (F.9) is partly because we start our analysis with an application of Wedin's $\sin \Theta$ Theorem, which gives the following result in Lemma 15:

$$\max \{ \|UR_U - U^*\|, \|VR_V - V^*\| \} \leq c_5 \frac{\sigma}{\sigma_r^*} \sqrt{n}. \tag{F.19}$$

This bound does not capture any potential imbalances of the data matrix, namely when n_1 and n_2 are not on the same order. However, in light of Lemmas 15 and 18, we can sharpen the above estimate to the following: there exists some universal constant $c'_5 > 0$ such that for any given $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$

$$\|U_{\mathcal{I},\cdot} R_U - U_{\mathcal{I},\cdot}^*\| \leq c'_5 \frac{\sigma^2 n}{\sigma_r^{*2}} \|U_{\mathcal{I},\cdot}^*\| + c'_5 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n}, \tag{F.20a}$$

$$\|V_{\mathcal{J},\cdot} R_V - V_{\mathcal{J},\cdot}^*\| \leq c'_5 \frac{\sigma^2 n}{\sigma_r^{*2}} \|V_{\mathcal{J},\cdot}^*\| + c'_5 \frac{\sigma}{\sigma_r^*} \sqrt{J+r+\log n}. \tag{F.20b}$$

See Appendix F.2.5 for the proof of the claim (F.20). By taking $\mathcal{I} = [n_1]$ and $\mathcal{J} = [n_2]$, we have

$$\|UR_U - U^*\| \leq c'_5 \frac{\sigma^2 n}{\sigma_r^{*2}} + 2c'_5 \frac{\sigma}{\sigma_r^*} \sqrt{n_1 + \log n}, \tag{F.21a}$$

$$\|VR_V - V^*\| \leq c'_5 \frac{\sigma^2 n}{\sigma_r^{*2}} + 2c'_5 \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n}, \tag{F.21b}$$

which improves the original bound (F.19). We can also obtain the following sharper bounds:

$$\|\Sigma(R_U - H_U)\| \leq c'_5 (\sigma \sqrt{n_1 + \log n} + \frac{\sigma^2 n_2}{\sigma_r^*}) \frac{\sigma}{\sigma_r^*} \sqrt{n_1 + \log n}, \quad \text{and} \tag{F.22a}$$

$$\|\Sigma(R_V - H_V)\| \leq c'_5 (\sigma \sqrt{n_2 + \log n} + \frac{\sigma^2 n_1}{\sigma_r^*}) \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n}. \tag{F.22b}$$

See Appendix F.2.5 for the proof of the claim (F.22).

Based on these refined perturbation bounds, we can sharpen the bounds in [Lemma 17](#) to

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{H}_{\mathbf{V}} - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| &\leq c'_7 \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n} \right)^2 \sigma_r^* \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \\ &\quad + c'_7 \sigma \sqrt{I + r + \log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{I + n_2 + \log n} \right), \end{aligned} \quad (\text{F.23a})$$

$$\begin{aligned} \|\mathbf{V}_{\mathcal{J},\cdot} \boldsymbol{\Sigma} \mathbf{H}_{\mathbf{U}} - \mathbf{M}_{\mathcal{J},\cdot}^\top \mathbf{U}^*\| &\leq c'_7 \left(\frac{\sigma^2 n_2}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{n_1 + \log n} \right)^2 \sigma_r^* \|\mathbf{V}_{\mathcal{J},\cdot}^*\|_2 \\ &\quad + c'_7 \sigma \sqrt{J + r + \log n} \left(\frac{\sigma^2 n_2}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{J + n_1 + \log n} \right), \end{aligned} \quad (\text{F.23b})$$

where $c'_7 > 0$ is some universal constant. See [Appendix F.2.5](#). for the proof of the claim [\(F.23\)](#).

With this set of refined bounds in place, we are ready to finish the proof of [Proposition 2](#) and [Proposition 3](#). Recall the definitions of $\boldsymbol{\Delta}_{\mathbf{U}}$ and $\boldsymbol{\Psi}_{\mathbf{U}}$ from equations [\(F.14\)](#) and [\(F.17\)](#) respectively. We have

$$\begin{aligned} \|(\boldsymbol{\Delta}_{\mathbf{U}})_{\mathcal{I},\cdot}\| &= \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{R}_{\mathbf{V}} - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| \leq \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{H}_{\mathbf{V}} - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} (\mathbf{H}_{\mathbf{V}} - \mathbf{R}_{\mathbf{V}})\| \\ &\leq \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{H}_{\mathbf{V}} - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot}\| \|\boldsymbol{\Sigma} (\mathbf{H}_{\mathbf{V}} - \mathbf{R}_{\mathbf{V}})\| \\ &\stackrel{(i)}{\leq} c'_7 \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n} \right)^2 \sigma_r^* \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c'_7 \sigma \sqrt{I + r + \log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{I + n_2 + \log n} \right) \\ &\quad + (3 \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n}) \cdot c'_5 (\sigma \sqrt{n_2 + \log n} + \frac{\sigma^2 n_1}{\sigma_r^*}) \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n} \\ &\stackrel{(ii)}{\leq} 2c'_7 \sigma \sqrt{I + r + \log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{I + n_2 + \log n} \right) + (4c'_7 + 7c'_5) \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\|. \end{aligned} \quad (\text{F.24})$$

Here (i) utilizes from [\(F.23a\)](#), [\(F.12\)](#) and [\(F.22b\)](#), while (ii) holds provided that $\sigma \sqrt{n} \leq c_n \sigma_r^*$ for some sufficiently small $c_n > 0$. By taking $c_2 = 4c'_7 + 7c'_5$, we have proved [\(F.5a\)](#), and [\(F.5b\)](#) can be verified similarly. This finishes the proof of [Proposition 3](#). We can also improve [\(F.18\)](#) to

$$\begin{aligned} \|(\boldsymbol{\Psi}_{\mathbf{U}})_{\mathcal{I},\cdot}\| &\leq \frac{1}{\sigma_r^*} \|(\boldsymbol{\Delta}_{\mathbf{U}})_{\mathcal{I},\cdot}\| + \frac{1}{\sigma_r^*} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n}) \|\boldsymbol{\Delta}_{\boldsymbol{\Sigma}}\| \\ &\stackrel{(a)}{\leq} c_2 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_2 \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{I + n_2 + \log n} \right) \\ &\quad + \frac{1}{\sigma_r^*} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n}) \cdot c_6 \left(\frac{\sigma^2 n}{\sigma_r^*} + \sigma \sqrt{r + \log n} \right) \\ &\stackrel{(b)}{\leq} c_1 \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n} \left(\frac{\sigma}{\sigma_r^*} \sqrt{I + n_2 + \log n} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right) + c_1 \left(\frac{\sigma^2 n}{\sigma_r^{*2}} + \frac{\sigma \sqrt{r + \log n}}{\sigma_r^*} \right) \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \end{aligned}$$

where we take $c_1 = c_2 + 2c_6$. Here step (a) follows from [\(F.5a\)](#) and [\(F.16\)](#), while step (b) holds provided that $\sigma \sqrt{n} \leq c_n \sigma_r^*$ for some sufficiently small $c_n > 0$. This has established [\(F.2a\)](#). Similarly we can verify [\(F.2b\)](#), which completes the proof of [Proposition 2](#).

F.2 Proof of auxiliary lemmas for matrix denoising

This section provides omitted proofs in [Appendix F.1](#).

F.2.1 Proof of Lemma 15

To begin with, we find it convenient to write the full SVD of M as

$$M = \begin{bmatrix} \mathbf{U} & \mathbf{U}_\perp \end{bmatrix} \begin{bmatrix} \boldsymbol{\Sigma} & \mathbf{0} \\ \mathbf{0} & \boldsymbol{\Sigma}_\perp \end{bmatrix} \begin{bmatrix} \mathbf{V} & \mathbf{V}_\perp \end{bmatrix}^\top = \mathbf{U}\boldsymbol{\Sigma}\mathbf{V}^\top + \mathbf{U}_\perp\boldsymbol{\Sigma}_\perp\mathbf{V}_\perp^\top, \quad (\text{F.25})$$

where $\mathbf{U}_\perp \in \mathbb{R}^{n_1 \times (n_1 - r)}$ and $\mathbf{V}_\perp \in \mathbb{R}^{n_2 \times (n_2 - r)}$ are the orthogonal complements of \mathbf{U} and \mathbf{V} , and $\boldsymbol{\Sigma}_\perp \in \mathbb{R}^{(n_1 - r) \times (n_2 - r)}$ contains the smaller singular values (which is not necessarily a diagonal matrix). We also define where $\mathbf{U}_\perp^* \in \mathbb{R}^{n_1 \times (n_1 - r)}$ and $\mathbf{V}_\perp^* \in \mathbb{R}^{n_2 \times (n_2 - r)}$ to be the orthogonal complements of \mathbf{U}^* and \mathbf{V}^* .

Lemma 19 shows that with probability exceeding $1 - O(n^{-10})$,

$$\|\mathbf{E}\| \leq C_g \sigma \sqrt{n}. \quad (\text{F.26})$$

Then Weyl's inequality immediately gives

$$\frac{1}{2}\sigma_r^* \leq \sigma_r^* - \|\mathbf{E}\| \leq \sigma_r(\mathbf{M}) \leq \sigma_1^* + \|\mathbf{E}\| \leq 2\sigma_1^* \quad (\text{F.27})$$

since $\|\mathbf{E}\| \leq C_g c_n \sigma_r^* \leq \sigma_r^*/2$ with the proviso that c_n is sufficiently small. In view of Wedin's $\sin \Theta$ Theorem (see [Wed72] or [Che+21a, Theorem 2.3.1]), we have

$$\frac{1}{\sqrt{2}}\|\mathbf{U}\mathbf{R}_U - \mathbf{U}^*\| \leq \|\mathbf{U}\mathbf{U}^\top - \mathbf{U}^*\mathbf{U}^{*\top}\| = \|\mathbf{U}^{*\top}\mathbf{U}_\perp\| \leq \frac{\|\mathbf{E}\|}{\sigma_r^* - \|\mathbf{E}\|} \leq 2C_g \frac{\sigma}{\sigma_r^*} \sqrt{n}, \quad (\text{F.28})$$

where the last step follows from $\sigma_{r+1}(\mathbf{M}^*) = 0$ and (F.27). The first three quantities in the above formula are all metrics between subspace \mathbf{U} and \mathbf{U}^* and are equivalent up to constant multiple; see [Che+21a, Section 2.2.3] for detailed discussions. Similarly we can show that

$$\frac{1}{\sqrt{2}}\|\mathbf{V}\mathbf{R}_V - \mathbf{V}^*\| \leq \|\mathbf{V}\mathbf{V}^\top - \mathbf{V}^*\mathbf{V}^{*\top}\| = \|\mathbf{V}^{*\top}\mathbf{V}_\perp\| \leq 2C_g \frac{\sigma}{\sigma_r^*} \sqrt{n}. \quad (\text{F.29})$$

Now we move on to establishing the proximity between \mathbf{H}_U and \mathbf{R}_U . Since \mathbf{U} and \mathbf{U}^* both have orthonormal columns, the SVD of $\mathbf{H}_U = \mathbf{U}^\top \mathbf{U}^*$ can be written as

$$\mathbf{H}_U = \mathbf{X}(\cos \Theta)\mathbf{Y}^\top,$$

where $\mathbf{X}, \mathbf{Y} \in \mathbb{R}^{r \times r}$ are orthogonal matrices and $\Theta = \text{diag}\{\theta_1, \dots, \theta_r\}$ is a diagonal matrix where $0 \leq \theta_1 \leq \dots \leq \theta_r \leq \pi/2$ are the principal (or canonical) angles between the two subspaces \mathbf{U} and \mathbf{U}^* (see [Che+21a, Section 2.1] for a systematic introduction). Here $\cos \Theta = \text{diag}\{\cos \theta_1, \dots, \cos \theta_r\}$ applies the cosine function to the diagonal entries, and $\sin \Theta = \text{diag}\{\sin \theta_1, \dots, \sin \theta_r\}$ is defined analogously. Then by definition, we have $\mathbf{R}_U = \text{sgn}(\mathbf{H}_U) = \mathbf{X}\mathbf{Y}^\top$. This immediately gives

$$\mathbf{R}_U - \mathbf{H}_U = \mathbf{X}\mathbf{Y}^\top - \mathbf{X}(\cos \Theta)\mathbf{Y}^\top = \mathbf{X}(\mathbf{I}_r - \cos \Theta)\mathbf{Y}^\top = 2\mathbf{X} \sin^2(\Theta/2)\mathbf{Y}^\top.$$

On the other hand, we have

$$\mathbf{U}^\top \mathbf{U}_\perp^* (\mathbf{U}_\perp^\top \mathbf{U}_\perp^*) = \mathbf{U}^\top (\mathbf{I}_{n_1} - \mathbf{U}^* \mathbf{U}^{*\top}) \mathbf{U} = \mathbf{I}_r - \mathbf{H}_U \mathbf{H}_U^\top = \mathbf{I}_r - \mathbf{X}(\cos \Theta)^2 \mathbf{X}^\top = \mathbf{X}(\sin \Theta)^2 \mathbf{X}^\top.$$

This suggest that, there exists a matrix $\mathbf{Z} \in \mathbb{R}^{(n_1-r) \times r}$ with orthonormal columns such that

$$\mathbf{U}^\top \mathbf{U}_\perp^* = \mathbf{X} (\sin \Theta) \mathbf{Z}^\top.$$

Taking the above two equations collectively yields

$$\Sigma (\mathbf{R}_U - \mathbf{H}_U) = 2\Sigma \mathbf{X} \sin^2(\Theta/2) \mathbf{Y}^\top = 2\Sigma \mathbf{U}^\top \mathbf{U}_\perp^* \mathbf{Z} (\sin \Theta)^{-1} \sin(\Theta/2) \sin(\Theta/2) \mathbf{Y}^\top. \quad (\text{F.30})$$

Here we define $0/0 = 0$ such that in the above equation,

$$(\sin \Theta)^{-1} \sin(\Theta/2) = \text{diag} \left\{ \frac{\sin(\theta_1/2)}{\sin \theta_1}, \dots, \frac{\sin(\theta_r/2)}{\sin \theta_r} \right\}$$

is always well defined. In view of the relation between the principal angles and subspace distances (see e.g. [Che+21a, Lemma 2.5]) and (F.28), we have

$$\|\sin \Theta\| = \|\mathbf{U}\mathbf{U}^\top - \mathbf{U}^*\mathbf{U}^{*\top}\| \leq 2C_g \frac{\sigma}{\sigma_r^*} \sqrt{n}. \quad (\text{F.31})$$

Since $\sin(\theta/2) \leq \sin(\theta)$ for any $0 \leq \theta \leq \pi/2$, we know that

$$\|\sin(\Theta/2)\| \leq \|\sin \Theta\| \quad \text{and} \quad \|(\sin \Theta)^{-1} \sin(\Theta/2)\| \leq 1. \quad (\text{F.32})$$

These relations allow us to obtain

$$\begin{aligned} \|\Sigma (\mathbf{R}_U - \mathbf{H}_U)\| &\stackrel{(i)}{\leq} 2 \|\Sigma \mathbf{U}^\top \mathbf{U}_\perp^*\| \|\mathbf{Z}\| \|(\sin \Theta)^{-1} \sin(\Theta/2)\| \|\sin(\Theta/2)\| \\ &\stackrel{(ii)}{\leq} 2 \|\Sigma \mathbf{U}^\top \mathbf{U}_\perp^*\| \|\sin \Theta\| \stackrel{(iii)}{\leq} 4C_g \frac{\sigma}{\sigma_r^*} \sqrt{n} \|\Sigma \mathbf{U}^\top \mathbf{U}_\perp^*\|, \end{aligned} \quad (\text{F.33})$$

where (i) follows from (F.30), (ii) utilizes (F.32), and (iii) makes use of (F.31). Then we write

$$\begin{aligned} \Sigma \mathbf{U}^\top \mathbf{U}_\perp^* &= \mathbf{V}^\top (\mathbf{U} \Sigma \mathbf{V}^\top)^\top \mathbf{U}_\perp^* = \mathbf{V}^\top (\mathbf{M} - \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top)^\top \mathbf{U}_\perp^* \\ &= \mathbf{V}^\top (\mathbf{U}^* \Sigma^* \mathbf{V}^{*\top} + \mathbf{E} - \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top)^\top \mathbf{U}_\perp^* = \mathbf{V}^\top \mathbf{E}^\top \mathbf{U}_\perp^{*\top}. \end{aligned}$$

Here the last step holds since $\mathbf{U}^{*\top} \mathbf{U}_\perp^* = \mathbf{0}$ and $\mathbf{V}^\top \mathbf{V}_\perp = \mathbf{0}$. Therefore we have

$$\begin{aligned} \|\Sigma \mathbf{U}^\top \mathbf{U}_\perp^*\| &\leq \|\mathbf{E}\mathbf{V}\| \leq \|\mathbf{E}\mathbf{V}^*\| + \|\mathbf{E}(\mathbf{V}\mathbf{R}_V - \mathbf{V}^*)\| \leq \|\mathbf{E}\mathbf{V}^*\| + \|\mathbf{E}\| \|\mathbf{V}\mathbf{R}_V - \mathbf{V}^*\| \\ &\stackrel{(i)}{\leq} C_g \sigma \sqrt{n_1 + \log n} + 2\sqrt{2}C_g \frac{\sigma}{\sigma_r^*} \sqrt{n} \cdot C_g \sigma \sqrt{n} \leq C_g \sigma \sqrt{n_1 + \log n} + 3C_g^2 \frac{\sigma^2 n}{\sigma_r^*}. \end{aligned} \quad (\text{F.34})$$

Here (i) follows from Lemma 19, (F.26) and (F.29). Taking (F.33) and (F.34) collectively yields

$$\|\Sigma (\mathbf{R}_U - \mathbf{H}_U)\| \leq 4C_g \frac{\sigma}{\sigma_r^*} \sqrt{n} (C_g \sigma \sqrt{n_1 + \log n} + 3C_g^2 \frac{\sigma^2 n}{\sigma_r^*}) \leq 8C_g^2 \frac{\sigma^2 n}{\sigma_r^*}$$

provided that c_n is sufficiently small. This taken collectively with (F.27) yields

$$\|\mathbf{H}_U - \mathbf{R}_U\| \leq \frac{\|\Sigma (\mathbf{R}_U - \mathbf{H}_U)\|}{\sigma_r(\mathbf{M})} \leq 16C_g^2 \frac{\sigma^2 n}{\sigma_r^{*2}}. \quad (\text{F.35})$$

Given that \mathbf{R}_U is an orthogonal matrix, all its r singular values are one. Thus by Weyl's inequality,

$$\frac{1}{2} \leq \sigma_r(\mathbf{R}_U) - \|\mathbf{H}_U - \mathbf{R}_U\| \leq \sigma_r(\mathbf{H}_U) \leq \sigma_1(\mathbf{H}_U) \leq \sigma_1(\mathbf{R}_U) + \|\mathbf{H}_U - \mathbf{R}_U\| \leq 2,$$

provided that $\sigma\sqrt{n} \leq c_n\sigma_r^*$ with a sufficiently small $c_n > 0$. Similarly, we can show that

$$\|\Sigma(\mathbf{R}_V - \mathbf{H}_V)\| \leq 8C_g^2 \frac{\sigma^2 n}{\sigma_r^*}, \quad \|\mathbf{H}_V - \mathbf{R}_V\| \leq 16C_g^2 \frac{\sigma^2 n}{\sigma_r^{*2}}, \quad \text{and} \quad \frac{1}{2} \leq \sigma_r(\mathbf{H}_U) \leq \sigma_1(\mathbf{H}_U) \leq 2.$$

Let $c_5 = 16C_g^2$ to complete the proof.

F.2.2 Proof of Lemma 16

Let us begin by upper bounding the error of interest using the triangle inequality

$$\|\mathbf{R}_U^\top \Sigma \mathbf{R}_V - \Sigma^*\| \leq \underbrace{\|\mathbf{R}_U^\top \Sigma \mathbf{R}_V - \mathbf{H}_U^\top \Sigma \mathbf{H}_V\|}_{=: \alpha_1} + \underbrace{\|\mathbf{H}_U^\top \Sigma \mathbf{H}_V - \mathbf{U}^{*\top} \mathbf{M} \mathbf{V}^*\|}_{=: \alpha_2} + \underbrace{\|\mathbf{U}^{*\top} \mathbf{M} \mathbf{V}^* - \Sigma^*\|}_{=: \alpha_3}.$$

It then suffices to bound the three terms α_1 , α_2 and α_3 .

We first invoke Lemma 15 to bound α_1 , which gives

$$\begin{aligned} \alpha_1 &\leq \|(\mathbf{H}_U - \mathbf{R}_U)^\top \Sigma \mathbf{H}_V\| + \|\mathbf{R}_U^\top \Sigma (\mathbf{H}_V - \mathbf{R}_V)\| \\ &\leq 2\|\Sigma(\mathbf{H}_U - \mathbf{R}_U)\| + \|\Sigma(\mathbf{H}_V - \mathbf{R}_V)\| \leq 3c_5 \frac{\sigma^2 n}{\sigma_r^*}. \end{aligned}$$

Regarding α_2 , recall the full SVD of \mathbf{M} in (F.25), which allows us to write

$$\mathbf{H}_U^\top \Sigma \mathbf{H}_V - \mathbf{U}^{*\top} \mathbf{M} \mathbf{V}^* = \mathbf{U}^{*\top} (\mathbf{U} \Sigma \mathbf{V}^\top - \mathbf{M}) \mathbf{V}^* = \mathbf{U}^{*\top} \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top \mathbf{V}^*.$$

In view of Weyl's inequality and (F.26), we have

$$\|\Sigma_\perp\| \leq \|\mathbf{E}\| \leq C_g \sigma \sqrt{n}. \quad (\text{F.36})$$

Taking together (F.36), (F.28) and (F.29) gives

$$\alpha_2 = \|\mathbf{U}^{*\top} \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top \mathbf{V}^*\| \leq \|\mathbf{U}^{*\top} \mathbf{U}_\perp\| \|\Sigma_\perp\| \|\mathbf{V}^{*\top} \mathbf{V}_\perp\| \leq 4C_g^3 \frac{(\sigma\sqrt{n})^3}{\sigma_r^{*2}}.$$

To control α_3 , we first observe that

$$\mathbf{U}^{*\top} \mathbf{M} \mathbf{V}^* - \Sigma^* = \mathbf{U}^{*\top} \mathbf{M}^* \mathbf{V}^* + \mathbf{U}^{*\top} \mathbf{E} \mathbf{V}^* - \Sigma^* = \Sigma^* + \mathbf{U}^{*\top} \mathbf{E} \mathbf{V}^* - \Sigma^* = \mathbf{U}^{*\top} \mathbf{E} \mathbf{V}^*.$$

It suffices to bound the spectral norm of $\mathbf{U}^{*\top} \mathbf{E} \mathbf{V}^*$. Invoke Lemma 19 shows that with probability exceeding $1 - O(n^{-10})$,

$$\alpha_3 = \|\mathbf{U}^{*\top} \mathbf{E} \mathbf{V}^*\| \leq C_g \sigma \sqrt{r + \log n}. \quad (\text{F.37})$$

Taking the bounds on α_1 , α_2 and α_3 collectively yields

$$\|\mathbf{R}_U^\top \Sigma \mathbf{R}_V - \Sigma^*\| \leq 3c_5 \frac{\sigma^2 n}{\sigma_r^*} + 4C_g^3 \frac{(\sigma\sqrt{n})^3}{\sigma_r^{*2}} + C_g \sigma \sqrt{r + \log n} \leq c_6 \frac{\sigma^2 n}{\sigma_r^*} + c_6 \sigma \sqrt{r + \log n}$$

as long as $c_6 \geq 3c_5 + 4C_g^3c_n + C_g$. In addition, we also have the following by-product:

$$\begin{aligned} \|\mathbf{H}_U^\top \Sigma \mathbf{H}_V - \Sigma^*\| &\leq \|\mathbf{H}_U^\top \Sigma \mathbf{H}_V - \mathbf{U}^{*\top} \mathbf{M} \mathbf{V}^*\| + \|\mathbf{U}^{*\top} \mathbf{M} \mathbf{V}^* - \Sigma^*\| = \alpha_2 + \alpha_3 \\ &\leq 4C_g^3 \frac{(\sigma\sqrt{n})^3}{\sigma_r^{*2}} + C_g \sigma \sqrt{r + \log n} \leq c_6 \frac{(\sigma\sqrt{n})^3}{\sigma_r^{*2}} + c_6 \sigma \sqrt{r + \log n}, \end{aligned}$$

which holds provided that $c_6 \geq 4C_g^3 + C_g$.

F.2.3 Proof of Lemma 17

Let $I = |\mathcal{I}|$ be the cardinality of \mathcal{I} . We start by the following decomposition

$$\|\mathbf{U}_{\mathcal{I}, \Sigma} \mathbf{H}_V - \mathbf{M}_{\mathcal{I}, \mathbf{V}^*}\| = \|\mathbf{M}_{\mathcal{I}, (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*)}\| \leq \underbrace{\|\mathbf{M}_{\mathcal{I}, (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*)}\|}_{=:\alpha} + \underbrace{\|\mathbf{E}_{\mathcal{I}, (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*)}\|}_{=:\beta},$$

where the first relation follows from $\mathbf{U}\Sigma = \mathbf{M}\mathbf{V}$. The rest of the proof is devoted to bounding α and β .

Step 1: bounding α . To begin with, we first notice that

$$\alpha = \|\mathbf{U}_{\mathcal{I}, \Sigma}^* \Sigma^* \mathbf{V}^{*\top} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*)\| \leq \|\mathbf{U}_{\mathcal{I}, \Sigma}^*\| \|\Sigma^* \mathbf{V}^{*\top} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*)\|.$$

Using the full SVD of \mathbf{M} in (F.25), we can write

$$\begin{aligned} \Sigma^* \mathbf{V}^{*\top} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*) &= \Sigma^* \mathbf{V}^{*\top} (\mathbf{V} \mathbf{V}^\top \mathbf{V}^* - \mathbf{V}^*) = \mathbf{U}^{*\top} \mathbf{M}^* (\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^* \\ &= \mathbf{U}^{*\top} (\mathbf{M} - \mathbf{E}) (\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^* \\ &= \mathbf{U}^{*\top} (\mathbf{U} \Sigma \mathbf{V}^\top + \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top - \mathbf{E}) (\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^* \\ &= \mathbf{U}^{*\top} \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top \mathbf{V}^* - \mathbf{U}^{*\top} \mathbf{E} (\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^*. \end{aligned}$$

Invoke (F.28), (F.29) and (F.36) to bound the spectral norm of the first matrix

$$\|\mathbf{U}^{*\top} \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top \mathbf{V}^*\| \leq \|\mathbf{U}^{*\top} \mathbf{U}_\perp\| \|\Sigma_\perp\| \|\mathbf{V}^{*\top} \mathbf{V}_\perp\| \leq 4C_g^3 \frac{(\sigma\sqrt{n})^3}{\sigma_r^{*2}}. \quad (\text{F.38})$$

The spectral norm of the second matrix is upper bounded by

$$\begin{aligned} \|\mathbf{U}^{*\top} \mathbf{E} (\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^*\| &\leq \|\mathbf{U}^{*\top} \mathbf{E}\| \|\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}\| \|\mathbf{V}^*\| = \|\mathbf{U}^{*\top} \mathbf{E}\| \|\mathbf{V} \mathbf{H}_V - \mathbf{V}^*\| \\ &\leq C_g \sigma \sqrt{n_2 + \log n} \cdot 2C_g \frac{\sigma}{\sigma_r^*} \sqrt{n} = 2C_g^2 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(n_2 + \log n)}, \end{aligned} \quad (\text{F.39})$$

where the penultimate relation follows from Lemma 19 and a consequence of (F.29):

$$\|\mathbf{V} \mathbf{H}_V - \mathbf{V}^*\| = \|(\mathbf{V} \mathbf{V}^\top - \mathbf{V}^* \mathbf{V}^{*\top}) \mathbf{V}^*\| \leq \|\mathbf{V} \mathbf{V}^\top - \mathbf{V}^* \mathbf{V}^{*\top}\| \leq 2C_g \frac{\sigma}{\sigma_r^*} \sqrt{n}. \quad (\text{F.40})$$

Therefore by the triangle inequality,

$$\begin{aligned} \|\Sigma^* \mathbf{V}^{*\top} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*)\| &\leq \|\mathbf{U}^{*\top} \mathbf{U}_\perp \Sigma_\perp \mathbf{V}_\perp^\top \mathbf{V}^*\| + \|\mathbf{U}^{*\top} \mathbf{E} (\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^*\| \\ &\leq 4C_g^3 \frac{(\sigma\sqrt{n})^3}{\sigma_r^{*2}} + 2C_g^2 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(n_2 + \log n)} \leq 3C_g^2 \frac{\sigma^2 n}{\sigma_r^*} \end{aligned} \quad (\text{F.41})$$

provided that $\sigma\sqrt{n} \leq c_n\sigma_r^*$ with a sufficiently small $c_n > 0$. Hence we obtain

$$\alpha \leq 3C_g^2 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\|. \quad (\text{F.42})$$

Step 2: introducing leave- I -out estimates. The key technical difficulty for bounding β is the statistical dependency between $\mathbf{E}_{\mathcal{I},\cdot}$ and $\mathbf{V}\mathbf{H}_{\mathbf{V}} - \mathbf{V}^*$. Towards this, we define an auxiliary data matrix

$$\mathbf{M}^{(\mathcal{I})} := \mathbf{M}^* + \mathcal{P}_{-\mathcal{I},\cdot}(\mathbf{E}),$$

where the operator $\mathcal{P}_{-\mathcal{I},\cdot} : \mathbb{R}^{n_1 \times n_2} \rightarrow \mathbb{R}^{n_1 \times n_2}$ is defined as

$$[\mathcal{P}_{-\mathcal{I},\cdot}(\mathbf{A})]_{i,j} = \begin{cases} A_{i,j}, & \text{if } i \in \mathcal{I}, \\ 0, & \text{otherwise.} \end{cases}$$

We also define the auxiliary estimates $\mathbf{U}^{(\mathcal{I})}$, $\mathbf{V}^{(\mathcal{I})}$, $\mathbf{H}_{\mathbf{U}}^{(\mathcal{I})}$ and $\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})}$ w.r.t. the data matrix $\mathbf{M}^{(\mathcal{I})}$, in the same way as how we define \mathbf{U} , \mathbf{V} , $\mathbf{H}_{\mathbf{U}}$ and $\mathbf{H}_{\mathbf{V}}$ w.r.t. the data matrix \mathbf{M} . Introducing these auxiliary estimates have the following benefits:

- By construction $\mathbf{M}^{(\mathcal{I})}$ is independent of $\mathbf{E}_{\mathcal{I},\cdot}$, therefore all these auxiliary estimates are independent of $\mathbf{E}_{\mathcal{I},\cdot}$. It is easier to bound $\|\mathbf{E}_{\mathcal{I},\cdot}(\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}^*)\|$ than directly working on β .
- Since $\mathbf{M}^{(\mathcal{I})}$ is constructed by leaving a small amount of noise information from \mathbf{M} , we can expect that these auxiliary estimates are close to the original estimates, at least when I is small.

In what follows, we will exploit these statistical benefits to bound β .

Step 3: bounding β . We start by applying the triangle inequality to obtain

$$\beta \leq \underbrace{\|\mathbf{E}_{\mathcal{I},\cdot}(\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}^*)\|}_{=: \beta_1} + \underbrace{\|\mathbf{E}_{\mathcal{I},\cdot}(\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}\mathbf{H}_{\mathbf{V}})\|}_{=: \beta_2}.$$

Regarding β_1 , since $\mathbf{V}^{(\mathcal{I})}\mathbf{H}^{(\mathcal{I})} - \mathbf{V}^*$ and $\mathbf{E}_{\mathcal{I},\cdot}$ are independent, we can condition on $\mathbf{V}^{(\mathcal{I})}\mathbf{H}^{(\mathcal{I})} - \mathbf{V}^*$ and invoke [Lemma 19](#), thereby finding that

$$\begin{aligned} \beta_1 &\stackrel{(i)}{\leq} C_g\sigma\sqrt{I+r+\log n}\|\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}^*\| \\ &\leq C_g\sigma\sqrt{I+r+\log n}\|\mathbf{V}\mathbf{H}_{\mathbf{V}} - \mathbf{V}^*\| + C_g\sigma\sqrt{I+r+\log n}\|\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}\mathbf{H}_{\mathbf{V}}\| \\ &\stackrel{(ii)}{\leq} 2C_g^2\frac{\sigma^2}{\sigma_r^*}\sqrt{n(I+r+\log n)} + C_g\sigma\sqrt{I+r+\log n}\|\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}\mathbf{H}_{\mathbf{V}}\|. \end{aligned} \quad (\text{F.43})$$

Here (i) holds with probability exceeding $1 - O(n^{-10})$ without conditioning, while (ii) follows from [\(F.40\)](#). Regarding β_2 , we have

$$\beta_2 \leq \|\mathbf{E}_{\mathcal{I},\cdot}\| \|\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}\mathbf{H}_{\mathbf{V}}\| \leq C_g\sigma\sqrt{I+n_2+\log n}\|\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}\mathbf{H}_{\mathbf{V}}\|, \quad (\text{F.44})$$

where the last relation follows from [Lemma 19](#). It suffices to bound the spectral norm of $\mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_{\mathbf{V}}$, which appears in both upper bounds [\(F.43\)](#) and [\(F.44\)](#) for β_1 and β_2 .

We first observe that

$$\|\mathbf{U}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U} \mathbf{H}_{\mathbf{U}}\| = \|(\mathbf{U}^{(\mathcal{I})} \mathbf{U}^{(\mathcal{I})\top} - \mathbf{U} \mathbf{U}^\top) \mathbf{U}^*\| \leq \|\mathbf{U}^{(\mathcal{I})} \mathbf{U}^{(\mathcal{I})\top} - \mathbf{U} \mathbf{U}^\top\|, \quad \text{and} \quad (\text{F.45a})$$

$$\|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_{\mathbf{V}}\| = \|(\mathbf{V}^{(\mathcal{I})} \mathbf{V}^{(\mathcal{I})\top} - \mathbf{V} \mathbf{V}^\top) \mathbf{V}^*\| \leq \|\mathbf{V}^{(\mathcal{I})} \mathbf{V}^{(\mathcal{I})\top} - \mathbf{V} \mathbf{V}^\top\|. \quad (\text{F.45b})$$

The right hand side of the above two inequalities are the subspace distance between \mathbf{V} and $\mathbf{V}^{(\mathcal{I})}$, as well as \mathbf{U} and $\mathbf{U}^{(\mathcal{I})}$. This motivates us to apply Wedin's Θ Theorem to achieve

$$\begin{aligned} & \max \left\{ \|\mathbf{U}^{(\mathcal{I})} \mathbf{U}^{(\mathcal{I})\top} - \mathbf{U} \mathbf{U}^\top\|, \|\mathbf{V}^{(\mathcal{I})} \mathbf{V}^{(\mathcal{I})\top} - \mathbf{V} \mathbf{V}^\top\| \right\} \\ & \leq \frac{\sqrt{2} \max \left\{ \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M}) \mathbf{V}^{(\mathcal{I})}\|, \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M})^\top \mathbf{U}^{(\mathcal{I})}\| \right\}}{\sigma_r(\mathbf{M}^{(\mathcal{I})}) - \sigma_{r+1}(\mathbf{M}) - \|\mathbf{M}^{(\mathcal{I})} - \mathbf{M}\|} \\ & \stackrel{(i)}{\leq} \frac{\sqrt{2} \max \left\{ \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M}) \mathbf{V}^{(\mathcal{I})}\|, \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M})^\top \mathbf{U}^{(\mathcal{I})}\| \right\}}{\sigma_r(\mathbf{M}^*) - \|\mathcal{P}_{-\mathcal{I}, \cdot}(\mathbf{E})\| - \sigma_{r+1}(\mathbf{M}^*) - \|\mathbf{E}\| - \|\mathbf{E}_{\mathcal{I}, \cdot}\|} \\ & \stackrel{(ii)}{\leq} \frac{2\sqrt{2} \max \left\{ \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M}) \mathbf{V}^{(\mathcal{I})}\|, \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M})^\top \mathbf{U}^{(\mathcal{I})}\| \right\}}{\sigma_r^*}. \end{aligned} \quad (\text{F.46})$$

Here (i) follows from Weyl's inequality, while (ii) follows from the following consequence of [\(F.26\)](#):

$$\max \left\{ \|\mathcal{P}_{-\mathcal{I}, \cdot}(\mathbf{E})\|, \|\mathbf{E}_{\mathcal{I}, \cdot}\| \right\} \leq \|\mathbf{E}\| \leq C_g \sigma \sqrt{n} \leq c_n C_g \sigma_r^* \leq \frac{1}{2} \sigma_r^*,$$

where the last relation holds provided that c_n is sufficiently small. Then we shall bound the spectral norm of $(\mathbf{M}^{(\mathcal{I})} - \mathbf{M}) \mathbf{V}^{(\mathcal{I})}$ and $(\mathbf{M}^{(\mathcal{I})} - \mathbf{M})^\top \mathbf{U}^{(\mathcal{I})}$. The first one is upper bounded by

$$\begin{aligned} \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M}) \mathbf{V}^{(\mathcal{I})}\| &= \|\mathbf{E}_{\mathcal{I}, \cdot} \mathbf{V}^{(\mathcal{I})}\| \stackrel{(i)}{\leq} 2 \|\mathbf{E}_{\mathcal{I}, \cdot} \mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})}\| \leq 2 \|\mathbf{E}_{\mathcal{I}, \cdot} \mathbf{V}^*\| + 2 \|\mathbf{E}_{\mathcal{I}, \cdot} (\mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}^*)\| \\ &\stackrel{(ii)}{\leq} C_g \sigma \sqrt{I + r + \log n} + 2\beta_1, \end{aligned} \quad (\text{F.47})$$

and the second one is upper bounded by

$$\begin{aligned} \|(\mathbf{M}^{(\mathcal{I})} - \mathbf{M})^\top \mathbf{U}^{(\mathcal{I})}\| &= \|\mathbf{E}_{\mathcal{I}, \cdot}^\top \mathbf{U}_{\mathcal{I}, \cdot}^{(\mathcal{I})}\| \stackrel{(iii)}{\leq} 2 \|\mathbf{E}_{\mathcal{I}, \cdot}^\top \mathbf{U}_{\mathcal{I}, \cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})}\| \leq 2 \|\mathbf{E}_{\mathcal{I}, \cdot}^\top \mathbf{U}_{\mathcal{I}, \cdot}^*\| + 2 \|\mathbf{E}_{\mathcal{I}, \cdot}\| \|\mathbf{U}_{\mathcal{I}, \cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I}, \cdot}^*\| \\ &\stackrel{(iv)}{\leq} 2C_g \sigma \sqrt{n_2 + \log n} \|\mathbf{U}_{\mathcal{I}, \cdot}^*\| + C_g \sigma \sqrt{I + n_2 + \log n} \|\mathbf{U}_{\mathcal{I}, \cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I}, \cdot}^*\| \end{aligned} \quad (\text{F.48})$$

Here (i) and (iii) hold true when $\sigma_r(\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})}) \geq 1/2$ and $\sigma_r(\mathbf{H}_{\mathbf{U}}^{(\mathcal{I})}) \geq 1/2$, which can be verified following the same analysis for proving [Lemma 15](#) (where we showed that $\sigma_r(\mathbf{H}_{\mathbf{V}}) \geq 1/2$ and $\sigma_r(\mathbf{H}_{\mathbf{U}}) \geq 1/2$), and is omitted here for brevity; while (ii) and (iv) follow from [Lemma 19](#). Collecting together equations [\(F.45\)](#), [\(F.46\)](#), [\(F.47\)](#) and [\(F.48\)](#), we arrive at

$$\begin{aligned} & \max \left\{ \|\mathbf{U}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U} \mathbf{H}_{\mathbf{U}}\|, \|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_{\mathbf{V}}\| \right\} \\ & \leq \frac{20C_g}{\sigma_r^*} \left[\sigma \sqrt{I + r + \log n} + \sigma \sqrt{n_2} (\|\mathbf{U}_{\mathcal{I}, \cdot}^*\| + \|\mathbf{U}_{\mathcal{I}, \cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I}, \cdot}^*\|) \right] + \frac{6\beta_1}{\sigma_r^*}. \end{aligned} \quad (\text{F.49})$$

Here we use the facts that $\|\mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 1$ and $\|\mathbf{U}_{\mathcal{I},\cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 3$. We want to replace the spectral norm of $\mathbf{U}_{\mathcal{I},\cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I},\cdot}^*$ in (F.49) with that of $\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*$. This can be done by a self-bounding trick: by the triangle inequality and (F.49), we have

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I},\cdot}^*\| &\leq \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U} \mathbf{H}_{\mathbf{U}}\| \\ &\leq \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\| + \frac{20C_g}{\sigma_r^*} [\sigma\sqrt{I+r+\log n} + \sigma\sqrt{n_2}(\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I},\cdot}^*\|)] + \frac{6\beta_1}{\sigma_r^*}. \end{aligned}$$

Notice that both sides of the above inequality include $\|\mathbf{U}_{\mathcal{I},\cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I},\cdot}^*\|$. Since $\sigma\sqrt{n_2} \leq c_n\sigma_r^*$ for some sufficiently small $c_n > 0$, we can rearrange terms to reach

$$\|\mathbf{U}_{\mathcal{I},\cdot}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 2\|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\| + \frac{30C_g}{\sigma_r^*} (\sigma\sqrt{I+r+\log n} + \sigma\sqrt{n_2}\|\mathbf{U}_{\mathcal{I},\cdot}^*\|) + \frac{9\beta_1}{\sigma_r^*}. \quad (\text{F.50})$$

Substituting (F.50) into (F.49) gives

$$\begin{aligned} &\max \left\{ \|\mathbf{U}^{(\mathcal{I})} \mathbf{H}_{\mathbf{U}}^{(\mathcal{I})} - \mathbf{U} \mathbf{H}_{\mathbf{U}}\|, \|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_{\mathbf{V}}\| \right\} \\ &\leq \frac{40C_g}{\sigma_r^*} [\sigma\sqrt{I+r+\log n} + \sigma\sqrt{n_2}(\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\|)] + \frac{9\beta_1}{\sigma_r^*} \end{aligned} \quad (\text{F.51})$$

provided that $\sigma\sqrt{n_2} \leq c_n\sigma_r^*$ holds for some sufficiently small $c_n > 0$.

Finally, we can substitute (F.51) back to (F.43) to reach

$$\begin{aligned} \beta_1 &\leq 2C_g^2 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} + C_g\sigma\sqrt{I+r+\log n} \|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_{\mathbf{V}}\| \\ &\leq 250C_g^2 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} + 9C_g \frac{\sigma\sqrt{I+r+\log n}}{\sigma_r^*} \beta_1, \end{aligned}$$

where we have used the facts that $\|\mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 1$ and $\|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 3$. Notice that β_1 appears in both side of the inequality. Given that $\sigma\sqrt{I+r+\log n} \leq 2\sigma\sqrt{n} \leq 2c_n\sigma_r^*$ for some sufficiently small $c_n > 0$, we can rearrange terms to deduce that

$$\beta_1 \leq 300C_g^2 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)}. \quad (\text{F.52})$$

Then we substitute (F.51) back to (F.44) to get

$$\begin{aligned} \beta_2 &\leq C_g\sigma\sqrt{I+n_2+\log n} \|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_{\mathbf{V}}\| \\ &\leq C_g\sigma\sqrt{I+n_2+\log n} \left[\frac{40C_g}{\sigma_r^*} [\sigma\sqrt{I+r+\log n} + \sigma\sqrt{n_2}(\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\|)] + \frac{9\beta_1}{\sigma_r^*} \right] \\ &\leq 50C_g^2\sigma\sqrt{I+n_2+\log n} \left[\frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} + \frac{\sigma}{\sigma_r^*} \sqrt{n_2} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\|) \right]. \end{aligned} \quad (\text{F.53})$$

Here the last step follows from (F.52) and it holds true as long as $c_n > 0$ is sufficiently small. Taking the bound (F.52) on β_1 and the bound (F.53) on β_2 collectively, we have

$$\beta \leq \beta_1 + \beta_2 \leq 400C_g^2 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} + 100C_g^2 \frac{\sigma^2 n_2}{\sigma_r^*} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_{\mathbf{U}} - \mathbf{U}_{\mathcal{I},\cdot}^*\|). \quad (\text{F.54})$$

Step 4: putting everything together. Combining the bounds in the preceding steps, we know that with probability exceeding $1 - O(n^{-10})$

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\| &\leq \alpha + \beta \\ &\leq 3C_g^2 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 400C_g^2 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} + 100C_g^2 \frac{\sigma^2 n_2}{\sigma_r^*} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|) \\ &\leq c_7 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(I+r+\log n)} + c_7 \frac{\sigma^2 n_2}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|, \end{aligned}$$

where $c_7 = 400C_g^2$. Following the same analysis, we can also show that

$$\|\mathbf{V}_{\mathcal{J},\cdot} \boldsymbol{\Sigma} \mathbf{H}_U - \mathbf{M}_{\mathcal{J},\cdot}^\top \mathbf{U}^*\| \leq c_7 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{V}_{\mathcal{J},\cdot}^*\| + c_7 \frac{\sigma^2}{\sigma_r^*} \sqrt{n(J+r+\log n)} + c_7 \frac{\sigma^2 n_1}{\sigma_r^*} \|\mathbf{V}_{\mathcal{J},\cdot} \mathbf{H}_V - \mathbf{V}_{\mathcal{J},\cdot}^*\|,$$

where $J = |\mathcal{J}|$. These complete the proof.

F.2.4 Proof of Lemma 18

We first use the triangle inequality to achieve

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| &\leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U \boldsymbol{\Sigma}^* - \mathbf{U}_{\mathcal{I},\cdot}^* \boldsymbol{\Sigma}^*\| \\ &\leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{U}_{\mathcal{I},\cdot}^* \boldsymbol{\Sigma}^*\| + \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U \boldsymbol{\Sigma}^*\| \\ &\leq \underbrace{\frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} \boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{M}_{\mathcal{I},\cdot} \mathbf{V}^*\|}_{=: \alpha_1} + \underbrace{\frac{1}{\sigma_r^*} \|\mathbf{E}_{\mathcal{I},\cdot} \mathbf{V}^*\|}_{=: \alpha_2} + \underbrace{\frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} (\boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{H}_U \boldsymbol{\Sigma}^*)\|}_{=: \alpha_3}, \end{aligned}$$

where the last line follows from $\mathbf{M} \mathbf{V}^* = \mathbf{U}^* \boldsymbol{\Sigma}^* + \mathbf{E} \mathbf{V}^*$. Thus, the problem reduces to bounding α_1 , α_2 and α_3 . Applying Lemma 17 yields

$$\alpha_1 \leq c_7 \frac{\sigma^2 n}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_7 \frac{\sigma^2}{\sigma_r^{*2}} \sqrt{n(I+r+\log n)} + c_7 \frac{\sigma^2 n_2}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|. \quad (\text{F.55})$$

Regarding α_2 , we can invoke Lemma 19 to show that, with probability exceeding $1 - O(n^{-10})$,

$$\alpha_2 \leq C_g \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n}. \quad (\text{F.56})$$

We are now left with bounding α_3 , and it is seen from Lemma 16 that

$$\begin{aligned} \alpha_3 &\leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}\| \|\boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{H}_U \boldsymbol{\Sigma}^*\| \leq \frac{2}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U\| \|\boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{H}_U \boldsymbol{\Sigma}^*\| \\ &\leq \frac{2}{\sigma_r^*} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|) \|\boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{H}_U \boldsymbol{\Sigma}^*\|, \end{aligned} \quad (\text{F.57})$$

where we used $\sigma_r(\mathbf{H}_U) \geq 1/2$ (see Lemma 15). Recall the full SVD of \mathbf{M} in (F.25), which gives

$$\begin{aligned} \boldsymbol{\Sigma} \mathbf{H}_V - \mathbf{H}_U \boldsymbol{\Sigma}^* &= \boldsymbol{\Sigma} \mathbf{V}^\top \mathbf{V}^* - \mathbf{U}^\top \mathbf{U}^* \boldsymbol{\Sigma}^* = \mathbf{U}^\top (\mathbf{U} \boldsymbol{\Sigma} \mathbf{V}^\top - \mathbf{U}^* \boldsymbol{\Sigma}^* \mathbf{V}^{*\top}) \mathbf{V}^* \\ &= \mathbf{U}^\top (\mathbf{E} - \mathbf{U}_\perp \boldsymbol{\Sigma}_\perp \mathbf{V}_\perp^\top) \mathbf{V}^* = \mathbf{U}^\top \mathbf{E} \mathbf{V}^*. \end{aligned}$$

This allows us to bound

$$\begin{aligned}
\|\Sigma \mathbf{H}_V - \mathbf{H}_U \Sigma^*\| &\stackrel{(i)}{\leq} 2\|(\mathbf{U} \mathbf{H}_U)^\top \mathbf{E} \mathbf{V}^*\| \leq 2\|(\mathbf{U} \mathbf{H}_U - \mathbf{U}^*)^\top \mathbf{E} \mathbf{V}^*\| + 2\|\mathbf{U}^{*\top} \mathbf{E} \mathbf{V}^*\| \\
&\leq 2\|\mathbf{U} \mathbf{H}_U - \mathbf{U}^*\| \|\mathbf{E} \mathbf{V}^*\| + 2\|\mathbf{U}^{*\top} \mathbf{E} \mathbf{V}^*\| \\
&\stackrel{(ii)}{\leq} 4C_g \frac{\sigma}{\sigma_r^*} \sqrt{n} \cdot C_g \sigma \sqrt{n_1 + r + \log n} + 2C_g \sigma \sqrt{r + \log n} \\
&\stackrel{(iii)}{\leq} 2C_g \sigma \sqrt{r + \log n} + 7C_g^2 \frac{\sigma^2 n}{\sigma_r^*}. \tag{F.58}
\end{aligned}$$

Here (i) holds since $\sigma_r(\mathbf{H}_U) \geq 1/2$ (see [Lemma 15](#)), (ii) follows from [Lemma 19](#) as well as a direct consequence of [\(F.28\)](#):

$$\|\mathbf{U} \mathbf{H}_U - \mathbf{U}^*\| = \|\mathbf{U} \mathbf{U}^\top \mathbf{U}^* - \mathbf{U}^*\| = \|(\mathbf{U} \mathbf{U}^\top - \mathbf{U}^* \mathbf{U}^{*\top}) \mathbf{U}^*\| \leq \|\mathbf{U} \mathbf{U}^\top - \mathbf{U}^* \mathbf{U}^{*\top}\| \leq 2C_g \frac{\sigma}{\sigma_r^*} \sqrt{n};$$

while (iii) holds with the proviso that $\sigma \sqrt{n} \lesssim \sigma_r^*$. Substituting [\(F.58\)](#) into [\(F.57\)](#) gives

$$\alpha_3 \leq \frac{2}{\sigma_r^*} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|) (2C_g \sigma \sqrt{r + \log n} + 7C_g^2 \frac{\sigma^2 n}{\sigma_r^*}). \tag{F.59}$$

Taking the bounds [\(F.55\)](#) on α_1 , [\(F.56\)](#) on α_2 and [\(F.59\)](#) on α_3 collectively, we achieve

$$\begin{aligned}
\|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| &\leq \alpha_1 + \alpha_2 + \alpha_3 \\
&\leq (17C_g + c_7 c_n) \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n} + (c_7 + 14C_g^2) \frac{\sigma^2 n}{\sigma_r^{*2}} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|).
\end{aligned}$$

Here we have used the facts that $\|\mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 1$ and $\|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 3$. Notice that the term $\|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|$ appears on both side of the above inequality. Since we assume that $\sigma \sqrt{n} \leq c_n \sigma_r^*$ for some sufficiently small $c_n > 0$, we can use the self-bounding trick to obtain

$$\|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 20C_g \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n} + (2c_7 + 20C_g^2) \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\|$$

as claimed, if we take $c_8 = 2c_7 + 20C_g^2$. Following the same analysis, we can also show that

$$\|\mathbf{V}_{\mathcal{J},\cdot} \mathbf{H}_V - \mathbf{V}_{\mathcal{J},\cdot}^*\| \leq c_8 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{V}_{\mathcal{J},\cdot}^*\| + c_8 \frac{\sigma}{\sigma_r^*} \sqrt{J + r + \log n}.$$

F.2.5 Proof of equations in [Appendix F.1.3](#)

This section provides the proof of equations [\(F.20\)](#) and [\(F.23\)](#) in [Appendix F.1.3](#).

Proof of [\(F.20\)](#). Applying the triangle inequality yields

$$\begin{aligned}
\|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{R}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| &\leq \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot} (\mathbf{H}_U - \mathbf{R}_U)\| \leq \|\mathbf{U}_{\mathcal{I},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot}\| \|\mathbf{H}_U - \mathbf{R}_U\| \\
&\stackrel{(i)}{\leq} c_8 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n} + (3\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n}) \cdot 16C_g^2 \frac{\sigma^2 n}{\sigma_r^{*2}} \\
&\stackrel{(ii)}{\leq} (c_8 + 48C_g^2) \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n}.
\end{aligned}$$

Here (i) follows from [Lemma 18](#), [\(F.12\)](#) and [\(F.35\)](#), while (ii) holds true when $\sigma \sqrt{n} \leq c_n \sigma_r^*$ for some sufficiently small $c_n > 0$. We have proved [\(F.20a\)](#), and [\(F.20b\)](#) can be verified analogously.

Proof of (F.23). A weaker version of this result was established in Lemma 17. Here we refine several steps in its proof using sharper perturbation bounds (F.20). We first improve (F.38) to

$$\begin{aligned} \|\mathbf{U}^{\star\top} \mathbf{U}_\perp \boldsymbol{\Sigma}_\perp \mathbf{V}_\perp^\top \mathbf{V}^\star\| &\leq \|\mathbf{U}^{\star\top} \mathbf{U}_\perp\| \|\boldsymbol{\Sigma}_\perp\| \|\mathbf{V}^{\star\top} \mathbf{V}_\perp\| \stackrel{(i)}{\leq} \|\mathbf{U} \mathbf{R}_U - \mathbf{U}^\star\| \|\boldsymbol{\Sigma}_\perp\| \|\mathbf{V} \mathbf{R}_V - \mathbf{V}^\star\| \\ &\stackrel{(ii)}{\leq} \left(\frac{\sigma^2 n}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^\star} \sqrt{n_1 + \log n} \right) \left(\frac{\sigma^2 n}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^\star} \sqrt{n_2 + \log n} \right) \sigma \sqrt{n}, \end{aligned} \quad (\text{F.60})$$

Here (i) follows from the relation between subspace distances (see e.g. [Che+21a, Section 2.2.2]), while (ii) utilizes the refined bound (F.20). We can improve (F.39) to

$$\begin{aligned} \|\mathbf{U}^{\star\top} \mathbf{E}(\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^\star\| &\leq \|\mathbf{U}^{\star\top} \mathbf{E}\| \|\mathbf{V} \mathbf{H}_V - \mathbf{V}^\star\| \\ &\leq C_g \sigma \sqrt{n_2 + \log n} \cdot c'_5 \left(\frac{\sigma^2 n}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^\star} \sqrt{n_2 + \log n} \right), \end{aligned} \quad (\text{F.61})$$

where we use the a direct consequence of (F.20a):

$$\|\mathbf{V} \mathbf{H}_V - \mathbf{V}^\star\| \stackrel{(a)}{\leq} \|\mathbf{V} \mathbf{V}^\top - \mathbf{V}^\star \mathbf{V}^{\star\top}\| \stackrel{(b)}{\leq} \|\mathbf{V} \mathbf{R}_V - \mathbf{V}^\star\| \stackrel{(c)}{\leq} c'_5 \frac{\sigma^2 n}{\sigma_r^{\star 2}} + c'_5 \frac{\sigma}{\sigma_r^\star} \sqrt{n_2 + \log n}. \quad (\text{F.62})$$

Here (a) utilizes (F.40), (b) follows again from the relation between subspace distances, while (c) uses the refined bound in (F.21b). Putting (F.60) and (F.61) together, we can improve (F.41) to

$$\begin{aligned} \|\boldsymbol{\Sigma}^\star \mathbf{V}^{\star\top} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^\star)\| &\leq \|\mathbf{U}^{\star\top} \mathbf{U}_\perp \boldsymbol{\Sigma}_\perp \mathbf{V}_\perp^\top \mathbf{V}^\star\| + \|\mathbf{U}^{\star\top} \mathbf{E}(\mathbf{V} \mathbf{V}^\top - \mathbf{I}_{n_2}) \mathbf{V}^\star\| \\ &\leq 2C_g c'_5 \left(\frac{\sigma^2 n_1}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^\star} \sqrt{n_2 + \log n} \right)^2 \sigma_r^\star, \end{aligned} \quad (\text{F.63})$$

provided that $\sigma \sqrt{n} \leq c_n \sigma_r^\star$ for some sufficiently small $c_n > 0$. This allows us to improve our original bound on α in (F.42) to

$$\alpha \leq 2C_g c'_5 \left(\frac{\sigma^2 n_1}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^\star} \sqrt{n_2 + \log n} \right)^2 \sigma_r^\star \|\mathbf{U}_{\mathcal{L}}^\star\|. \quad (\text{F.64})$$

We also need to refine our bound on β , where we recall that $\beta \leq \beta_1 + \beta_2$ with

$$\beta_1 = \|\mathbf{E}_{\mathcal{L},\cdot}(\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V}^\star)\| \quad \text{and} \quad \beta_2 = \|\mathbf{E}_{\mathcal{L},\cdot}(\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_V)\|.$$

We improve our bound on β_1 to

$$\begin{aligned} \beta_1 &\stackrel{(i)}{\leq} C_g \sigma \sqrt{I + r + \log n} \|\mathbf{V} \mathbf{H}_V - \mathbf{V}^\star\| + C_g \sigma \sqrt{I + r + \log n} \|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_V\| \\ &\stackrel{(ii)}{\leq} C_g c'_5 \sigma \sqrt{I + r + \log n} \left(\frac{\sigma^2 n}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^\star} \sqrt{n_2 + \log n} \right) + C_g \frac{\sigma}{\sigma_r^\star} \sqrt{I + r + \log n} \cdot \beta_1 \\ &\quad + 40C_g^2 \sigma \sqrt{I + r + \log n} \left[\frac{\sigma}{\sigma_r^\star} \sqrt{I + r + \log n} + \frac{\sigma}{\sigma_r^\star} \sqrt{n_2} (\|\mathbf{U}_{\mathcal{L},\cdot}^\star\| + \|\mathbf{U}_{\mathcal{L},\cdot} \mathbf{H}_U - \mathbf{U}_{\mathcal{L},\cdot}^\star\|) \right] \\ &\stackrel{(iii)}{\leq} (100C_g^2 + 3C_g c'_5) \sigma \sqrt{I + r + \log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{\star 2}} + \frac{\sigma}{\sigma_r^\star} \sqrt{I + n_2 + \log n} \right). \end{aligned} \quad (\text{F.65})$$

Here (i) uses our original derivation in (F.43), (ii) utilizes (F.62) and (F.51), and (iii) follows $\|\mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 1$, $\|\mathbf{U}_{\mathcal{I},\cdot}\mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 2$, as well as the self-bounding trick that appeared several times in the proof of Lemma 17, which holds provided that $\sigma\sqrt{I+r+\log n} \leq 2\sigma\sqrt{n} \leq 2c_n\sigma_r^*$ for some sufficiently small $c_n > 0$. Then we can improve (F.51) to

$$\begin{aligned}
\|\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}\mathbf{H}_{\mathbf{V}}\| &\stackrel{(a)}{\leq} \frac{40C_g}{\sigma_r^*} [\sigma\sqrt{I+r+\log n} + \sigma\sqrt{n_2}(\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + \|\mathbf{U}_{\mathcal{I},\cdot}\mathbf{H}_U - \mathbf{U}_{\mathcal{I},\cdot}^*\|)] + \frac{9\beta_1}{\sigma_r^*} \\
&\stackrel{(b)}{\leq} 40C_g \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} + 40C_g \frac{\sigma}{\sigma_r^*} \sqrt{n_2} (\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_8 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n}) \\
&\quad + (900C_g^2 + 27C_g c'_5) \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{I+n_2+\log n} \right) \\
&\stackrel{(c)}{\leq} 50C_g \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} + 50C_g \frac{\sigma}{\sigma_r^*} \sqrt{n_2} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \tag{F.66}
\end{aligned}$$

Here step (a) utilizes (F.51), step (b) follows from (F.65) and Lemma 18, while step (c) holds provided that $\sigma\sqrt{I+n_2+\log n} \leq \sigma\sqrt{n} \leq c_n\sigma_r^*$ for some sufficiently small $c_n > 0$. This allows us to improve the bound (F.53) on β_2 and obtain

$$\begin{aligned}
\beta_2 &\leq C_g \sigma \sqrt{I+n_2+\log n} \|\mathbf{V}^{(\mathcal{I})}\mathbf{H}_{\mathbf{V}}^{(\mathcal{I})} - \mathbf{V}\mathbf{H}_{\mathbf{V}}\| \\
&\leq 50C_g^2 \sigma \sqrt{I+n_2+\log n} \left(\frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} + \frac{\sigma}{\sigma_r^*} \sqrt{n_2} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \right).
\end{aligned}$$

Putting the above refined bounds on β_1 and β_2 together yields

$$\beta \leq (150C_g^2 + 3C_g c'_5) \sigma \sqrt{I+r+\log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{I+n_2+\log n} \right) + 50C_g^2 \frac{\sigma^2 n_2}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\|, \tag{F.67}$$

where we use the fact that $\|\mathbf{U}_{\mathcal{I},\cdot}^*\| \leq 1$. Finally, taking the refined bound (F.64) on α and the refined bound (F.67) on β collectively yields

$$\begin{aligned}
\|\mathbf{U}_{\mathcal{I},\cdot}\boldsymbol{\Sigma}\mathbf{H}_{\mathbf{V}} - \mathbf{M}_{\mathcal{I},\cdot}\mathbf{V}^*\| &\leq \alpha + \beta \\
&\leq c'_7 \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{n_2+\log n} \right)^2 \sigma_r^* \|\mathbf{U}_{\mathcal{I},\cdot}^*\| + c'_7 \sigma \sqrt{I+r+\log n} \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{I+n_2+\log n} \right).
\end{aligned}$$

where $c'_7 = 150C_g^2 + 3C_g c'_5$. We have proved (F.23a), and the other claimed bound (F.23b) can be proved analogously.

F.3 Proof of corollaries for matrix denoising

This section provide the proof for Corollaries 1 and 2.

F.3.1 Proof of Corollary 1

To begin with, we decompose the error of interest as follows

$$\begin{aligned}
\mathbf{U}\boldsymbol{\Sigma}\mathbf{V}^\top - \mathbf{M}^* &= \mathbf{U}\boldsymbol{\Sigma}\mathbf{V}^\top - \mathbf{U}^*\boldsymbol{\Sigma}^*\mathbf{V}^{*\top} = \mathbf{U}\mathbf{R}_U\mathbf{R}_U^\top\boldsymbol{\Sigma}\mathbf{V}^\top - \mathbf{U}^*\boldsymbol{\Sigma}^*\mathbf{V}^{*\top} \\
&= (\mathbf{U}^* + \mathbf{E}\mathbf{V}^*(\boldsymbol{\Sigma}^*)^{-1} + \boldsymbol{\Psi}_U)(\mathbf{M}^\top\mathbf{U}^* + \boldsymbol{\Delta}_V)^\top - \mathbf{U}^*\boldsymbol{\Sigma}^*\mathbf{V}^{*\top} \\
&= \mathbf{U}^*\mathbf{U}^{*\top}\mathbf{E} + \mathbf{E}\mathbf{V}^*\mathbf{V}^{*\top} + \underbrace{\boldsymbol{\Psi}_U\boldsymbol{\Sigma}^*\mathbf{V}^{*\top} + (\mathbf{U}\mathbf{R}_U - \mathbf{U}^*)\mathbf{U}^{*\top}\mathbf{E} + \mathbf{U}\mathbf{R}_U\boldsymbol{\Delta}_V^\top}_{=: \boldsymbol{\Phi}}.
\end{aligned}$$

For any $\mathcal{I} \subseteq [n_1]$ and $\mathcal{J} \subseteq [n_2]$, with probability exceeding $1 - O(n^{-10})$ we have

$$\begin{aligned}
\|\Phi_{\mathcal{I},\mathcal{J}}\| &\leq \|(\Psi_U)_{\mathcal{I},\Sigma^*}\| \|V_{\mathcal{J},\cdot}^*\| + \|U_{\mathcal{I},\cdot} R_U - U_{\mathcal{I},\cdot}^*\| \|U^{*\top} E_{\cdot,\mathcal{J}}\| + \|U_{\mathcal{I},\cdot}\| \|(\Delta_V)_{\mathcal{J},\cdot}\| \\
&\stackrel{(i)}{\leq} c'_1 \sigma \sqrt{I+r+\log n} \left(\frac{\sigma}{\sigma_r^*} \sqrt{I+n_2+\log n} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right) \|V_{\mathcal{J},\cdot}^*\| + c'_1 \left(\frac{\sigma^2 n}{\sigma_r^*} + \sigma \sqrt{r+\log n} \right) \|U_{\mathcal{I},\cdot}^*\| \|V_{\mathcal{J},\cdot}^*\| \\
&\quad + c'_5 \left(\frac{\sigma^2 n}{\sigma_r^{*2}} \|U_{\mathcal{I},\cdot}^*\| + \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} \right) \cdot C_g \sigma \sqrt{J+r+\log n} \\
&\quad + \left(3 \|U_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} \right) \cdot c_2 \left[\sigma \sqrt{J+r+\log n} \left(\frac{\sigma}{\sigma_r^*} \sqrt{J+n_1+\log n} + \frac{\sigma^2 n}{\sigma_r^{*2}} \right) + \frac{\sigma^2 n}{\sigma_r^*} \|V_{\mathcal{J},\cdot}^*\| \right] \\
&\stackrel{(ii)}{\leq} c'_3 \frac{\sigma^2}{\sigma_r^*} \sqrt{(I+r+\log n)(J+r+\log n)} + c'_3 \sigma \sqrt{J+r+\log n} \left(\frac{\sigma}{\sigma_r^*} \sqrt{J+n_1} + \frac{\sigma^2 n_2}{\sigma_r^{*2}} \right) \|U_{\mathcal{I},\cdot}^*\| \\
&\quad + c'_3 \sigma \sqrt{I+r+\log n} \left(\frac{\sigma}{\sigma_r^*} \sqrt{I+n_2} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right) \|V_{\mathcal{J},\cdot}^*\| + c'_3 \left(\frac{\sigma^2 n}{\sigma_r^*} + \sigma \sqrt{r+\log n} \right) \|U_{\mathcal{I},\cdot}^*\| \|V_{\mathcal{J},\cdot}^*\|.
\end{aligned}$$

Here (ii) holds by taking $c'_3 = 2c'_1 + 2c'_5 + 6c_2 + 4c_8 c_2$, and (i) follows from the equations (F.20), (F.12), Lemma 19, and Proposition 3, as well as

$$\begin{aligned}
\|(\Psi_U)_{\mathcal{I},\Sigma^*}\| &\stackrel{(iii)}{=} \|(\Delta_U)_{\mathcal{I},\cdot} + U_{\mathcal{I},\cdot} R_U \Delta_\Sigma\| \leq \|(\Delta_U)_{\mathcal{I},\cdot}\| + \|U_{\mathcal{I},\cdot}\| \|\Delta_\Sigma\| \\
&\stackrel{(iv)}{\leq} c_2 \sigma \sqrt{I+r+\log n} \left[\frac{\sigma}{\sigma_r^*} \sqrt{I+n_2+\log n} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right] + c_2 \frac{\sigma^2 n}{\sigma_r^*} \|U_{\mathcal{I},\cdot}^*\| \\
&\quad + \left(3 \|U_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I+r+\log n} \right) \cdot c_6 \left[\frac{\sigma^2 n}{\sigma_r^*} + \sigma \sqrt{r+\log n} \right] \\
&\stackrel{(v)}{\leq} c'_1 \sigma \sqrt{I+r+\log n} \left[\frac{\sigma}{\sigma_r^*} \sqrt{I+n_2+\log n} + \frac{\sigma^2 n_1}{\sigma_r^{*2}} \right] + c'_1 \left(\frac{\sigma^2 n}{\sigma_r^*} + \sigma \sqrt{r+\log n} \right) \|U_{\mathcal{I},\cdot}^*\|,
\end{aligned}$$

where (iii) follows from (F.17), (iv) follows from Proposition 3, equations (F.16) and (F.12), while (v) holds by taking $c'_1 = c_2 + 3c_6 + 4c_6 c_8$. Next, applying Lemma 19 yields

$$\|U_{\mathcal{I},\cdot}^* U^{*\top} E_{\cdot,\mathcal{J}} + E_{\mathcal{I},\cdot} V^* V_{\mathcal{J},\cdot}^{*\top}\| \leq C_g \sigma \sqrt{J+r+\log n} \|U_{\mathcal{I},\cdot}^*\| + C_g \sigma \sqrt{I+r+\log n} \|V_{\mathcal{J},\cdot}^*\|.$$

Combining the above bounds with the triangle inequality yields

$$\begin{aligned}
\|(U\Sigma V^\top - M^*)_{\cdot,\mathcal{J}}\| &\leq \|U_{\mathcal{I},\cdot}^* U^{*\top} E_{\cdot,\mathcal{J}} + E_{\mathcal{I},\cdot} V^* V_{\mathcal{J},\cdot}^{*\top}\| + \|\Phi_{\mathcal{I},\mathcal{J}}\| \\
&\leq c_3 \sigma \sqrt{J+r+\log n} \|U_{\mathcal{I},\cdot}^*\| + c_3 \sigma \sqrt{I+r+\log n} \|V_{\mathcal{J},\cdot}^*\| \\
&\quad + c_3 \frac{\sigma^2 n}{\sigma_r^*} \|U_{\mathcal{I},\cdot}^*\| \|V_{\mathcal{J},\cdot}^*\| + c_3 \frac{\sigma^2}{\sigma_r^*} \sqrt{(I+r+\log n)(J+r+\log n)},
\end{aligned}$$

where we take $c_3 = 2C_g + 2c'_3$. Here we use the fact that $\|U_{\mathcal{I},\cdot}^*\| \leq 1$ and the assumption that $\sigma\sqrt{n} \leq c_n \sigma_r^*$ holds for some sufficiently small $c_n > 0$.

F.3.2 Proof of Corollary 2

In view of Proposition 2, we can use the triangle inequality to obtain

$$\|U_{\mathcal{I},\cdot}^{*\top} (U_{\mathcal{I},\cdot} R_U - U_{\mathcal{I},\cdot}^*)\| \leq \underbrace{\|U_{\mathcal{I},\cdot}^{*\top} E_{\mathcal{I},\cdot} V^* (\Sigma^*)^{-1}\|}_{=:\alpha} + \underbrace{\|U_{\mathcal{I},\cdot}^{*\top} (\Psi_U)_{\mathcal{I},\cdot}\|}_{=:\beta}.$$

Thus, we have reduced the problem to bounding α and β .

We begin by bounding α . Applying [Lemma 19](#) yields

$$\alpha \leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{E}_{\mathcal{I},\cdot} \mathbf{V}^*\| \leq C_g \frac{\sigma}{\sigma_r^*} \sqrt{r + \log n} \|\mathbf{U}_{\mathcal{I},\cdot}^*\|.$$

Recalling the definition (19) of Ψ_U , we can upper bound β by

$$\begin{aligned} \beta &= \|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} (\Delta_U)_{\mathcal{I},\cdot} (\Sigma^*)^{-1} + \mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{U}_{\mathcal{I},\cdot} \mathbf{R}_U \Delta_\Sigma (\Sigma^*)^{-1}\| \\ &\leq \underbrace{\|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{U}_{\mathcal{I},\cdot}^* \Sigma^* \mathbf{V}^{*\top} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*) (\Sigma^*)^{-1}\|}_{=:\beta_1} + \underbrace{\|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{E}_{\mathcal{I},\cdot} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*) (\Sigma^*)^{-1}\|}_{=:\beta_2} \\ &\quad + \underbrace{\|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{U}_{\mathcal{I},\cdot} \mathbf{R}_U \Delta_\Sigma (\Sigma^*)^{-1}\|}_{=:\beta_3}. \end{aligned}$$

We now need to bound the three terms β_1 , β_2 and β_3 . Beginning with β_1 , applying the relation (F.63) yields

$$\beta_1 \leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\|^2 \|\Sigma^* \mathbf{V}^{*\top} (\mathbf{V} \mathbf{H}_V - \mathbf{V}^*)\| \leq 2C_g c'_5 \left(\frac{\sigma^2 n_1}{\sigma_r^{*2}} + \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n} \right)^2 \|\mathbf{U}_{\mathcal{I},\cdot}^*\|^2.$$

Turning to β_2 , applying the triangle inequality yields

$$\beta_2 \leq \underbrace{\|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{E}_{\mathcal{I},\cdot} (\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V}^*) (\Sigma^*)^{-1}\|}_{=:\beta_{2,1}} + \underbrace{\|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{E}_{\mathcal{I},\cdot} (\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_V) (\Sigma^*)^{-1}\|}_{=:\beta_{2,2}}.$$

The definition of $\mathbf{V}^{(\mathcal{I})}$ and $\mathbf{H}_V^{(\mathcal{I})}$ can be found in [Appendix F.2.3](#). Since $\mathbf{E}_{\mathcal{I},\cdot}$ and $\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V}^*$ are independent, we can invoke [Lemma 19](#) to obtain

$$\beta_{2,1} \leq C_g \frac{\sigma}{\sigma_r^*} \sqrt{r + \log n} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V}^*\| \leq C_g c'_5 \frac{\sigma}{\sigma_r^*} \sqrt{r + \log n} \left(\frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n} + \frac{\sigma^2 n}{\sigma_r^{*2}} \right) \|\mathbf{U}_{\mathcal{I},\cdot}^*\|,$$

where we have used the fact that

$$\|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V}^*\| \leq c'_5 \frac{\sigma^2 n}{\sigma_r^{*2}} + c'_5 \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n}.$$

(This bound can be established by an argument similar to that used in proving the bound (F.62)).

In addition, applying [Lemma 19](#) and equation (F.66) yields

$$\begin{aligned} \beta_{2,2} &\leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{E}_{\mathcal{I},\cdot}\| \|\mathbf{V}^{(\mathcal{I})} \mathbf{H}_V^{(\mathcal{I})} - \mathbf{V} \mathbf{H}_V\| \\ &\leq 50C_g^2 \frac{\sigma}{\sigma_r^*} \sqrt{n_2 + \log n} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \left(\frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n} + \frac{\sigma}{\sigma_r^*} \sqrt{n_2} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \right). \end{aligned}$$

Lastly, turning to β_3 , it can be upper bounded as

$$\begin{aligned} \beta_3 &\leq \|\mathbf{U}_{\mathcal{I},\cdot}^{*\top} \mathbf{U}_{\mathcal{I},\cdot} \mathbf{R}_U \Delta_\Sigma (\Sigma^*)^{-1}\| \leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \|\mathbf{U}_{\mathcal{I},\cdot}\| \|\Delta_\Sigma\| \\ &\leq \frac{1}{\sigma_r^*} \|\mathbf{U}_{\mathcal{I},\cdot}^*\| \left(3\|\mathbf{U}_{\mathcal{I},\cdot}^*\| + 2c_8 \frac{\sigma}{\sigma_r^*} \sqrt{I + r + \log n} \right) \cdot c_6 \left(\frac{\sigma^2 n}{\sigma_r^*} + \sigma \sqrt{r + \log n} \right), \end{aligned}$$

where the last relation follows from the relations (F.12) and (F.16).

By combining the above bounds, we find that

$$\begin{aligned} \|\mathbf{U}_{\mathcal{I},\cdot}^{\star\top}(\mathbf{U}_{\mathcal{I},\cdot}\mathbf{R}\mathbf{U} - \mathbf{U}_{\mathcal{I},\cdot}^{\star})\| &\leq \alpha + \beta \leq \alpha + \beta_1 + \beta_2 + \beta_3 \leq \alpha + \beta_1 + \beta_{2,1} + \beta_{2,2} + \beta_3 \\ &\leq c_4 \left[\frac{\sigma}{\sigma_r^*} \sqrt{r + \log n} + \frac{\sigma^2}{\sigma_r^{*2}} \sqrt{n_2(n_2 + I)} + \frac{\sigma^3 n_1}{\sigma_r^{*3}} \sqrt{I} \right] \|\mathbf{U}_{\mathcal{I},\cdot}^{\star}\| + c_4 \frac{\sigma^2 n}{\sigma_r^{*2}} \|\mathbf{U}_{\mathcal{I},\cdot}^{\star}\|^2, \end{aligned}$$

where we take $c_4 = 5C_g c'_5 + 100C_g^2 + 5c_6 + 5c_8 c_6$. Here we use the condition that $\sigma\sqrt{n} \leq c_n \sigma_r^*$ for some sufficiently small constant $c_n > 0$.

G Technical lemma

This section provides some technical lemmas that are useful in the paper.

Lemma 19. *Let $\mathbf{E} \in \mathbb{R}^{n_1 \times n_2}$ be a random matrix with independent entries, and $\|E_{i,j}\|_{\psi_2} \leq \sigma$. Then there exists some sufficiently large constant $C_g > 0$ such that*

$$\|\mathbf{E}\| \leq C_g \sigma \sqrt{n} \quad \text{where } n = \max\{n_1, n_2\} \quad (\text{G.1})$$

holds with probability at least $1 - O(n^{-10})$. For any fixed orthonormal matrices $\mathbf{U} \in \mathbb{R}^{n_1 \times r_1}$ and $\mathbf{V} \in \mathbb{R}^{n_2 \times r_2}$, we have

$$\|\mathbf{U}^\top \mathbf{E} \mathbf{V}\| \leq \frac{1}{2} C_g \sigma \sqrt{r_1 + r_2 + \log n} \quad (\text{G.2})$$

with probability at least $1 - O(n^{-10})$. As a consequence, for any $\mathbf{X} \in \mathbb{R}^{n_1 \times m_1}$ and $\mathbf{Y} \in \mathbb{R}^{n_2 \times m_2}$,

$$\|\mathbf{X}^\top \mathbf{E} \mathbf{Y}\| \leq \frac{1}{2} C_g \sigma \|\mathbf{X}\| \|\mathbf{Y}\| \sqrt{\text{rank}(\mathbf{X}) + \text{rank}(\mathbf{Y}) + \log n} \quad (\text{G.3})$$

holds with probability at least $1 - O(n^{-10})$.

Proof. The first result (G.1) is standard and is a special case of (G.2) by setting $\mathbf{U} = \mathbf{I}_{n_1}$ and $\mathbf{V} = \mathbf{I}_{n_2}$. To bound the spectral norm of $\mathbf{U}^\top \mathbf{E} \mathbf{V}$, we first write it as

$$\|\mathbf{U}^\top \mathbf{E} \mathbf{V}\| = \sup_{\mathbf{x} \in \mathcal{S}^{r_1}, \mathbf{y} \in \mathcal{S}^{r_2}} \mathbf{x}^\top \mathbf{U}^\top \mathbf{E} \mathbf{V} \mathbf{y} = \sup_{\mathbf{x} \in \mathcal{S}^{r_1}, \mathbf{y} \in \mathcal{S}^{r_2}} \sum_{i=1}^{n_1} \sum_{j=1}^{n_2} E_{i,j}(\mathbf{U}_i, \mathbf{x})(\mathbf{V}_j, \mathbf{y})$$

where $\mathcal{S}^r = \{\mathbf{x} \in \mathbb{R}^r : \|\mathbf{x}\|_2 = 1\}$ denotes the unit sphere in \mathbb{R}^r . By Hoeffding's inequality [Ver17, Theorem 2.6.3], we know that with probability exceeding $1 - \delta$,

$$\sum_{i=1}^{n_1} \sum_{j=1}^{n_2} E_{i,j}(\mathbf{U}_i, \mathbf{x})(\mathbf{V}_j, \mathbf{y}) \leq C_b \sigma \|\mathbf{U} \mathbf{x}\|_2 \|\mathbf{V} \mathbf{y}\|_2 \sqrt{\log(\delta^{-1})} = C_b \sigma \sqrt{\log(\delta^{-1})}$$

for some universal constant $C_b > 0$. Here the last relation holds since \mathbf{U} and \mathbf{V} both have orthonormal columns. The 1/4-covering number of \mathcal{S}^r is upper bounded by 9^r [Ver17, Corollary

4.2.13]. Let \mathcal{N}_{r_1} and \mathcal{N}_{r_2} be the $1/4$ -covering of \mathcal{S}^{r_1} and \mathcal{S}^{r_2} satisfying $|\mathcal{N}_{r_1}| \leq 9^{r_1}$ and $|\mathcal{N}_{r_2}| \leq 9^{r_2}$. Then with probability exceeding $1 - 9^{r_1+r_2}\delta$,

$$\|\mathbf{U}^\top \mathbf{E} \mathbf{V}\| \stackrel{(i)}{\leq} 2 \max_{\mathbf{x} \in \mathcal{N}_{r_1}, \mathbf{y} \in \mathcal{N}_{r_2}} \sum_{i=1}^{n_1} \sum_{j=1}^{n_2} E_{i,j}(\mathbf{U}_i, \cdot; \mathbf{x})(\mathbf{V}_j, \cdot; \mathbf{y}) \stackrel{(ii)}{\leq} 2C_b \sigma \sqrt{\log(\delta^{-1})}.$$

Here (i) is the standard technique for bounding the spectral norm using nets, see [Ver17, Section 4.4.1] for details; and (ii) follows from a union bound argument. By taking $\delta = n^{-10}9^{-r_1-r_2}$, we reach the desired bound (G.2). Finally it is straightforward to establish the last result (G.3) by considering the SVDs of \mathbf{X} and \mathbf{Y} and applying (G.2), and we omit the proof here for brevity. \square